

Oct. 17, 1975

3286-43

FOR ECONOMIC POISON REGISTERED
UNDER NO.

With Comments

CAUTION:

Harmful if swallowed. Avoid prolonged breathing of spray mist or dust. Avoid contact with skin. Wash thoroughly after using. Avoid contamination of feed and foodstuffs. Keep out of reach of children.

Note to Physician: Malathion, upon use may cause cholinesterase inhibition. Atropine is antidotal.

FLIES: For reduction of fly population, use 32 to 40 pounds per 100 gallons of water, in dairy barns, horse and sheep barns, poultry houses, corrals, manure piles, garbage heaps, city dumps, etc. Apply spray thoroughly to wet ceilings, walls and other places where flies congregate.

For a bait spray, use 32 pounds with 2 gallon unsulfurized molasses or corn syrup, or 20 pounds of sugar per 100 gallons of water. Repeat application as necessary.

NOTE: Do not apply to freshly whitewashed surfaces. Wait 14 days before using. Recommend continued agitation of mixtures to insure even dispersion through solution.

Ornamentals: Use 2½ pounds Malathion 25% Wettable Powder per 100 gallons of water for control of Aphids, Spider Mites, Whitefly, Mealybugs, Thrips, Japanese Beetle adults, Four-Lined Leaf Bug, Tarnished Plant Bug, Rose Leafhopper, Potato Leafhopper, Scurfy Scale, Euonymous Scale and European Pine Shoot Moth.

Make one or more applications as needed. Use 6 pounds Malathion 25% wettable Powder per 100 gallons of water for control of Soft Brown Scale, Monterey Pine Scale and Black Scale Crawlers. Make one or more applications as needed. Use 6 pounds Malathion 25% Wettable Powder per 100 gallons of water for control of Juniper Scale, Fletcher Scale, Bagworms, Oak Kermes, Florida Red Scale, Azalea Scale, Magnolia Scale, Pine Leaf Scale, Tent Caterpillar, Birch Leaf Miner and Boxwood Leaf Miner. Make one or more applications as needed after crawlers have settled on the foliage.

NOTE: Make thorough full coverage sprays. Repeat application as needed. Do not use on Boston, Maidenhair, Pteris Ferns or species of Crassula. Do not apply on Petunias.

Chicken Red Mites and Northern Fowl Mite in Poultry Houses: Apply 2½ oz. of 25% MALATHION WETTABLE per gal. per 100 to 150 birds. Repeat application in 4 to 8 weeks or when necessary. Do not apply Malathion directly to birds. Avoid contamination of food and water troughs.

Do not reuse container. Destroy when empty.

WARNING!

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Oct. 17, 1975

3286-43

10/17/75
WITH COMMENTS

CAUTION:

Harmful if swallowed. Avoid prolonged breathing of spray mist or dust. Avoid contact with skin. Wash thoroughly after using. Avoid contamination of feed and foodstuffs. Keep out of reach of children.

Note to Physician: Malathion, upon use may cause cholinesterase inhibition. Atropine is antidotal.

FLIES: For reduction of fly population, use 32 to 40 pounds per 100 gallons of water, in dairy barns, horse and sheep barns, poultry houses, corrals, manure piles, garbage heaps, city dumps, etc. Apply spray thoroughly to wet ceilings, walls and other places where flies congregate.

For a bait spray, use 32 pounds with 2 gallon unsulfurized molasses or corn syrup, or 20 pounds of sugar per 100 gallons of water. Repeat application as necessary.

NOTE: Do not apply to freshly whitewashed surfaces. Wait 14 days before using. Recommend continued agitation of mixtures to insure even dispersion through solution.

Ornamentals: Use 2½ pounds Malathion 25% Wettable Powder per 100 gallons of water for control of Aphids, Spider Mites, Whitefly, Mealybugs, Thrips, Japanese Beetle adults, Four-Lined Leaf Bug, Tarnished Plant Bug, Rose Leafhopper, Potato Leafhopper, Scurfy Scale, Euonymous Scale and European Pine Shoot Moth.

Make one or more applications as needed. Use 6 pounds Malathion 25% wettable Powder per 100 gallons of water for control of Soft Brown Scale, Monterey Pine Scale and Black Scale Crawlers. Make one or more applications as needed. Use 6 pounds Malathion 25% Wettable Powder per 100 gallons of water for control of Juniper Scale, Fletcher Scale, Bagworms, Oak Kermes, Florida Red Scale, Azalea Scale, Magnolia Scale, Pine Leaf Scale, Tent Caterpillar, Birch Leaf Miner and Boxwood Leaf Miner. Make one or more applications as needed after crawlers have settled on the foliage.

NOTE: Make thorough full coverage sprays. Repeat application as needed. Do not use on Boston, Maidenhair, Pteris Ferns or species of Crassula. Do not apply on Petunias.

Chicken Red Mites and Northern Fowl Mite in Poultry Houses: Apply 2½ oz. of 25% MALATHION WETTABLE per gal. per 100 to 150 birds. Repeat application in 4 to 8 weeks or when necessary. Do not apply Malathion directly to birds. Avoid contamination of food and water troughs.

Do not reuse container. Destroy when empty.

WARNING!

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.



MALATHION WETTABLE



CAUTION

Note to Physician

For reduction of fly population

For a bait spray

Ornamentals

WARNING!

NET WEIGHT 25 LBS

MALATHION 25% WETTABLE POWDER

WARNING!

KEEP OUT OF REACH OF CHILDREN!

See side panel for additional caution

Active Ingredient: 25%
Malathion* 25%
Inert Ingredients 75%
*O-O-dimethyl dithiophosphate of diethyl mercaptosuc-
cinate.

Ferd Staffel Co.

331 Burnet Street

San Antonio, Texas 78298

EPA Reg. No. 106-1066