





BIO DUST IS A BIOLOGICAL INSECTICIDE INTENDED FOR USE AGAINST CHEWING

JUL 27 1977

Futo dela us . registrate EPA Rec.



PRECAUTIONARY STATEMENT CAUTION **ENVIRONMENTAL HAZARDS**

LARVAE (CATERPILLARS) OF CERTAIN LEPIDOPTEROUS SPECIES.

Keep out of lakes, streams or ponds. Do not contaminate water by cleaning or by disposal of wastes.

DIRECTIONS FOR USE

GENERAL CLASSIFICATION: It is a violation of federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL DIRECTIONS

Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited. Dispose of in an incinerator or landfill approved for pesticide containers, or bury in a safe place.

BIQ DUST is recommended for use by ground or by air. Apply when pests are first noticed and repeat as necessary. Thorough plant coverage is necessary for control.

APPLICATION RATE

FOR the control of Cabbage Looper on Broccoli, Cabbage, Cauliflower, Collards, Celery, Lettuce, Melons, Potatoes, Spinach, Tomatoes and Cucumbers, apply 25 to 50 lbs. per acre. GROUND APPLICATION ONLY: Cabbage Looper on Kale, Mustard Greens and Turnip Greens apply at 25 to 50 pounds per acre.

FOR the control of Imported Cabbage Worm on Broccoli, Cabbage, Cauliflower, Collards and Turnip Greens, apply 15 to 25 lbs. per acre.

GROUND APPLICATION ONLY: imported Cabbageworm on Kale and Mustard Greens apply at 15 to 25 pounds per acre.

FOR the control of Hornworm on Tobacco, apply 12 to 25 lbs. per acre.

FERD STAFFEL COMPANY, San Antonio, TX 78298

ACTIVE INGREDIENTS:

Bacillus thuringiensis, Berliner, Potency of 320 International Units per mg. (at least *Equivalent to potency of 0.15 billion international Units per pound of this product.

CAUTION

Keep Out Of Reach Of Children. See back panel for additional cautions. Read All Directions Before Using.

E.P.A. Reg. No. 3286-46 E.P.A. Est. No. 3286-TX-1 Net Weight 1 lb.