



# <Staffel's><sup>TM</sup>

*There is no Alternate*  
**SPECIAL**

# lawn food

**15-10-10**

FS2-45A

## FERTILIZER - INSECTICIDE

Total Nitrogen	15%
Available Phosphoric Acid (P <sub>2</sub> O <sub>5</sub> )	10%
Soluble Potash (K <sub>2</sub> O)	10%
Sulfur (S)	13%
Iron (Fe)	0.35%
Zinc (Zn)	0.27%

CONTAINS DURSBAN\*

FOR THE CONTROL OF SOIL INSECTS SUCH AS ANTS, ARMYWORMS, CUTWORMS, SOD WEBWORMS, CHINCH BUGS, EARWIGS, SOWBUGS, CRICKETS, AND CERTAIN WHITE GRUBS

ACTIVE INGREDIENT  
Chlorpyrifos (O,O-diethyl O-(3,5,6-Trichloro-2-pyridyl)  
phosphorothioate  
INERT INGREDIENTS

0.3  
99.7  
TOTAL 100.0

DURSBAN® Reg. Trademark of the Dow Chemical Company  
U.S. Patent No. 3,244,586

KEEP OUT OF REACH OF CHILDREN  
**CAUTION**  
READ ALL DIRECTIONS BEFORE USING  
SEE BACK PANEL FOR ADDITIONAL CAUTIONS

EPA Reg. No. 3286-50

EPA Est. No. 3286 TX 1

**NET WT. 50 LBS. (22.7 kg)**

MANUFACTURED BY  
THE STAFFEL COMPANY, San Antonio, TX 78298



*'There is no Alternate'*

#### PRECAUTIONARY STATEMENTS

##### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION** May be harmful if swallowed. Avoid eye contact. Wash thoroughly after handling. Wash contaminated clothing before reuse. Flush contaminated eyes with plenty of water and get medical attention if irritation persists. Keep away from food, feedstuffs and domestic water supplies. Keep children and pets off of treated area until watered in, and the area is dry.

**NOTE TO PHYSICIAN:** Active ingredient is a cholinesterase inhibitor. Treat symptomatically. Atropine is an antidote.

##### ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. Keep out of any body of water. Fish and crustaceans may be killed if their waters are contaminated with this product. Do not contaminate water by cleaning equipment or disposing of wastes.

#### DIRECTIONS FOR USE

##### GENERAL CLASSIFICATION

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

##### DISPOSAL DIRECTIONS

Dispose of contents and container in accordance with applicable regulations. Do not reuse or refill this container.

**NOTICE:** Seller's quantity shall be limited to the terms of the label and the user. Buyer assumes any risk to person or property, and to the environment, in using this product in these conditions.

#### HOW TO USE

To control chinch bugs, sod webworms, ants, armyworms, crickets, cutworms, earwigs, and sowbugs, uniformly apply the contents of this 50 lb. bag to cover 6500 sq. ft. of noncrop turfgrass area using any hand- or power-operated gravity or rotary type spreader designed for application of granular materials. Treat when insects first appear; re-treat as needed. To help obtain uniform distribution, apply one-half the required amount of product in one direction; then apply the balance at a right angle. *Immediately after application* thoroughly water the treated turf to wash the insecticide into the zone of insect infestation.

To control certain white grubs (larvae of European chafer and Japanese beetle) use 25 to 50 pounds of this product per 1600 sq. ft. Apply the granules uniformly over the area treated followed immediately by thorough watering to wash the insecticide into the turf. For best results treat when grubs are young and actively feeding near the soil surface, usually during late July and August. Consult your local Agricultural Extension Service specialist for proper time to treat in your area.

##### SPREADER SETTINGS

	Heavy
Cyclone	6-6 1/2
Ortho Drop Spreader	6
Ortho Whirlbird	4
Scotts	3