

**GENERAL DIRECTIONS FOR USE** — Place bait in dry containers next to burrows, in runways, or in locations where rats and mice are accustomed to feeding, drinking or traveling. Desirable bait placements are along walls, in corners, and in concealed locations where rats and mice can feed unobserved. Replace stale, damp or moldy bait with fresh bait if necessary. Continue baiting until all signs of feeding have stopped.

**RATS** - Use 4 or more ounces of bait at each location and maintain an uninterrupted supply of bait for at least 10 days.

**MICE** - Place tablespoon amounts 1/4 to 1/2 ounce of bait in shallow containers at 8-10 foot intervals and maintain an uninterrupted supply of bait for at least 15 days.

  
*Colonial* 42  
**KILLS**  
**RATS and MICE**

NET WT. lbs.

**WARFARIN**

**ACTIVE INGREDIENT:** Warfarin

3-alpha-acetonyl-benzyl 4-hydroxycoumarin	0.025%
<b>INERT INGREDIENTS</b>	99.975%
	<hr/>
	100.000%

**COLONIAL PRODUCTS, INC.**

1830 — 10th AVE. No.

LAKE WORTH, FLA.

**CAUTION:** KEEP OUT OF REACH OF CHILDREN  
SEE SIDE PANEL FOR ADDITIONAL CAUTIONS

**CAUTION** — Protect children, pets and domestic animals from bait. If swallowed by humans, animals or pets, this material may reduce the clotting ability of the blood and cause bleeding. In such cases, a physician or veterinarian should be called immediately. Intravenous and oral administration of vitamin K combined with blood transfusions are indicated as in the case of hemorrhage caused by overdoses of bishydroxycoumarin.

This rodenticide contains Warfarin, an anticoagulant chemical that reduces the clotting ability of the blood and, upon repeated feedings, causes fatal hemorrhage in rats and mice.

U.S.D.A. REG. NO. 3314-1

