


COLONIAL  
**PES-TA-REST**

E.P.A. Reg. No. 3314-29  
E.P.A. Est. No. 3314-F1-1

Active Ingredients

Terpyridine	0.001%
Orthophenylphenol (Oxydiazole)	0.006%
Related ingredients	0.001%
Other Aromatic hydrocarbons	0.001%
Related compounds	0.001%
Aromatic petroleum hydrocarbons	99.338%
Inert Ingredients	100.000% w/w

Manufactured by  
Colonial Products, Inc.  
1830 Tenth Avenue, P.O. Box 3200  
Lake Worth, Florida 33458



**DIRECTIONS FOR USE**

Shake well before each use. Remove protective cap. Aim spray, keeping away from persons. Push button to spray.

Foods should be removed or covered during treatment. A person should not eat food treated or recovered during treatment or thoroughly cleaned before eating. When using this product in these areas, apply only when the facility is not in operation.

**CONTROL HOUSEHOLD PESTS**

**For Houseflies, Mosquitoes and Gnats** — Spray directly onto the walls, windows, doors, window frames, and base paths of the entire residence. Wait until the spray has dried completely before re-entering the room. For the same area, the spray may be repeated every two hours. For gnats, spray around the growing insects such as flowers, shrubs, trees, etc., and allow to dry.

**Ants** — Direct spray onto the trails and nests of the ants. Repeat as necessary.


**OUTDOORS — TO REDUCE ANNOYANCE FROM FLYING INSECTS**

For the reduction of flying insects such as gnats, mosquitoes, houseflies, etc., spray directly onto the foliage of the plants. Hold container 10 inches from the plants. Spray in short bursts of one to three seconds or spray at the rate of one linear foot of growth per second. Direct spray to all sides of the foliage until slightly moist. Avoid heavy wetting. Repeat as necessary.

**OUTDOORS — TO PROTECT ORNAMENTAL PLANTS**

To control aphids, exposed thrips, adult white flies, leafhoppers, two-spotted mites, plant bugs, spittlebugs and Japanese beetles on smaller rhododendrons, pines, spruce, hemlock, spirea, dogwood, palm, mimosa, toyon, viburnum, euonymus, oaks, maples, locust, forsythia and althea; and mimosa webworm larvae on mimosa, spray directly on insects. Hold container 10 inches from the infected plants. Spray in short spray bursts of one to three seconds or spray at the rate of one linear foot of growth per second. Direct spray to all sides covering foliage until slightly moist. Avoid heavy wetting. Repeat as necessary to obtain control.

COLONIAL  
**PES-TA-REST**



Manufactured by  
Colonial Products, Inc.  
230 Tenth Avenue, North  
Orlando, Florida 33460

Cis/trans isomers ratio: max. 30% ( $\pm$ ) cis and min. 70% ( $\pm$ ) trans.  
U.S. Patent No. 3,465,007.

**SHAKE WELL BEFORE USING.**

**CAUTION**  
KEEP OUT OF REACH OF CHILDREN

DO NOT SPRAY ON PLANTS OR FLOWERS IN BLOOM. DO  
NOT SPRAY ON PLANTS OR FLOWERS IN BLOOM. DO  
NOT SPRAY ON PLANTS OR FLOWERS IN BLOOM. DO  
NOT SPRAY ON PLANTS OR FLOWERS IN BLOOM.

**DO BODS — IN FLUID FORM FOR PLANTS**

Spray peonies, iris, zinnia, snapdragons, ageratum, chrysanthemums, nasturtiums, daisy, geraniums, marigolds, daphne, fothergilla, ivy, lobelia, Queen Anne's lace and broad leaf acacia for control of aphids, especially thrips, adult white flies, leafhoppers, plant bugs, spittlebugs and Japanese beetles. Spray as directed above.

USE WITH CAUTION on azaleas, junipers, roses, including hybrid tea roses, flowering quince, ash and camellia. These plants may be subject to leaf or flower burn under certain conditions of use.

For adult white flies, two-spotted mites and mealybugs, be sure to direct spray to contact these pests. Thoroughly cover the infested area. Use several repeat applications at two day intervals if necessary.

**PROTECTIVE PLANTS:**

Do not spray on hollyhocks, phlox, geraniums, begonias, coleus for rapid pests. Leave the leaves dry until dark hours, and when cool after treatment is completed.

**MISCELLANEOUS PLANTS — ON THE EXTERIOR BUILDINGS:**

When spraying on trees, shrubs, vines, etc., when leaves are dry, spray liberally. Do not spray on plants which are in bloom. Spray liberally. Do not spray on plants which are in bloom. Spray liberally.

When spraying on trees, shrubs, vines, etc., when leaves are dry, spray liberally. Spray liberally. Do not spray on plants which are in bloom. Spray liberally.

When spraying on trees, shrubs, vines, etc., when leaves are dry, spray liberally.

When spraying on trees, shrubs, vines, etc., when leaves are dry, spray liberally.

HARMFUL IF SWALLOWED. Avoid ingestion of food and household. Remove birds, pets and cover aquaria with glass. Do not leave in direct sunlight.