

The 1.5 liter canister can treat an undivided area up to 12,000 cubic feet (15' x 40' x 10' ceiling). DO NOT USE IN AN AREA SMALLER THAN 1,000 CUBIC FEET (15' x 20' x 10' ceiling). For smaller rooms, use a CAN-1 quart, glass, 7.5 ounce spray. Treats room up to 6,000 cubic feet (20' x 30' x 10' ceiling).

DISINFECTION

CLOSE all windows, when treated floors and treated furniture which may be present with tape or rags. (Cleaning pads may be used.)

PUT away papers, documents, framed and unframed art in areas to be treated.

REMOVE pots and covers or remove fish tanks and bowls; REMOVE aprons, towels, linens, etc. and articles made of cloth or thermocloth.

CLOSE all doors, doors and windows; turn off fans and air conditioners. Turn on lights, all open flamed except water heater.

APPENDIX NO.

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

1. Set can on a surface open, such as a table, chair or stand at least 12 inches from outlet light. Spread paper dinner cup to cover area directly below the nozzle. Remove cover.

2. Tilt top of can away from you.	3. Press lever down firmly until it clicks.	4. Set can upright on paper...	5. ...and leave the room closing the door behind you.
-----------------------------------	---	--------------------------------	---

6. Leave room being treated closed for 4 hours.

7. After 4 hours, open all doors and windows, turn on air conditioner and fans, and let the treated area air for 30 minutes.

Re-treat "Treatment in two weeks or as necessary".

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Avoid inhalation of vapors. Avoid contact with skin, eyes, or clothing. Wash contaminated skin promptly with soap and warm water. For eyes, flush with plenty of water. Get medical attention if irritation persists. Avoid contamination of food and food utensils. Do not used in commercial food processing, preparation, service or storage areas. In the home, cover or remove exposed food. Avoid contamination of cooking utensils and food preparation surfaces. Stay out of rooms being treated for 4 hours, ventilate the rooms as directed.

PHYSICAL AND CHEMICAL HAZARDS

Contents under pressure. Do not use or store near heat or open flame. Do not puncture or incinerate container. Exposure to temperatures above 120°F may cause bursting.

DISPOSAL

1. Remove product from can. Wrap canister and put in trash collection.

THE RICOH COMPANY, INC., HUNTSVILLE,

100 University Street, Huntsville, Alabama, U.S.A. 35801
DIA Reg. No. 11715-CT-1 EPA Reg. No. 11598-CT-1, DIA, Huntsville, AL

(See code on bottom) 11715-CT-1, 11598-CT-1

EPA-PR

GENERAL DIRECTIONS

Shake well before each use. Remove protective cap away from person. Push button to spray.

Foods should be removed or covered during treatment. Surfaces should be covered during treatment before using. When using the product in these areas the facility is not in operation.

CONTROL HOUSEHOLD PESTS

For Houseflies, Mosquitoes, and Gnats: and windows. Point nozzle upward, direct spray in room, especially windows and other light sources. Sectors. Fill the room with mist, then leave the room closed for at least 15 minutes. Ventilate the room when mist has disappeared.

For Crawling Insects: Roaches, earwigs, spiders and centipedes. Spray thoroughly into cracks, crevices, moist areas, openings around refrigerators, baseboards, and storage areas. Spray where possible.

Ants: Spray trails, nests and points of entry. Spray as possible. Repeat as necessary.

Animal Pests: Dogs, Cats - Fleas. Apply directly under forelegs and on other areas. Part hair to get good coverage. Hold nozzle about one foot. Apply until hair is thoroughly moist. Do not directly on mouth of animal. To reduce reinfestation, spray animal's bedding and other resting areas. Treat at rate of two seconds per linear foot while spraying at a distance. Also spray entire inside surface of dog kennels. Repeat as necessary.

OUTDOORS — TO REDUCE A NUISANCE FROM FLYING INSECTS

Reduce mosquitoes, gnats, midges, houseflies, and nocturnal flying moths by spraying their resting sites listed above. Hold container 18 to 24 inches from grassy areas. Spray these areas with a slow spray rate of from 1 to 3 seconds per cubic yard of space from the treated areas. To reduce annoyance from these open porches and patios in a similar manner air is still. Allow spray to dissipate before occupying. Repeat as necessary.

Physical and Chemical H

Contents under pressure. Do not puncture. Store near heat or open flame. Never throw can or incinerator. Exposure to temperatures above cause bursting.

COLONIAL PES-TA-REST

FLEA FLIP[®] DOG & CAT SPRAY



Active Ingredients

Resmethrin (5-Ethenyl-3-methyl-2-isopropenyl-3-(2-methyl-1-propenyl)-cyclopentene-1-carboxylate)	0.200% w/w
Related Compounds	0.028%
Dimethyl Aluminim Acetate (Acetyl Dimethyl Aluminim)	0.150%
Related Compounds	0.012%
Alcohol Petroleum Hydrocarbons	0.272%
Isobutane	.99.338%
Total	100.000% w/w

Inert Ingredients

KEEP OUT OF REACH OF CHILDREN CAUTION

Precautionary Statement

Hazards to Humans and Domestic Animals

Avoid contact with eyes, skin or clothing.

Avoid contamination of feed and foodstuffs.

Avoid breathing.

NET WT 16 OZ

EPA Reg. No. 3514-29 EPA Est. No. 1-623-58-1

Manufactured for



COLONIAL PRODUCTS, INC.
1830 Tenth Avenue North
Lake Worth, Florida 33461

A

FL.

Directions for Use

For use on food processing equipment and ornamental plants.
With or after cleaning.

Storage and Disposal

Keep out of reach of children.
Do not contaminate water supplies.

**OUTDOORS — TO PROTECT
ORNAMENTAL PLANTS**

To control gnats, aphids, plant lice, white flies, leafhoppers, fall-saucer scale, soft scales and lace bugs on smaller shrubs, vines, roses, lilacs, holly, azaleas, dogwood, palm trees, rosemary, yucca, agave, citrus, fruit trees, fruit, banana, caladium, and many other warm weather annuals. Spray directly on stems. Hold container 18 inches from the treated plants. Spray in early morning or late evening. Reapply at 10 to 14 days intervals depending on growth per season. Treat so as to avoid overspray covering leaves until slightly moist. Avoid heavy spraying. Repeat as necessary to maintain control.

Day Plants: Use 1/2 to 1 oz. per square yard. Use 1/2 oz. per square foot for trees, shrubs, and ground covers. Use 1/2 oz. per square foot for flowers, vines, and other annuals. Use 1/2 oz. per square foot for tropicals and palms.

USE WITH CAUTION: Do not spray on blossoms, fruits, seed pods, or ripening vegetables. Do not spray on young seedlings. Do not spray on flowers, vines, or other plants that are to be eaten raw. Do not spray on lawns, grass, or other areas where people walk.

DO NOT SPRAY ON CHILDREN OR PETS: Do not spray on children or pets. Do not spray on dogs, cats, or other animals. Do not spray on birds, bees, butterflies, or other insects.

PROTECT HOUSE PLANTS:

DO NOT SPRAY ON CHILDREN OR PETS: Do not spray on children or pets. Do not spray on dogs, cats, or other animals. Do not spray on birds, bees, butterflies, or other insects.

**NUISANCE PESTS — ON OR
ENTERING BUILDINGS:**

Wasps: Apply direct to nests, larvae, and larvae. Do not spray on flying insects. Do not spray on flowers, shrubs, trees, or other plants. Do not spray on birds, bees, butterflies, or other insects.

Ants: Use 1/2 oz. per square foot. Use 1/2 oz. per square foot for lawns, gardens, and other areas.

Spiders: Use 1/2 oz. per square foot. Use 1/2 oz. per square foot for lawns, gardens, and other areas.

Earwigs: Use 1/2 oz. per square foot. Use 1/2 oz. per square foot for lawns, gardens, and other areas.

3314-191

COLONIAL PES-TA-REST**GNAT GNIP****Active Ingredients**

Resmethrin 1% Phenylmethoxy-3,4-dimethyl-2,2-dimethyl-3,2-dimethyl-1-propenyl cyclopropane carbamate	0.20%
Related Compounds	0.08%
3,4-Dihydroxy-2-methyl-1-propanol	0.15%
Related Compounds	0.22%
Aromatic petroleum hydrocarbons	0.22%
Inert Ingredients	94.36%

KEEP OUT OF REACH OF CHILDREN**CAUTION****Precautionary Statement****Hazards to Humans and Domestic Animals**

Avoid contact with eyes, skin or clothing.

Avoid contamination of feed and foodstuffs.

Avoid breathing vapors.

NET WT 1/2 LITER

E.P.A. Reg. No. 3314-24 E.P.A. Est. No. 1162363-1

Manufactured for

COLONIAL PRODUCTS, INC.
1830 Tenth Avenue North
Lake Worth, Florida 33461

CYAN-PB

12.000

GENERAL DIRECTIONS

Shake well before each use. Remove protective cap. Aim spray opening away from person. Push button to spray.

Foods should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using. When using the product in these areas, apply only when the facility is not in operation.

CONTROL HOUSEHOLD PESTS:

For Houseflies, Mosquitoes, and Gnats: Close all doors and windows. Point nozzle upward, direct spray mist to all parts of the room, especially windows and other light sources which attract these insects. Fill the room with mist, then leave the treated area. Keep the room closed for at least 15 minutes. Ventilate the room when treatment is completed.

For Crawling Insects: Roaches, earwigs, green bugs, sowbugs, spiders and centipedes. Spray thoroughly into hiding places such as cracks, crevices, moist areas, openings around pipes and sinks, under refrigerators, baseboards and storage areas. Spray directly on insects where possible.

Ants: Spray trails, nests and points of entry. Spray on ants where possible. Repeat as necessary.

Animal Pests: Dogs, Cats — Fleas. Apply directly to animals along back under forelegs and on other areas. Part hair to contact skin and insure good coverage. Hold nozzle about one foot away while spraying. Apply until hair is thoroughly moist. Do not directly spray into eyes, ears or mouth of animal. To reduce reinestation thoroughly spray the animal's bedding and other resting areas. Treat these areas at the rate of two seconds per linear foot while spraying at a distance of 12 to 18 inches. Also spray entire inside surface of dog houses. Repeat as necessary.

**OUTDOORS — TO REDUCE ANNOYANCE
FROM FLYING INSECTS:**

Reduce mosquitoes, gnats, midges, houseflies, biting flies and pyralid and noctuid flying moths by spraying their resting areas. Follow precautions listed above. Hold container 18 to 24 inches from shrubs, bushes and grassy areas. Spray these areas with a slow sweeping motion at the rate of from 1 to 3 seconds per cubic yard of space while moving away from the treated areas. To reduce annoyance from these flying insects treat open porches and patios in a similar manner. Spray only when the air is still. Allow spray to dissipate before occupying the treated area. Repeat as necessary.

Physical and Chemical Hazards

Contents under pressure. Do not puncture. Do not use or store near heat or open flame. Never throw container into fire or incinerator. Exposure to temperatures above 130°F may cause bursting.

COLONIAL

PES-TA-REST

FLY FLIP

Directions for Use

Storage and Disposal

OUTDOORS — TO PROTECT ORNAMENTAL PLANTS

For ornamental plants and other outdoor areas, spray mist directly onto foliage, flowers and stems. Do not spray on blossoms or fruit. Do not spray on plants which are sensitive to insecticides. Do not spray on plants which are sensitive to insecticides. Do not spray on plants which are sensitive to insecticides.

USE WITH CAUTION

Do not spray on lawns, trees, shrubs, vines, flowers, fruit, vegetables, or other plants which are sensitive to insecticides. Do not spray on plants which are sensitive to insecticides. Do not spray on plants which are sensitive to insecticides.

PROTECT HOUSE PLANTS

Do not spray on houseplants. Do not spray on plants which are sensitive to insecticides. Do not spray on plants which are sensitive to insecticides.

NUISANCE PESTS — ON OR ENTERING BUILDINGS

Wasps: Spray mist directly onto wasps and nests. Do not spray on birds, pets or people. Do not spray on birds, pets or people.

Ants: Spray mist directly onto ant colonies and nests. Do not spray on birds, pets or people. Do not spray on birds, pets or people.

Spiders: Spray mist directly onto spiders and webs. Do not spray on birds, pets or people. Do not spray on birds, pets or people.

Earwigs: Spray mist directly onto earwigs and nests. Do not spray on birds, pets or people. Do not spray on birds, pets or people.

COLONIAL PES-TA-REST

FLY FLIP



Active Ingredients

Residual Insecticide: 2.5% Pyrethrin, 0.25% Pirethrin, 0.25% Cypermethrin	0.0075%
Residual Insecticide: 0.25% Resmethrin	0.0025%
Other Active Ingredients: 0.0005%	0.0005%
Related Compounds: 0.012%	0.012%
A small amount of inert materials	0.272%
Inert Ingredients:	0.318%
	1.000%

KEEP OUT OF REACH OF CHILDREN CAUTION

PRECAUTIONARY STATEMENT

HAZARDS TO Humans and Domestic Animals

Avoid contact with eyes, skin or clothing.

Avoid inhalation of feed and foodstuffs.

Avoid fire and explosion.

NOTIFICATION

REGULATORY INFORMATION

MANUFACTURER


COLONIAL PRODUCTS, INC.
1851 North Avenue North
Lake Worth, Florida 33461

GENERAL DIRECTIONS

Shake well before each use. Remove protective cap. Aim spray opening 3-way from person. Push button to spray.

Foods should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using. When using the product in these areas, apply only when the facility is not in operation.

CONTROL HOUSEHOLD PESTS:

For Household, Mosquitoes, and Gnats: Close all doors and windows. Point nozzle upward; direct spray mist to all parts of the room, especially windows and other light sources which attract these insects. Fill the room with mist, then leave the treated area. Keep the room closed for at least 15 minutes. Ventilate the room when treatment is completed.

For Crawling Insects: Roaches, earwigs, grain mites, sowbugs, spiders and centipedes. Spray thoroughly into hiding places such as cracks, crevices, moist areas, openings around pipes and sinks, under refrigerators, baseboards, and storage areas. Spray directly on insects where possible.

Ants: Spray basis nests and points of entry. Spray on ants where possible. Repeat as necessary.

Animal Pests: Dogs, Cats — Fleas. Apply directly to animals along back, under forelegs and on other areas. Part hair to contact skin and insure good coverage. Hold nozzle about one foot away while spraying. Apply until hair is thoroughly moist. Do not directly spray into eyes, ears or mouth of animal. To reduce reinfestation thoroughly spray the animal's bedding and other resting areas. Treat these areas at the rate of two seconds per linear foot while spraying at a distance of 12 to 18 inches. Also spray entire inside surface of dog houses. Repeat as necessary.

OUTDOORS — TO REDUCE ANNOYANCE FROM FLYING INSECTS:

Reduce mosquitoes, gnats, midges, houseflies, biting flies and pygmy and common flying moths by spraying their resting areas. Follow previous listed above. Hold nozzle 12 to 14 inches from shrubs, bushes and grassy areas. Spray these areas with a slow sweeping motion at the rate of four to six seconds per linear yard of space when moving away from the treated areas. To reduce annoyance from these flying insects, heat oven, porch and patio in a similar manner. Spray only when the insects are active. Allow spray to dissipate before occupying the treated area. Repeat as necessary.

Physical and Chemical Hazards

Contents under pressure. Do not puncture. Do not use or store near heat or open flame. Never throw container into fire or incinerator. Exposure to temperatures above 130°F may cause bursting.

13574-129