

3314-29

Directions for Use

For a complete Federal Insecticide, Fungicide, and Rodenticide Act registration number, see the label on the container.

Storage and Disposal

Keep this and all other pesticides in their original containers. Do not reuse empty containers. Do not mix with other pesticides.

OUTDOORS — TO PROTECT ORNAMENTAL PLANTS

For control of aphids, exposed thrips, adult white flies, leafhoppers, two-spotted spider mites, plant bugs, sunnyside bugs, Japanese beetles, scale insects, cicadas, grasshoppers, spruce bark beetles, dogwood borer, chrysomelids, leaf-miners, leaf beetles, sawflies, scarab beetles, leaf beetles, and other insects, and minute wet worm larvae on ornamentals, spray directly on insects in a closed container 18 inches from the infested plants. Spray in short spray bursts of one to three seconds at spray at the rate of one linear foot of growth per second. Direct spray to all sides covering foliage until it is lightly moist. Avoid heavy wetting. Repeat as necessary to obtain control.

For control of aphids, exposed thrips, adult white flies, leafhoppers, two-spotted spider mites, plant bugs, sunnyside bugs, Japanese beetles, scale insects, cicadas, grasshoppers, spruce bark beetles, dogwood borer, chrysomelids, leaf-miners, leaf beetles, sawflies, scarab beetles, leaf beetles, and other insects, and minute wet worm larvae on ornamentals, spray directly on insects in a closed container 18 inches from the infested plants. Spray in short spray bursts of one to three seconds at spray at the rate of one linear foot of growth per second. Direct spray to all sides covering foliage until it is lightly moist. Avoid heavy wetting. Repeat as necessary to obtain control.

USE WITH CAUTION — This product is extremely toxic to fish and aquatic life. Do not use in or near water. Do not use in or near swimming pools, ponds, lakes, streams, rivers, or other bodies of water. Do not use in or near fish hatcheries, nurseries, or other aquatic facilities. Do not use in or near areas where water is used for drinking or irrigation.

Do not use on plants that are sensitive to this product. Do not use on plants that are stressed or diseased. Do not use on plants that are in bloom. Do not use on plants that are in fruit. Do not use on plants that are in seed.

PROTECT HOUSE PLANTS:

For control of aphids, exposed thrips, adult white flies, leafhoppers, two-spotted spider mites, plant bugs, sunnyside bugs, Japanese beetles, scale insects, cicadas, grasshoppers, spruce bark beetles, dogwood borer, chrysomelids, leaf-miners, leaf beetles, sawflies, scarab beetles, leaf beetles, and other insects, and minute wet worm larvae on ornamentals, spray directly on insects in a closed container 18 inches from the infested plants. Spray in short spray bursts of one to three seconds at spray at the rate of one linear foot of growth per second. Direct spray to all sides covering foliage until it is lightly moist. Avoid heavy wetting. Repeat as necessary to obtain control.

NUISANCE PESTS — ON OR ENTERING BUILDINGS:

Wasps — Accumulations should be removed from eaves, porches, and other areas. Accumulations should be removed from eaves, porches, and other areas. Accumulations should be removed from eaves, porches, and other areas. Accumulations should be removed from eaves, porches, and other areas.

Ants — Accumulations should be removed from eaves, porches, and other areas. Accumulations should be removed from eaves, porches, and other areas. Accumulations should be removed from eaves, porches, and other areas.

Spiders — Accumulations should be removed from eaves, porches, and other areas. Accumulations should be removed from eaves, porches, and other areas. Accumulations should be removed from eaves, porches, and other areas.

Earwigs — Accumulations should be removed from eaves, porches, and other areas. Accumulations should be removed from eaves, porches, and other areas. Accumulations should be removed from eaves, porches, and other areas.

COLONIAL PES-TA-REST

PLANT PROTECTOR



Active Ingredients

Resmethrin (0.1% w/v), Deltamethrin (0.1% w/v), Cyfluthrin (0.1% w/v), Permethrin (0.1% w/v), Deltamethrin (0.1% w/v), Cyfluthrin (0.1% w/v), Permethrin (0.1% w/v), Deltamethrin (0.1% w/v), Cyfluthrin (0.1% w/v), Permethrin (0.1% w/v).

Inert Ingredients

KEEP OUT OF REACH OF CHILDREN CAUTION

Precautionary Statement: Hazards to Humans and Domestic Animals

Avoid contact with eyes, skin, or clothing. Avoid contamination of feed and feed stuffs. Avoid breathing vapors.

Net Weight 1.12 lb.

EPA Reg. No. 1014-01-01 EEA-Reg. No. 112-01-01

Manufactured by



COLONIAL PRODUCTS, INC.
1810 North Avenue North
Lake Worth, Florida 33461

EPA 48

12-001

GENERAL DIRECTIONS

Shake well before each use. Remove protective cap. Aim spray opening away from person. Push button to spray.

Funks should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using. When using the product in these areas, apply only when the facility is not in operation.

CONTROL HOUSEHOLD PESTS:

For Houseflies, Mosquitoes, and Gnats: Close all doors and windows. Point nozzle upward. Direct spray mist to all parts of the room, especially windows and other light sources which attract these insects. Fill the room with mist, then leave the treated area. Keep the room closed for at least 15 minutes. Ventilate the room when treatment is completed.

For Crawling Insects: Roaches, earwigs, grain mites, sawbugs, spiders, and centipedes. Spray thoroughly into hiding places such as cracks, crevices, moist areas, openings around pipes and sinks, under refrigerators, baseboards, and storage areas. Spray directly on insects where possible.

Ants: Spray trails, nests and points of entry. Spray on ants where possible. Repeat as necessary.

Animal Pests: Dogs, Cats, Fleas. Apply directly to animals along back under forelegs and on other areas. Part hair to contact skin and insure good coverage. Hold nozzle about one foot away while spraying. Apply until hair is thoroughly moist. Do not directly spray into eyes, ears or mouth of animal. To reduce reinfestation, thoroughly spray the animal's bedding and other resting areas. Treat these areas at the rate of two seconds per linear foot while spraying at a distance of 12 to 18 inches. Also spray entire inside surface of dog houses. Repeat as necessary.

OUTDOORS — TO REDUCE ANNOYANCE FROM FLYING INSECTS:

Reduce mosquitoes, gnats, midges, houseflies, biting flies and pyralid and noctuid flying moths by spraying their resting areas. Follow precautions listed above. Hold container 18 to 24 inches from shrubs, bushes and grassy areas. Spray these areas with a slow sweeping motion at the rate of from 1 to 3 seconds per cubic yard of space while moving away from the treated areas. To reduce annoyance from these flying insects, treat open porches and patios in a similar manner. Spray only when the insects stir. Allow spray to dissipate before occupying the treated area. Repeat as necessary.

Physical and Chemical Hazards

Contents under pressure. Do not puncture. Do not use or store near heat or open flame. Never throw container into fire or incinerator. Exposure to temperatures above 130°F may cause bursting.