

E.P.A. Est. No. 3314-FL-1
E.P.A. Reg. No. 3314-31

COLONIAL P.T.O. INDUSTRIAL RESIDUAL SPRAY

For use in and around the home, schools, theaters, warehouses,
motels, hotels, and other industrial establishments.
Do not use as a space spray.

ACTIVE INGREDIENTS:

Pyrethrins	0.05%
*Piperonyl butoxide, technical	0.05%
†N-octyl bicycloheptene dicarboximide	0.100%
**2-(1-methylethoxy) phenol methylcarbamate	1.000%
Petroleum distillate	80.684%

INERT INGREDIENTS

18.000%
100.000%

ACCEPTED
SEP 18 1975
U.S. DEPT. OF AGRICULTURE
FEDERAL INSECTICIDE
AND RODENT
RECORDS
RECORDED
INDEXED
3314-31

DIRECTIONS FOR USE

GENERAL: Spray only surfaces. Spray to wetting. Point spray away from people, plants and pets.


INDOOR USE: For control of ants, cockroaches (palmetto bugs, water bugs), crickets, silverfish, spiders, scorpions, saw-tooth grain beetles (exposed stages), spray into cracks, crevices, around baseboards and other places where these insects hide. Apply behind and beneath cabinets, refrigerators, sinks, stoves, and in and around waste containers. Repeat as needed. Do not spray foodstuffs or house plants. For ants, spray trails and places where they enter the building. Do not allow children to contact treated surfaces until surfaces are completely dry.

OUTDOOR USE: For control of ants, clover mites, crickets, earwigs, millipedes, sowbugs, and brown dog ticks, spray infested areas thoroughly including outside surfaces of screens, doors, window frames, foundations, patios or other places insects may use to enter the building. For ants, spray hills and runways. Repeat as necessary. Do not spray vegetation.

BROWN DOG TICKS: Spray sleeping quarters of pets, around baseboards, window and door frames, wall cracks and local areas of floors. Repeat as needed. Fresh bedding should be placed in animal quarters after spraying. Do not spray animals.

CAUTION

May be harmful if swallowed, inhaled or absorbed through the skin. Avoid breathing of spray mist and provide adequate ventilation of area being treated. Contact with skin, eyes, or clothing should also be avoided. Wash thoroughly with soap and warm water after handling. Avoid contamination of food, utensils and food preparation areas.



Manufactured by
Colonial Products, Inc.
1830 Tenth Avenue, North
Lake Worth, Florida 33460

EPA Est. No. 3314-FL-1
EPA Reg. No. 3314-31

COLONIAL P.T.O. INDUSTRIAL RESIDUAL SPRAY

For use in and around the home, schools, theaters, warehouses,
motels, hotels, and other industrial establishments.
Do not use as a space spray.

ACTIVE INGREDIENTS:

Pyrethrins	0.000%
*Piperonyl butoxide, technical	0.000%
†N-octyl bicycloheptene dicarboximide	0.100%
**2-(1-methylethoxy) phenol methylcarbamate	1.000%
Petroleum distillate	80.684%
INERT INGREDIENTS	18.000%
	100.000%

ACCEPTED
SEP 18 1975
Under the Federal Insecticide, Fungicide, and Rodenticide Act, this product is registered for the control of insects under EPA Reg. No. 3314-31

DIRECTIONS FOR USE

GENERAL: Spray only surfaces. Spray to wetting. Point spray away from people, plants and pets.

INDOOR USE: For control of ants, cockroaches (palmetto bugs, water bugs), crickets, silverfish, spiders, scorpions, saw-tooth grain beetles (exposed stages), spray into cracks, crevices, around baseboards and other places where these insects hide. Apply behind and beneath cabinets, refrigerators, sinks, stoves, and in and around waste containers. Repeat as needed. Do not spray foodstuffs or house plants. For ants, spray trails and places where they enter the building. Do not allow children to contact treated surfaces until surfaces are completely dry.

OUTDOOR USE: For control of ants, clover mites, crickets, earwigs, fleas, grasshoppers, sowbugs, and brown dog ticks, spray infested areas thoroughly including outside surfaces of screens, doors, window frames, foundations, patios or other places insects may use to enter the building. For ants, spray hills and runways. Repeat as necessary. Do not spray vegetation.

BROWN DOG TICKS: Spray sleeping quarters of pets, around baseboards, window and door frames, wall cracks and local areas of floors. Repeat as needed. Fresh bedding should be placed in animal quarters.