

# 3314-34

CAUTION: Keep away from humans, domestic animals and pets. If swallowed or absorbed by humans, domestic animals or pets, this material may reduce the clotting ability of the blood and cause bleeding. In that case, intravenous and oral administration of Vitamin K combined with blood transfusions are indicated as in the case of hemorrhage caused by overdoses of bishydroxycoumarin. Call a physician immediately.

Baits must be placed in areas inaccessible to children, pets, wildlife and domestic animals or in tamper proof bait boxes. Do not allow treated bait to be exposed on bare ground.

Do not reuse container. Destroy when empty.

Keep out of lakes, streams or ponds. Do not contaminate

GENERA specified INDOOR behind openings


OUTDOC and agri

FOR NO placement 10 days

FOR HO ment wit supply o

Contamin

Treated pets. wil



## 3314-34

- ~\*

LETS

CAUTION: Keep away from humans, domestic animals and pets. If swallowed or absorbed by humans, domestic animals or pets, this material may reduce the clotting ability of the blood and cause bleeding. In that case, intravenous and oral administration of Vitamin K combined with blood transfusions are indicated as in the case of hemorrhage caused by overdoses of bishydroxycoumarin. Call a physician immediately.

Baits must be placed in areas inaccessible to children, pets, wildlife and domestic animals or in tamper proof bait boxes. Do not allow treated bait to be exposed on bare ground.

Do not reuse container. Destroy when empty.

Keep out of lakes, streams or ponds. Do not contaminate

GENERAL DIRECTIONS: Can be used in wet and damp areas as specified below:

INDOORS: Place bait in runways, at feeding and watering areas, behind boards, along walls, between walls, and behind gnawed openings.

OUTDOORS: place bait in burrows, in sewers above high water line, and agricultural premises.


FOR NORWAY AND ROOF RATS: Place 4 to 16 ounces of bait per placement. Maintain constant supply of bait at stations for at least 10 days as long as there is evidence of rat population.

FOR HOUSE MICE: Place  $\frac{1}{2}$  oz. (tablespoon amounts) per placement with stations located at 8 to 12 ft. intervals. Maintain constant supply of bait for at least 15 days.

Contaminated or spoiled baits should be replaced promptly.

Treated baits must be placed in locations not accessible to children, pets. wildlife and domestic animals, or in tamper-proof bait boxes.

	,	, , ,		
		•		



the state

100.000%

Active Ingredient: 2-[(p-chlorophenyl) phenylacelyl] - 1, 3-indandione* Inert Ingredient	0.005% 99.995%

# 3314-34

CAUTION: Keep away from humans, domestic animals and pets. If swallowed or absorbed by humans, domestic animals or pets, this material may reduce the clotting ability of the blood and cause bleeding. In that case, intravenous and oral administration of Vitamin K combined with blood transfusions are indicated as in the case of hemorrhage caused by overdoses of bishydroxycoumarin. Call a physician immediately.

Baits must be placed in areas inaccessible to children, pets, wildlife and domestic animals or in tamper proof bait boxes. Do not allow treated bait to be exposed on bare ground.

Do not reuse container. Destroy when empty.

Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

## CP BAITS ARE OIL IMPREGNATED.

To insure the complete penetration of the bait by the toxicant, this product has been prepared in a two-step operation:

GENERA specified

INDOOI behind opening

OUTDO and ag

FOR NC placeme 10 days

FOR HC ment wi supply c

Contami

Treated pets, wi

Do not NOTICE material

# 3314-34

TS

5%

5%

0%

CAUTION: Keep away from humans, domestic animals and pets. If swallowed or absorbed by humans, domestic animals or pets, this material may reduce the clotting ability of the blood and cause bleeding. In that case, intravenous and oral administration of Vitamin K combined with blood transfusions are indicated as in the case of hemorrhage caused by overdoses of bishydroxycoumarin. Call a physician immediately.

Baits must be placed in areas inaccessible to children, pets, wildlife and domestic animals or in tamper proof bait boxes. Do not allow treated bait to be exposed on bare ground.

Do not reuse container. Destroy when empty.

Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

CP BAITS ARE OIL IMPREGNATED.

To insure the complete penetration of the bait by the toxicant, this product has been prepared in a two-step operation:

specified below:

openings.

OUTDOORS: place bait in burrows, in sewers above high water line, and agricultural premises.

FOR NORWAY AND ROOF RATS: Place 4 to 16 ounces of bait per placement, Maintain constant supply of bait at stations for at least 10 days as long as there is evidence of rat population.

FOR HOUSE MICE: Place  $\frac{1}{4} - \frac{1}{2}$  oz. (tablespoon amounts) per placement with stations located at 8 to 12 ft, intervals. Maintain constant supply of bait for at least 15 days.

Contaminated or spoiled baits should be replaced promptly.

Treated baits must be placed in locations not accessible to children, pets, wildlife and domestic animals, or in tamper-proof bait boxes.


NOTICE: Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.


KEEP AWAY FROM FOODSTUFFS. STORE IN A COOL PLACE. EPA Reg. No. 3314-34 EPA Ed. No. 3314-FL-1


GENERAL DIRECTIONS: Can be used in wet and damp areas as

INDOORS: Place bait in runways, at feeding and watering areas, behind boards, along walls, between walls, and behind gnawed

#### Do not reuse empty container. Destroy when empty.







1,	3-indandione*	
		100.000%



ground.

CAUTION: Keep away from humans, domestic animals and pets. If swallowed or absorbed by humans, domestic animals or pets, this material may reduce the clotting ability of the blood and cause bleeding. In that case, intravenous and oral administration of Vitamin K combined with blood transfusions are indicated as in the case of hemorrhage caused by overdoses of bishydroxycoumarin. Call a physician immediately.

Baits must be placed in areas inaccessible to children, pets, wildlife and domestic animals or in tamper proof bait boxes. Do not allow ireated bait to be exposed on bare

Do not reuse container. Destroy when empty.

Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

### CP BAITS ARE OIL IMPREGNATED.

To insure the complete penetration of the bait by the toxicant, this product has been prepared in a two-step operation:

GENERAL specified be

INDOORS: behind boo openings.

OUTDOORS and agricul

FOR NORW placement. 10 days as

FOR HOUS ment with supply of b

Contaminate

Treated ba pets, wildlif

Do not rea NOTICE: 6 material m

KEEP AW/

CAUTION: Keep away from humans, domestic animals and pets. If swallowed or absorbed by humans, domestic animals or pets, this material may reduce the clotting ability of the blood and cause bleeding. In that case, intravenous and oral administration of Vitamin K combined with blood transfusions are indicated as in the case of hemorrhage caused by overdoses of bishydroxycoumarin. Call a physician immediately.

Baits must be placed in areas inaccessible to children, pets, wildlife and domestic animals or in tamper proof bait boxes. Do not allow treated bait to be exposed on bare ground.

Do not reuse container. Destroy when empty.

LETS

0.005%

D,995%

0.000%

Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

## CP BAITS ARE OIL IMPREGNATED.

To insure the complete penetration of the bait by the toxicant, this product has been prepared in a two-step operation:

GENERAL DIRECTIONS: Can be used in wet and damp areas as specified below:

**INDOORS:** Place bait in runways, at feeding and watering areas, behind boards, along walls, between walls, and behind gnawed openings.

OUTDOORS: place bait in burrows, in sewers above high water line, and agricultural premises.

FOR NORWAY AND ROOF RATS: Place 4 to 16 ounces of bait per placement. Maintain constant supply of bait at stations for at least 10 days as long as there is evidence of rat population.

FOR HOUSE MICE: Place  $\frac{1}{2}$  oz. (tablespoon amounts) per placement with stations located at 8 to 12 ft, intervals. Maintain constant supply of bait for at least 15 days.

Contaminated or spoiled baits should be replaced promptly.

Treated baits must be placed in locations not accessible to children, pets, wildlife and domestic animals, or in tamper-proof bait boxes.

Do not reuse empty container. Design when empty. NOTICE: Suyer comunes off risks of use storage or Multing of s moterial pot is strict accordance with chaptions picks berewith.

LEEP ANELY FROM FOCOSTUPYE I



## NERIE RESISTANT PELLER NO RATE NO MICE

		· · · )
2-Lp-chloruption, 1) pherylacely1] -		Kee
1, 3-indandione*	0.005%	w'ai
Inert Ingredient	<b>99.995%</b>	ply
	<u> </u>	CP
	100.000%	To
		pro:
	100.000%	T



#### 

ep out of lakes, streams or plant. But lot contains a iter by cleaning of equipment, or disposal of wastes. Apr this product only as specified on this label.

## P BAITS ARE OIL IMPREGNATED.

insure the complete penetration of the bait by the toxicant, this duct has been prepared in a two-step operction:

.

pets, wild

Do nc?

NOTICE: material

KEEP AV

# .

· · · · · · . . .

#### rot i se catto an inter anter on pt,

thep out of lakes, streams or punds. Do not contaminute water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

## CP BAITS ARE OIL IMPREGNATED.

.

 $[ \mathbb{N} ]$ 

%

%

%

To insure the complete penetration of the bait by the toxicant, this product has been prepared in a two-step operation:



## 

. . • • • • •

· · · · Million At and a

- + + · · · · ··· 🔥 • · · · · · • •

untamposted of the extension of the second second

Treated taits must be placed of table station of the table pets, wildlife and domestic animals, or in tamper-proof built be and

Do not reuse empty container. Destroy when empty. NOTICE: Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

KEEP AWAY FROM FOODST'JFFS. STORE IN A COOL PLACE. EPA Reg. No. 3314-34 EPA Est. No. 3314-FL-1