

**PRECAUTIONARY STATEMENTS
HAZARD TO HUMANS AND DOMESTIC ANIMALS**

CAUTION

... ..
... ..
... ..

... ..
... ..
... ..
... ..
... ..
... ..
... ..
... ..
... ..
... ..
... ..

STATEMENT OF PRACTICAL TREATMENT

- **Inhaled** — Call physician immediately
- **Ingested** — Remove victim to fresh air and apply respiration if indicated
- **On skin** — Wash promptly with soap and water. Rinse thoroughly
- **In eyes** — Rinse eyes for at least 15 minutes with water and call physician immediately

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and wildlife. Do not apply directly to water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label. This product is highly toxic to bees exposed to direct treatment or residue on plants. Precautionary information may be obtained from your Cooperative Agricultural Extension Service.

PHYSICAL OR CHEMICAL HAZARDS

Combustibility: Do not use, pour, spill or store near heat or open flames. Handle in a well-ventilated area.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Storage: Store in a cool, dry place.

Pesticide Disposal: Pesticide, spray mixture or residue that cannot be used according to label instructions must be disposed of according to Federal, State or local procedures under the Resource Conservation and Recovery Act.

Container Disposal: 30 to 55 gallon drums: (a) Reuse container and offer for reconditioning, or (b) Triple rinse (or equivalent) and offer for recycling or reconditioning, or dispose of in a sanitary landfill, or by other approved State and local procedures.

Smaller Containers: Treat as in (b) above.

General: Consult Federal, State or local disposal regulations for approved alternative procedures.

INDO-THRIN FOOD PLANT SPECIAL

Designed for Use in Edible Product Areas of Official Establishments Operating Under the Meat, Poultry, Shell Egg Grading and Egg Products Inspection Programs and in Other Food Processing Plants.

Apply with Equipment Designed and Adjusted to Deliver a True Aerosol Mist



SPACE SPRAY: To kill accessible stages of houseflies, including fruit flies, small flying moths, gnats, cheese skippers, and these common stored product pests granary weevils, rice weevils, confused flour beetles, saw toothed grain beetles, spider beetles, cigarette beetles, Drugstore beetles, Angoumois grain moths, Mediterranean flour moths, Indian meal moths, Tobacco moths, meal worms and grain mites. Close windows and doors and shut off ventilating systems. Use a fogger or vaporizer adjusted to deliver an aerosol spray (no droplets over 50 microns in diameter and 80% less than 30 microns). Apply at the rate of 1 ounce per 1000 cubic feet of space. Direct the spray toward the ceiling and upper corners of the area behind obstructions. Vacate and close off treated area. Do not reenter for at least 15 minutes after application. Ventilate before reoccupying.

In mushroom production and processing. To kill mushroom flies or fungus gnats in growing houses, apply at a rate of 10 oz per 1000 cubic feet of space. Fogger should be operated 8 to 10 feet away from beds and directed slightly upward over them. Circulating fans should be left in operation, but for best results, refrigeration should be turned off. In packing and canning plants, use at a rate of 1 ounce to 2000 cubic feet of space.

SPACE AND CONTACT SPRAY: To kill...

