

GRAYS, MOSQUITOES, BITTER FLIES, HORNETS, AND/OR HOUSE BROWN MOTHS, MEDITERRANEAN FLOUR MOTHS, INDIAN MEAL MOTHS, AND TOBACCO MOTHS: direct spray at 30 second angle, distributing it uniformly throughout the entire area at a rate of 1.4 ounces per 1000 cubic feet of space.

FOGGING: To kill exposed or accessible stages of the above named crawling insects, use a rate of one gallon to 20,000 to 40,000 cubic feet of space. Contact insects directly whenever possible.

ON ANIMALS: To protect cattle and horses from **HORN FLIES, HOUSE FLIES, MOSQUITOS, AND GRAYS**, apply a light mist sufficient to wet the surface of the hair. To control **STABLE FLIES, HORSE FLIES, AND BEEH FLIES**, apply at a rate of 2 ounces per adult cow or horse sufficient to wet the hair thoroughly. Repeat treatment daily or at intervals necessary to give continued protection.

TO CONTROL BLOOD-SUCKING LICE, apply to the infested areas of the animal using a tall brush to get the spray to the base of the hair. Repeat every 7 or 10 weeks if required.

OUTDOOR ADULT MOSQUITO CONTROL: Apply as a fine mist to thoroughly wet the entire area. For best results, spray when the wind velocity does not exceed 3 miles per hour. Treat shrubbery and vegetation where mosquitoes may rest, particularly around the perimeter of mosquito breeding sites such as swamps, marshes, stagnant pools, and ponds. Apply at a rate of 1 ounce per 1000 square feet (or 2 1/2 pints per acre).

Using brush-mounted conventional mixing equipment such as Rotomax[®] models SIC, VI, or H80 HT, working at 5 mph to treat a 100 foot swath, calibrate equipment to deliver 4.5 gallons per minute.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Avoid contamination of food and feedstuffs. Remove cats, birds and other fish species before spraying.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, ponds, streams, tidal marshes, and estuaries. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DISPOSAL

Do not reuse empty container. Wrap container and put in trash collection.

Buyer assumes all risks of use, storage or handling of this product not in strict accordance with directions given hereon.

FLEA FLIP[®] READY-TO-USE

A Concentrated Flea Flip[®] Micro Emulsion
Clear • Stable

Leaves No Oily or Objectionable Residue

In view of Maximum Energy Storage, this product is extremely low in petroleum distillates.

Effective, Fast Acting Insecticide

For Use in Homes, Institutions, Restaurants, Food Processing Plants, Dairies, Linnark and Outdoors for Mosquito Control

Apply with Conventional Mechanical or Compressed Air Equipment

ACTIVE INGREDIENTS:

Pyrethrin 0.1%

*Pyrethrin Butoxide, technical 1.0%

Petroleum Distillate 0.4%

INERT INGREDIENTS:

*Equivalent to 0.8% (butylcarbyl) (6 propylpropenyl) ether and to 0.2% related compounds

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

Harmful if swallowed

See side panel for additional precautionary statements

NET CONTENTS:

EPA Reg. No. 3314-44 EPA Est. No. 3314FL-1

Manufactured by: Colonial Products, Inc.
1830 Tenth Avenue, North, Lake Worth, Florida 33461

INDOOR APPLICATION

The use of this product in food processing establishments should be confined to one period when the plant is not in operation. Food should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using. Vessels treated area and washers before reapplying. For maximum effectiveness, a combination of spot surface treatment and fogging is recommended.

SURFACE SPRAYING: TO CONTROL FLEAS AND BROWN DOG TICKS:

Thoroughly spray ... areas, pet beds, living quarters, nearby cracks and crevices, along and behind baseboards, moldings, window and door frames, and localized areas of floor and floor covering. Fresh bedding should be placed in animal quarters following treatment. Concurrent treatment of animals with an approved acaricide is recommended. Repeat treatment as needed.

TO CONTROL ROACHES, ANTS, SILVERFISH, SPIDERS, CRACKS, CLOVER

WITTS, CHEESE WITTS, GRANARY WEEVILS, RICE WEEVILS, COMBUSTED FLOUR BEETLES, RUST RED FLOUR BEETLES, SAW-TOOTHED GRAIN BEETLES, SPIDER BEETLES, CHEMIST BEETLES, DRUGSTORE BEETLES, MEAL WORMS, GRAIN WITTS, AND CABELLER: Use a good sprayer adjusted to deliver a coarse mist spray. Direct the spray into hiding places, cracks, and crevices, under pallets, around corners of stored loads, around the base of machinery, behind the shelves and drawers. For ants spray cracks, nests and points of entry. If surface application only it is to be used, spray floors, walls, and other surfaces applying at a rate of 1 gallon to 750 square feet of surface.

TO CONTROL CARPET BEETLES:

Spray edges of carpeting and under carpeting and rug. Make localized application to floor and baseboards. Spray directly into cracks, closets and adjacent areas of drapery. Repeat treatment as needed.

TO CONTROL BEDBUGS:


Spray mattresses lightly, particularly rolls and seams. Take beds apart and spray in all areas of drapery. Repeat treatment as needed.

TO CONTROL CLOTHES MOTHS:


Remove any infested articles from storage, brush thoroughly and air for several hours in sunlight, if possible. Apply spray liberally to empty chests, closets, drawers, and other storage, directing spray into cracks, joints, and crevices. After using, infested articles may be treated lightly keeping inside at least three feet from fabric to avoid staining.

SPACE SPRAYING:

Close doors and windows and shut off ventilating systems. Apply with a good machine adjusted to deliver a fine mist or fog. For rapid control of **FLIES, BROWN FLIES,**



* SOLID BLACK



GRATE, MOSQUITO, SHIPPER FLIES, HORNETS, ANNOYING GRASS MOTHS, MEDITERRANEAN FLOUR MOTHS, INDIAN MEAL MOTHS, AND TOBACCO MOTHS: direct spray at an upward angle, distributing it uniformly throughout the entire area at a rate of 3-4 ounces per 1000 cubic feet of space. Contact insects directly whenever possible.

FOGGING: To kill exposed or accessible stages of the above named crawling insects, use a rate of one gallon to 20,000 to 40,000 cubic feet of space. Contact insects directly whenever possible.

ON ANIMALS: To protect cattle and horses from **HORN FLIES, HOUSE FLIES, MOSQUITOS, AND GRATE,** apply a light mist sufficient to wet the surface of the hair. To control **STABLE FLIES, HORSE FLIES, AND DOSE FLIES,** apply at a rate of 2 ounces per adult cow or horse sufficient to wet the hair thoroughly. Repeat treatment daily or at intervals necessary to give continued protection.

TO CONTROL BLOOD-SUCKING LICE, apply to the infested areas of the animal using a stiff brush to get the spray to the base of the hair. Repeat every 7 or 10 weeks if required.

OUTDOOR ADULT MOSQUITO CONTROL: Apply as a fine mist to thoroughly cover the entire area. For best results, treat when the wind velocity does not exceed 5 miles per hour. Treat shrubbery and vegetation where mosquitoes may rest, particularly around the perimeter of mosquito breeding areas such as swamps, marshes, stagnant pools, and ponds. Apply at a rate of 1 ounce per 1000 square feet (or 2 1/2 pints per acre).

Using brush-mounted conventional mixing equipment such as **Racoma®** models SIC, 91, or 100 HT, treating at 5 mph to treat a 100 foot swath. Calibrate equipment to deliver 4.5 gallons per minute.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS
CAUTION

Harmful if swallowed. Avoid contamination of food and feedstuffs. Remove pets, birds and cover fish tanks before spraying.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, ponds, streams, tidal marshes, and estuaries. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DISPOSAL

Do not reuse empty container. Wrap container and put in trash collection.

Buyer assumes all risks of use, storage or handling of this product not in strict accordance with directions given hereon.

FLEA FLIP® READY-TO-USE

**A Concentrated Flea Flip® Micro Emission
Clear • Stable
Leaves No Oily or Objectionable Residue**

In case of National Energy Shortage, this product is extremely low in petroleum content.

Effective, Fast Acting Insecticide

For Use in Homes, Institutions, Restaurants, Food Processing Plants, Diner, Livestock and Outdoors for Mosquito Control

Apply with Conventional Mechanical or Compressed Air Equipment.

ACTIVE INGREDIENTS:

Pyrethrin	0.1%
Pyrethrin Butoxide, technical	1.0%
Petroleum Distillate	98.9%
INERT INGREDIENTS:	99.9%

*Equivalent to 0.2% (bicyclohexyl) (6-propylpiperonyl) ether and to 0.2% related compounds.

KEEP OUT OF REACH OF CHILDREN
CAUTION

Harmful if swallowed

See side panel for additional precautionary statements

NET CONTENTS

EPA Reg. No. 3314-44 EPA Est. No. 3314FL-1

Manufactured by: Colonial Products, Inc.

1830 Tenth Avenue, North, Lake Worth, Florida 33461

INDOOR APPLICATION

The use of this product in food processing establishments should be confined to case periods when the plant is not in operation. Food should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using. Vases treated area and vacuums before reoccupying. For maximum effectiveness, a combination of spot surface treatment and fogging is recommended.

SURFACE SPRAYING: TO CONTROL FLEAS AND BROWN BOB TICKS: Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind bookshelves, moldings, window and door frames, and localized areas of floor and floor covering. Fresh bedding should be placed in animal quarters following treatment. Concurrent treatment of animals with an approved insecticide is recommended. Repeat treatment as needed.

TO CONTROL ROACHES, ANTS, SILVERFISH, SPIDERS, CRICKETS, CLOVER MITES, CHEESE MITES, GRANARY WEEVILS, RICE WEEVILS, CONFUSED FLOUR BEETLES, MUSTY RED FLOUR BEETLES, SAW-TOOTHED GRAM BEETLES, JEROME BEETLES, CRABAPPLE BEETLES, BRONSTONE BEETLES, MEAL WORMS, GRASS MITES, AND CABERLES: Use a good sprayer adjusted to deliver a coarse mist spray. Direct the spray into hiding places, cracks, and crevices, under pallets, around containers of stored foods, around the base of machinery, behind the shelves and drawers. For silverfish spray bookshelves. For ants spray trails, nests and points of entry. If surface application only is to be used, spray floors, walls, and other surfaces applying at a rate of 1 gallon to 750 square feet of surface.

TO CONTROL CARPET BEETLES: Spray edges of carpeting and under carpeting and rug. Make localized application to floor and bookshelves. Spray directly into cracks, closets and infested areas of shelving. Repeat treatment as needed.

TO CONTROL BEDBUGS: Spray mattresses lightly, particularly tufts and seams. Take beds apart and spray in all areas of shelving. Repeat treatment as needed.

TO CONTROL CLOTHES MOTHS: Remove any infested articles from storage, brush thoroughly and air for several hours in sunlight, if possible. Apply spray liberally to empty chests, closets, bureaus, and other storage, directing spray into cracks, joints, and crevices. After using, infested articles may be treated lightly keeping article at least three feet from fabric to avoid staining.

SPACE SPRAYING: Close doors and windows and shut off ventilation systems. Apply with a good machine adjusted to deliver a fine mist or fog. For rapid control of **FLIES, FLEET FLIES,**

ACCEPTED
APR 8 1981
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 3314-44

*SOLID BARK