

EPA Reg. No. 3314-50 EPA Est. No. 3314-FL-1

AS A SPACE SPRAY: Dilute at a rate of 1 part concentrate to 19 parts water or oil. Apply with mechanical or compressed air equipment (non thermal) adjusted to deliver a fine mist. Close doors and windows and shut off ventilating systems. **For rapid control of flies, fruit flies, gnats, mosquitoes, skipper flies, wasps, hornets, Angoumois grain moths, Mediterranean flour moths, Indian meal moths, and tobacco moths:** direct spray at an upward angle distributing it uniformly through the entire area at a rate of 1 ounce per 1000 cubic feet of space. Keep area closed for at least 10 minutes. Ventilate and sweep up dead insects before reoccupying.

For rapid kill of exposed or accessible stages of the insects named on this label dilute 1 part concentrate to 1 to 9 parts water or oil. Apply using mechanical or compressed air equipment (not thermal) follow directions for space spraying. Apply the above dilutions at rate of from 1 to 10 ounces per 1000 cubic feet of space.

When used in meat and poultry processing plants, a dilution of no less than 1 to 1 (50%) is acceptable and it must be dispersed from equipment adjusted to deliver an aerosol spray (no droplets over 50 microns in diameter and 80% less than 30 microns.)

DISPOSAL

Do not reuse empty container. Wrap container and put in trash collection.

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS**

CAUTION

Harmful if swallowed. The use of this product in food processing plants should be confined to time periods when the plant is not in operation. Foods should be removed or covered before treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using. Remove any birds, pets or fish aquarium before spraying.

ENVIRONMENTAL HAZARDS:

This product is toxic to fish. Keep out of lakes, streams, and ponds.

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

INDO-THRIN S CONCENTRATE

A Food Industry Insect Spray
May Be Diluted With Either Water Or Oil



ACTIVE INGREDIENTS:

Allethrin (Allyl Homolog of Cinerin i)	2.50%
Related Compounds	0.28%
*Piperonyl Butoxide, Technical	10.00%
Petroleum Distillate	77.22%
INERT INGREDIENTS	10.00%
	100.00%

*Equivalent to 8-(4-butylcarboxyl)-(6-Propyl piperonyl) ether and 2% related compounds

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See side panel for additional precautions

NET CONTENTS _____



Manufactured by
Colonial Products, Inc.
1830 Tenth Ave. North, Lake Worth, FL 33461

DIRT

It is a violation of Federal law if the label is inconsistent with its label. When diluting with water, use moderate use. Holding a recommended. Agitate or petroleum oil is recommended. Vacate premises after treatment. Repeat treatment when necessary.

For maximum effectiveness, use application and space treatment.

AS A SURFACE SPRAY:


For maximum effectiveness, use application and space treatment. This dilution may not be used on inspected meat and poultry processing plants. dilute 1 to 10.

To control roaches, ants, cheese mites, granary weevil, red flour beetles, and cigarette beetles, Drugstore beetles: Use a good spray. Direct the spray into hiding places around containers of stored food behind shelves and drawers. spray trails, nests and pour on. spray floors, walls, 750 square feet of surface.

To control carpet beetle: carpeting and rugs. Mice, baseboards. Spray directly on shelves. Repeat treatment.

To control bedbugs: Spray tufts and seams. Take down baseboards, moldings and

To control fleas and ticks: areas, pet beds, resting quarters and behind baseboards, in localized areas of floor and placed in animal quarters for treatment of animals with an approved treatment as needed.



INDO-THRIN S CONCENTRATE

A Food Industry Insect Spray
May Be Diluted With Either Water Or Oil



ACTIVE INGREDIENTS:

Allethrin (Allyl Homolog of Chlorthal)	2.50%
Related Compounds	0.28%
*Piperonyl Butoxide, Technical	10.00%
Petroleum Distillate	77.22%

INERT INGREDIENTS

100.00%

*Equivalent to 1.5% of active ingredient and 2% of total weight.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See side panel for additional precautions

NET CONTENTS _____



Manufactured by
Colonial Products, Inc.

1830 Tenth Ave. North, Lake Worth, FL 33461

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

When diluting with water, prepare only the amount needed for immediate use. Holding a water dilution more than 24 hours is not recommended. Agitate occasionally. Final dilutions of a deodorized petroleum distillate are recommended.

Evacuate premises after treatment and ventilate before reoccupying.

Repeat treatment when reinfestation occurs.

For maximum effectiveness, a combination of localized surface application and space treatment is recommended.

AS A SURFACE SPRAY: For initial clean-up of severe insect infestations, dilute at a rate of 1 part concentrate in 4 parts water or oil. This dilution may not be used in food product areas of federally inspected meat and poultry plants, in these plants and for normal infestations dilute 1 part concentrate in 9 parts water or oil.

To control roaches, ants, silverfish, spiders, crickets, clover mites, cheese mites, granary weevils, rice weevils, confused flour beetles, rust-red flour beetles, sawtoothed grain beetles, spider beetles, cigarette beetles, drugstore beetles, meal worms, grain mites, and casidites: Use a good sprayer adjusted to deliver a coarse wet spray. Direct the spray into hiding places, cracks and crevices, under pallets around containers of stored foods, around the base of machinery behind shelves and drawers. For silverfish spray bookcases. For ants spray trails, nests and points of entry. If surface application only is to be used, spray floors, walls and other surfaces at a rate of 1 gallon to 150 square feet of surface.

To control carpet beetles: Spray edges of carpeting and under carpeting and rugs. Make localized application to floor and baseboards. Spray directly into cracks, closets and infested areas of sleeping. Repeat treatment as needed.

To control bedbugs: Spray mattresses lightly, particularly around tufts and seams. Take beds apart and spray in all joints. Treat baseboards, moldings and floors. Repeat treatment as needed.

To control fleas and brown dog ticks: Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind baseboards, moldings, window and door frames and localized areas of floor and floor covering. Fresh bedding should be placed in animal quarters following treatment. Concurrent treatment of animals with an approved insecticide is recommended. Repeat treatment as needed.