PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

Keep Out Of Reach Of Children. May be fatal if swallowed. May be absorbed through the skin. May be irritating to eyes and skin. Do not wear contaminated clothing. Do not get in eyes or on skin. Do not breathe vapor spray. Wash thoroughly after handling. Wash contaminated clothing before re-use. Discard contaminated shoes with gloves of rubber and get medical attention. Avoid breathing spray mist or vapors. Do not spray vegetables grown for food or other food plants except for food protection.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. Do not apply directly to water. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply when weather conditions favor drift from treated areas. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protect bee colonies. May be registered from your Cooperative Agricultural Extension Service.

PHYSICAL AND CHEMICAL HAZARDS

Do not cut, grind, weld or solder near heat or open flame. Keep container tightly sealed. Volatile concentrate is combustible.

STORAGE & DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Do not reuse empty container. Keep container and out of reach of children.

NOTICE Seller's warranty shall be limited to the terms of this label and specified formula. The buyer assumes any risk to persons or property arising out of the use or handling and accepts this product on these conditions.

PESTA REST

LAWN & ORNAMENTAL INSECT SPRAY

CONCENTRATE

Controls Chinch Bugs, Fleas, Ants, Clover Mites, Sod Webworms (Lawn Moth), Armyworms, Cutworms, Earwigs, Brown Dog Ticks, Aphids, Grasshoppers and other lawn and ornamental insects listed.

Keep Out Of Reach Of Children

WARNING

See Side Panel For Additional Precautionary Statements

STATEMENT OF PRACTICAL TREATMENT

If Swallowed — Do not induce vomiting. Call a Physician immediately.

If On Skin — In case of contact immediately remove contaminated clothing and immediately wash skin with soap and water.

If In Eyes — Flush eyes with plenty of water. Call a physician immediately.

Note To Physician — If symptoms of cholinesterase inhibition are present, atropine only by injection is antidotal.

ACTIVE INGREDIENTS

Chlorpyrifos, O,O-diethyl O-(3,5,6-trichloro-2-pyridinyl) phosphorothioate	16.00%
Inert Ingredients	77.25%
	6.75%
Total	100.00%

COLONIAL PRODUCTS, INC.
1630 South Avenue North, Lake Worth, FL 33461

DIRECTIONS FOR USE

Read entirely and follow this product in a manner consistent with the instructions.

Place of Lawns and other Ornamental and Recreational Turf Areas areas

INSECTS CONTROLLED Ants, Armyworms, Chiggers, Chinch Bugs, Clover Mites, Crickets, Cutworms, Earwigs, Fleas, Grasshoppers, Sod Webworms, Ticks

WHEN TO TREAT Treat when pests first appear or their damage to turf is observed. Retreat as needed for control. To prevent Chinch Bug damage apply as soon as daily temperatures reach 80°F and repeat every 7 months. For best results the turf should be mowed before application.

HOW TO TREAT – Apply spray as a coarse low pressure spray using sprinkler hose and spray or as pressure spray. Thoroughly water this spray after application to wash the insecticide into the turf, damage to sod webworms. Use a dilution of 3 to 5 gallons of water per 1,000 sq. ft. Details as follows:

AREA	AMT. TURF PRODUCT TO USE	AMT. WATER
1,000 Sq. Ft.	3.5 oz. (7 Tbsp.)	3.5 Gallons
5,000 Sq. Ft.	17.5 oz. (1 pint + 3 Tbsp.)	15-25 Gallons

For Sod Webworms delay watering and mowing for 24 hours afterwards.

DIRECTIONS FOR USE

Points of Ornamental Flowers, Shrubs, Vines, Shade & Flowering Trees

Use this product for the control of insect pests as listed when they are found on flowering plants. Use a hand fine spray or pressure sprayer to provide complete and uniform coverage of infested plants. Avoid spraying to point of run-off. Treat when pests appear and repeat at 7 to 10 day intervals as needed for control. NOTE: Do not use on Azaleas, Camellias, Pyracantha, Rose Bushes or evergreens by reason of possible injury to these plants.

INSECTS CONTROLLED Aphids, Bagworms, Eastern Tent Caterpillar, Fall Webworms, Grasshoppers, Mites, Orangestriped Oakworm, Spittlebug, Tuliptree Scale and Caterpillar

How to Mix: Mix 1 1/3 ounces in 3 gallons of water or 4 oz. in 10 gallons of water. Thoroughly spray all foliage.

INSECTS CONTROLLED Leafminers, Leafhoppers, Mites, Aphids, Webworms, Southernpine Coneworm, Tent Caterpillar (Eastern)

How to Mix: Mix 2 2/3 oz. in 3 gallons of water or 9.5 oz. in 10 gallons of water. Thoroughly spray all foliage.

INSECTS CONTROLLED Ash Borer, Lilac Borer, Oriental Beetle, Scale, Cicadas, Rose Curculio, Pine Needle, Tea, and Spruce Peach Scales

How to Mix: Mix 5 1/3 oz. in 3 gallons of water or 17.5 oz. in 10 gallons of water. Thoroughly spray all foliage.

INSECTS CONTROLLED Peach Tree Borer. Apply to Peach trees and shrubs of the genus Prunus as a trunk spray before early flight (trunk inside the tree). Mix 16 ounces in 3 gallons of water. In general use a coarse low pressure spray to thoroughly wet all portions of the trunk up to the first scaffold branches. Treat the first of May, June and July making three applications one month apart. Do not allow spray to get on fruit.

EPA Reg. No. 3314-64 EPA EST. NO. 37-[illegible]

See Contents

PROTECTIVE BAND & FOUNDATION SPRAY

BAND SPRAY TREATMENT To keep pests out of buildings by band treatment (Earwigs, Millipedes and Sowbugs (Wood lice)) mix 2 ounces of spray in 1/2 liter water (more or less adjusted to buildings). Take the building's circumference or height of 2 to 3 feet, make a plastic bag around building (wall, porch, etc.). Use 4 to 5 gallons of 1/2 gallon application to treat 1,000 sq. ft. (100 ft. square or 1/2 full long).

TURFGRASS WEEVILS (Hyperodes) For control during the period of adult activity in the Chief Hyperodes spots in late April and again in late May or as recommended by your local Agricultural Extension Service specialist or other area.

WHITE GRUBS European Chafer and Japanese Beetle Larvae. Use from 8 oz. to 16 oz. in 1 gallons of water to cover 1,000 sq. ft. Spray when grubs are young and actively feeding near the soil surface, usually during late July and August or as recommended by your local Agriculture Extension Service specialist or other area. Immediately after applying, irrigate the treated area with 1/2 to 1 inch of water to wash the insecticide into the underlying soil.

BLUEGRASS BILLBUG – Mix at the same rate as for White Grubs. Spray early in the season when adults first appear. Retreat when needed.

FIRE ANT CONTROL – Apply as a drench to pot materials infesting spray at the rate of 1 1/3 oz. in 4 gallons water. Gently agitate a total of 1 gallon of diluted solution over the surface of each mound using a sprinkling can. Thoroughly wet mound and surrounding area to a 4 ft. diameter (12 sq. ft.). For best results apply at soil surface. Apply early in the morning or late evening hours. Treat late afternoon. Apply 30°F or on days warming to late morning hours. Treat late afternoon. Dry weather. Protectant sprays may disturb the ants and cause drop-down, reducing product effectiveness.

ATTENTION Turf and Ornamental Plant Pests: Keep children and pets off treated areas until spray has dried. Keep out of lawn & garden hydrants and other sources of water. Do not allow livestock to graze in treated grass. Do not feed treated grass clippings or weed screenings to livestock nor use hay for livestock bedding.

NOTE Mosquitoes coming to rest on areas treated for control of turf and or surrounding plant pests will be killed for varying periods of time after treatment depending on exposure of treated area to weathering conditions.

ACCEPTED
3314-64
MAY 4 1982
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No.

BEST DOCUMENT AVAILABLE