

17


# P.T.O. DURSBAN - DDVP 1.25 E.G.

INDOOR & OUTDOOR  
PEST CONTROL

ROACHES, ANTS, CRICKETS, FLEAS,  
SPIDERS, CHINCH BUGS,  
SOD WEBWORMS.

ORNAMENTAL INSECT CONTROL

Only For Sale To, Use And Storage By  
Pest Control Operators or Other Servicepersons



**ACTIVE INGREDIENTS:**

Chlorpyrifos (0,0-diethyl 0-(3,5,6-trichloro 2-pyridyl) phosphorothioate) .....	12.79%
2,2-Dichlorovinyl dimethyl phosphate .....	3.19%
Related Compounds .....	0.24%
Xylene Range Aromatic Solvent .....	77.76%
<b>INERT INGREDIENTS:</b> .....	6.03%
<b>Total</b>	<b>100.00%</b>

Contains 1 lb. Chlorpyrifos & .25 lb. Dichlorvos (DDVP) per Gallon  
EPA REG. NO. 331467 EPA Est. No. 3314-FL-1

**KEEP OUT OF REACH OF CHILDREN  
DANGER**

See Side Panel For Additional Precautionary Statements  
**STATEMENT OF PRACTICAL TREATMENT**

**If Swallowed:** Call a physician or Poison Control Center immediately. Gastric lavage is indicated if material was taken internally. DO NOT INDUCE vomiting unless other treatment is not available. Vomiting may cause aspiration pneumonia. If it is necessary to induce vomiting, give victim one or two glasses of water and insert finger in back of throat. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious person.

**If Inhaled:** Remove victim to fresh air. Apply respiration if indicated.

**If On Skin:** Wash immediately with soap and water. Call a physician immediately.

**If In Eyes:** Flush eyes with plenty of water. Call a physician immediately.

**If Warning Symptoms Appear:** Keep patient prone and quiet. Start artificial respiration immediately, if patient is not breathing. Transport patient immediately to the nearest physician.

Net Contents \_\_\_\_\_ Gallon

BEFORE DOCUMENT AVAILABLE

**STORAGE AND DISPOSAL**

**Pesticide Disposal**

**Container Disposal**

**NOTICE**

**DIRECTIONS FOR USE**

**PESTS OUT-OF-DOORS**

**PESTS ON OUTSIDE SURFACES OF BUILDINGS**

**NOTE** — To help prevent infestation of buildings by Ants, Crickets, Earwigs, Millipedes, and Sowbugs (Pillbugs) treat a band of soil 6 to 10 feet wide around and adjacent to buildings, as to building's foundation to a height of 2 to 3 feet, where pests are active and may find entrance. Use 32 ounces (1 quart) in 100 gallons of water and apply as a coarse spray at the rate of about 10 gallons spray mixture per 1,000 sq. ft. to thoroughly and uniformly wet the band area.

**PESTS OF LAWNS, OTHER ORNAMENTAL AND RECREATIONAL TURF AREAS AND PERENNIAL TURF GRASSES GROWN FOR SEED**

Use this product to control insects listed in the following table by application at the recommended rate and in accordance with the directions given below. Dilute in enough water to obtain complete and uniform coverage of pest infested areas and apply as a coarse, wet pressure spray using suitable spray equipment. Thoroughly water after treatment to wash the insecticide into the turf spaces, if needed. For best results, the lawn should be moist at time of treatment.

**PESTS ON ORNAMENTALS:** Use this product to treat flowers, shrubs, vines, shade and flowering trees, and evergreens found to be infested with the pests listed in the following table. Dilute with water according to directions given in the table and apply using suitable hand or power spray equipment in a manner to provide complete coverage. For best results apply a wetting spray to both upper and lower leaf surfaces and infested limbs and trunk areas. Attempt to penetrate dense foliage but avoid overspraying. Treat when pests appear and repeat at 7 to 10 day intervals, if needed. **CAUTION:** Do not use on Azaleas, Camellias, Foinerlias, Rose Bushes, or varieties by reason of possible injury to foliage.

**DIRECTIONS FOR USE  
DOSAGES AND DIRECTIONS FOR USE ON TURF**

PEST	DOSAGE per 1,000 sq. ft.	REMARKS
Ants	32 oz. in 100 gal. water	Treat a band of soil 6 to 10 feet wide around and adjacent to buildings, as to building's foundation to a height of 2 to 3 feet, where pests are active and may find entrance.
Bluegrass Bribug	8 to 12 Ounces	Spray early in the season when adults first appear. Retreat when needed.
Cricket	32 oz. in 100 gal. water	Treat a band of soil 6 to 10 feet wide around and adjacent to buildings, as to building's foundation to a height of 2 to 3 feet, where pests are active and may find entrance.
Earwig	32 oz. in 100 gal. water	Treat a band of soil 6 to 10 feet wide around and adjacent to buildings, as to building's foundation to a height of 2 to 3 feet, where pests are active and may find entrance.
Millipede	32 oz. in 100 gal. water	Treat a band of soil 6 to 10 feet wide around and adjacent to buildings, as to building's foundation to a height of 2 to 3 feet, where pests are active and may find entrance.
Sowbug (Pillbug)	32 oz. in 100 gal. water	Treat a band of soil 6 to 10 feet wide around and adjacent to buildings, as to building's foundation to a height of 2 to 3 feet, where pests are active and may find entrance.
DOGWOOD BORER	1 qt. in 25 gal. water	Mix 1 gallon in 100 gallons water (1 quart in 25 gal.) and spray trunks and lower limbs of trees when adults begin to emerge. Consult your local Agricultural Extension Service for the proper timing.
PEACH TREE BORER	1 qt. in 25 gal. water	Mix 1 qt. in 25 gal. water. Apply to flowering trees and shrubs of the genus PRUNUS. Thoroughly spray the trunk at ground level up to the first limb. Apply as coarse wet spray 3 times May 1st, June 1st, and July 1st. Do not treat the fruit or upper portions of the tree. This treatment is for ornamentals only.

NOTE: Use 32 oz. in 100 gal. water for outdoor use.

3 Fluid Ounces/1,000 sq. ft. = 1 gallon per acre

**DOGWOOD BORER** — Mix 1 gallon in 100 gallons water (1 quart in 25 gal.) and spray trunks and lower limbs of trees when adults begin to emerge. Consult your local Agricultural Extension Service for the proper timing.

**PEACH TREE BORER** — Mix 1 qt. in 25 gal. water. Apply to flowering trees and shrubs of the genus PRUNUS. Thoroughly spray the trunk at ground level up to the first limb. Apply as coarse wet spray 3 times May 1st, June 1st, and July 1st. Do not treat the fruit or upper portions of the tree. This treatment is for ornamentals only.

**NOTE—MOSQUITOES** coming to rest on areas treated for control of Turf and Ornamental plant pests will be killed for varying periods of time after treatment, depending on exposure of treated areas to weathering conditions.

**AREA CONTROL OF TICKS AND CHIGGERS** — Use for noncrop areas such as roadsides, footpaths, and trails, picnic and camping sites, parks and other recreational areas where these pests are present and create a nuisance or possible public health problem. Apply 1 quart per acre in sufficient water to moist all areas including low underbrush, grassy areas, weeds, and ground surface and debris using enough spray volume to obtain thorough coverage, usually 40 to 100 gallons of water per acre.



**COLONIAL PRODUCTS, INC.**  
1830 Tenth Avenue North  
Lake Worth, FL 33461

Product literature available

**DIRECTIONS FOR USE**  
**ATTENTION:**

**DOSAGES AND DIRECTIONS FOR USE ON ORNAMENTAL PLANTS**

Pests	Amount of Concentrate to use in Water to Make
	3 Gal 10 Gal 100 Gal
Cockroaches, Ants, Crickets, Firebrats, Silverfish, Spiders, Fleas and Brown Dog Ticks	1/2 oz 1 1/2 oz 1 1/2 oz
Blattella, Cimex, Pulex, Periplaneta, Solenopoda, Formica, Lasius, Tetranychus, Ixodes, Diplolepis, Acanthosoma, and other insects	1/2 oz 1 1/2 oz 1 1/2 oz
Cockroaches, Ants, Crickets, Firebrats, Silverfish, Spiders, Fleas and Brown Dog Ticks	1/2 oz 1 1/2 oz 1 1/2 oz

**DIRECTIONS FOR USE**

**INDOOR PEST CONTROL**

Do not use this product in the production areas of food processing plants, restaurants or other areas where food is commercially prepared or processed. Do not use in serving areas where food is exposed. **USE ONLY IN NON-FOOD AREAS** as directed below.  
 Use this product as a spot treatment for the control of Cockroaches, Ants, Crickets, Firebrats, Silverfish, Spiders, Fleas and Brown Dog Ticks in and around residential, institutional and commercial buildings, including homes, apartment buildings, hospitals, stores, and warehouses. It may be used with water or a decolorized base oil for a maintenance spray or extra strength spray. Follow directions for control of pests listed and repeat as needed for control.

**DILUTION CHART**

Amount of Spray (Water or Oil)	Amount of Concentrate To Use	
	Maintenance Spray	Extra Strength Spray
1 Gallon	2 1/2 ounces	5 1/2 ounces
5 Gallon	13 ounces	28 ounces
10 Gallon	26 ounces	56 ounces

The lower strength should be used as a maintenance spray while the stronger rate should be used to quickly reduce heavy infestations of Cockroaches, Ants, Crickets, Firebrats, Silverfish, and Spiders.  
**HOW TO APPLY** — Apply as a coarse spray or with a paint brush to localized areas where the above mentioned pests are found or normally occur, including dark corners of rooms, and closets, cracks and crevices in walls, along and behind baseboards, beneath and behind sinks, stoves, refrigerators and cabinets; around plumbing and other utility installations. For Ants apply to ant trails, also around doors and windows and wherever these pests may find entry. For Brown Dog Ticks (Do not treat pets with this product) apply as a spot treatment to infested areas such as pet's bedding and resting quarters; nearby cracks and crevices along and behind baseboards, windows and door frames and other localized areas where pests may hide. Old bedding should be removed and replaced with fresh clean bedding after treatment.

**DIRECTIONS FOR APPLICATION IN NON-FOOD AREA**

Non-Food areas include: Garbage Rooms, Lavatories, floor drains (to sewer entries and vestibules) Offices, locker rooms, machine rooms, boiler rooms, garages, mop closets and storage (after canning or bottling). Use the "maintenance" rate (See Dilution Chart above) as a spot treatment by applying as a coarse spray or with a paint brush to areas where Cockroaches are normally found. Apply to baseboard areas, around water pipes, surfaces behind and beneath sinks, lockers, tables, panels, and similar areas where insects hide or through which they may enter. Do not let spray or mist contact food, feedstuffs and water supplies. Dishes and food handling equipment should be thoroughly washed with soap and water if they have become contaminated with this product. Repeat treatment as needed for control.  
**WARNING** — Use as spot treatment only for indoor pests. Do not use in poultry houses or other animal buildings. Do not allow children or pets to contact treated surfaces until spray has dried. Remove or cover fish bowls before spraying.

**DIRECTIONS FOR APPLICATION WITHIN FOOD AREAS**

**FOOD AREAS APPLICATION LIMITED TO CRACK AND CREVICE TREATMENT ONLY**

**USE ONLY**

Apply only to cracks and crevices in walls, floors, baseboards, window and door frames, and other similar areas where insects hide or through which they may enter. Do not apply to surfaces that are in contact with food, feedstuffs, or water supplies. Do not apply to equipment used for food processing or handling unless the equipment is thoroughly cleaned and dry before application. Do not apply to food, feedstuffs, or water supplies. Do not apply to dishes, glasses, or other food handling equipment unless they have been thoroughly washed with soap and water. Do not apply to surfaces that are in contact with food, feedstuffs, or water supplies.

**PRECAUTIONARY STATEMENTS**  
**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**  
**DANGER**

This product is extremely toxic to humans and domestic animals. It may cause severe irritation to the eyes, nose, throat, and respiratory system. It may also cause skin irritation. Do not breathe dust or spray. Do not get in eyes, nose, mouth, or on skin. If it gets on skin, wash with soap and water. If it gets in eyes, wash with water for 15 minutes. If it gets in mouth, do not induce vomiting. Seek medical attention immediately.

**NOTE TO PHYSICIAN — WARNING SYMPTOMS**

Symptoms include weakness, headache, tightness in chest, blurred vision, non-reactive pin-point pupils, salivation, sweating, nausea, vomiting, diarrhea, and abdominal cramps. **TREATMENT** — Atropine is the specific therapeutic antagonist of choice against parasympathetic nervous stimulation. If there are signs of parasympathetic stimulation Atropine Sulfate should be injected at 10-minute intervals, in doses of 1 to 2 milligrams until complete atropinization has occurred. Morphine is contraindicated. 2-PAM is also antidotal and may be administered in conjunction with Atropine. Clear chest by postural drainage.  
 Repeated exposure to cholinesterase inhibitors may without warning cause prolonged susceptibility to very small doses of any cholinesterase inhibitor. Allow no further exposure until time for cholinesterase regeneration has been allowed as determined by a blood test.

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish, birds, and other wildlife. Do not apply directly to lakes, streams or ponds. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes.

**PHYSICAL OR CHEMICAL HAZARDS**

Combustible Mixture — Do not use, pour, spill, or store near heat or open flame.