

# P.T.O. DURSBAN - DDVP E.S.E.

INDOOR & OUTDOOR  
PEST CONTROL

ROACHES, ANTS, CRICKETS, FLEAS,  
SPIDERS, CHINCH BUGS,  
SOD WEBWORMS.

ORNAMENTAL INSECT CONTROL

Only For Sale To, Use And Storage By  
Pest Control Operators or Other Servicepersons

ACCEPTED  
APR 27 1984  
Listed in the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, and Registered under  
registered under  
EPA Reg No 3314-68



**ACTIVE INGREDIENTS:**

Chlorpyrifos [0,0-diethyl 0-(3,5,6-trichloro 2-pyridyl) phosphorothioate] .....	24.00%
2,2-Dichlorovinyl dimethyl phosphate .....	8.04%
Related Compounds .....	0.48%
Xylene Range Aromatic Solvent .....	62.42%
<b>INERT INGREDIENTS:</b> .....	<b>7.08%</b>
	<b>Total 100.00%</b>

Contains 2 lb. Chlorpyrifos & 5 lb. Dichlorvos (DDVP) per Gallon  
EPA REG. NO. 3314-68 EPA Est No 3314-FL-1



**KEEP OUT OF REACH OF CHILDREN  
DANGER — POISON**



See Side Panel For Additional Precautionary Statements  
**STATEMENT OF PRACTICAL TREATMENT**

**If Swallowed:** Call a physician or Poison Control Center immediately. Gastric lavage is indicated if material was taken internally. DO NOT INDUCE vomiting unless other treatment is not available. Vomiting may cause aspiration pneumonia. If it is necessary to induce vomiting, give victim one or two glasses of water and insert finger in back of throat. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious person.

**If Inhaled:** Remove victim to fresh air. Apply respiration if indicated.

**If On Skin:** Wash immediately with soap and water. Call a physician immediately.

**If In Eyes:** Flush eyes with plenty of water. Call a physician immediately.

**If Warning Symptoms Appear:** Keep patient prone and quiet. Start artificial respiration immediately if patient is not breathing. Transport patient immediately to the nearest physician.

Net Contents      Gallon

BEST DOCUMENT AVAILABLE

**BEST DOCUMENT AVAILABLE  
STORAGE AND DISPOSAL**

**Pesticide Disposal**

**Container Disposal**

**NOTICE**

**DIRECTIONS FOR USE**

**PESTS OUT OF DOORS**

**PESTS ON OUTSIDE SURFACES OF BUILDINGS**

**NOTE** — To help prevent infestations of buildings by Ants, Crickets, Earwigs, Millipedes, and Sawflies (Millipede) treat a band of soil 6 to 10 feet wide around and adjacent to buildings, also building's foundation to a height of 2 to 3 feet, where pests are active and may find entrance. Use 16 ounces (1 pint) to 100 gallons of water and apply as a coarse spray at the rate of about 10 gallons spray mixture per 1,000 sq. ft. to thoroughly and uniformly wet the soil area.

**DIRECTIONS FOR USE  
DOSAGES AND DIRECTIONS FOR USE ON TURF**

PEST	DOSAGE per 1,000 sq. ft.	REMARKS
------	--------------------------	---------

Bluegrass Billbug	3 to 6 Ounces	Spray early in the season when adults first appear. Retreat when needed.
-------------------	---------------	--

1.5 Gallons (1.5000) R. is killed disease

**DOGWOOD BORER** — Use 16 ounces per 100 gallons water to treat both green and grey bark and inner bark of trees and shrubs in groups. Contact your local Agricultural Extension Office for the proper timing.

**PEACH TREE BORER** — Use 16 ounces per 100 gallons water to spray the trunk of peach trees up to 12 1/2 feet from the ground and spray 3 times May 1st, June 1st, and July 1st. Do not spray the lower branches of the tree. This treatment is for preventive only.

**NOTE** — MOSQUITOES — Apply to wet areas treated for control of Turf and Ornamental plant pests to control for mosquitoes. Do this after treatment, depending on amount of insecticide used and conditions.

**AREA CONTROL OF TICKS AND CHIGLINS** — Use for control areas such as residential, recreational, and public picnic and camping sites, parks and other recreational areas where these pests are present and create a nuisance or possible public health problem. Apply 1 pint per acre to sufficient water to moist all areas including low undrained, grassy areas, woods, and ground surface and dilute using enough spray volume to do rain through coverage, usually 40 to 100 gallons of water per acre.



**COLONIAL PRODUCTS, INC.**  
1830 Tenth Avenue North  
Lake Worth, FL 33461

# BEST DOCUMENT AVAILABLE

## DIRECTIONS FOR USE

**ATTENTION:** Do not use in food areas. For use in food areas, use only in non-food areas as directed below.

### DOSAGES AND DIRECTIONS FOR USE ON ORNAMENTAL PLANTS

Pests	Amount of Concentrate to use in Water to Make
	3 Gal 10 Gal 100 Gal
Ants, Aphids, Beetles, Fleas, Flies, Gnats, Grubs, House Flies, Mosquitoes, Spiders, Ticks, and Weevils	1/2 ounce
Cockroaches, Fleas, Flies, Gnats, Grubs, House Flies, Mosquitoes, Spiders, Ticks, and Weevils	1/2 ounce
Cockroaches, Fleas, Flies, Gnats, Grubs, House Flies, Mosquitoes, Spiders, Ticks, and Weevils	1/2 ounce
Cockroaches, Fleas, Flies, Gnats, Grubs, House Flies, Mosquitoes, Spiders, Ticks, and Weevils	1/2 ounce
Cockroaches, Fleas, Flies, Gnats, Grubs, House Flies, Mosquitoes, Spiders, Ticks, and Weevils	1/2 ounce
Cockroaches, Fleas, Flies, Gnats, Grubs, House Flies, Mosquitoes, Spiders, Ticks, and Weevils	1/2 ounce
Cockroaches, Fleas, Flies, Gnats, Grubs, House Flies, Mosquitoes, Spiders, Ticks, and Weevils	1/2 ounce
Cockroaches, Fleas, Flies, Gnats, Grubs, House Flies, Mosquitoes, Spiders, Ticks, and Weevils	1/2 ounce
Cockroaches, Fleas, Flies, Gnats, Grubs, House Flies, Mosquitoes, Spiders, Ticks, and Weevils	1/2 ounce
Cockroaches, Fleas, Flies, Gnats, Grubs, House Flies, Mosquitoes, Spiders, Ticks, and Weevils	1/2 ounce

## DIRECTIONS FOR USE

### INDOOR PEST CONTROL

Do not use the product in edible product areas of food processing plants, restaurants or other areas where food is commercially prepared or processed. Do not use in serving areas where food is exposed. **USE ONLY IN NON-FOOD AREAS** as directed below. Use this product as a spot treatment for the control of Cockroaches, Ants, Fleas, Flies, Gnats, Grubs, House Flies, Mosquitoes, Spiders, Ticks, and Weevils in and around residential, institutional, and commercial buildings including homes, apartment buildings, hospitals, stores, and warehouses. It may be used with cats. It is a dead-end base oil for a maintenance spray or extra strength spray. Follow directions for control of pests listed and repeat as directed for control.

### DOSE CHART

Amount of Spray	Amount of Concentrate To Use
1 Gallon	1/2 ounce
5 Gallons	2 1/2 ounces
10 Gallons	5 ounces
25 Gallons	12 1/2 ounces

This product should be used as a maintenance spray until the pest population is reduced to a level which is satisfactory. For control of Cockroaches, Ants, Fleas, Flies, Gnats, Grubs, House Flies, Mosquitoes, Spiders, Ticks, and Weevils, use this product as a spot treatment. Apply as a spot spray or with a paint brush to areas where Cockroaches, Ants, Fleas, Flies, Gnats, Grubs, House Flies, Mosquitoes, Spiders, Ticks, and Weevils are normally found. Apply in sheltered areas, around water pipes, drains, basins, and beneath sinks, lockers, tables, cabinets, and other areas where insects hide or through which they may enter. Do not let spray get on food, foodstuffs, and water supplies. Clean and food handling equipment should be thoroughly washed with soap and water if they have become contaminated with this product. Repeat treatment as needed for control.

### DIRECTIONS FOR APPLICATION IN NON-FOOD AREA

Non-food areas include Garbage rooms, Lavatories, floor drains, the back of entry and vestibule doors, locker rooms, machine rooms, boiler rooms, garages, mop closets and storage areas (after canning or bottling). Use the "maintenance" rate from Dose Chart above as a spot treatment by spraying as a spot spray or with a paint brush to areas where Cockroaches, Ants, Fleas, Flies, Gnats, Grubs, House Flies, Mosquitoes, Spiders, Ticks, and Weevils are normally found. Apply in sheltered areas, around water pipes, drains, basins, and beneath sinks, lockers, tables, cabinets, and other areas where insects hide or through which they may enter. Do not let spray get on food, foodstuffs, and water supplies. Clean and food handling equipment should be thoroughly washed with soap and water if they have become contaminated with this product. Repeat treatment as needed for control.

**PRECAUTIONS:** — Use as spot treatment only for indoor pests. Do not use in poultry houses or other animal buildings. Do not allow children or pets to contact treated surfaces until spray has dried. Remove or cover fish ponds before spraying.

## DIRECTIONS FOR APPLICATION WITHIN FOOD AREAS

FOOD AREAS APPLICATION LIMITED TO CRACK AND CREVICE TREATMENT ONLY.

USE ONLY IN NON-FOOD AREAS as directed below. Use this product as a spot treatment for the control of Cockroaches, Ants, Fleas, Flies, Gnats, Grubs, House Flies, Mosquitoes, Spiders, Ticks, and Weevils in and around residential, institutional, and commercial buildings including homes, apartment buildings, hospitals, stores, and warehouses. It may be used with cats. It is a dead-end base oil for a maintenance spray or extra strength spray. Follow directions for control of pests listed and repeat as directed for control.

## PRECAUTIONARY STATEMENTS

**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**  
**DANGER**

## NOTE TO PHYSICIAN — WARNING SYMPTOMS

Symptoms include weakness, headache, tightness in chest, blurred vision, non-reactive pin-point pupils, salivation, sweating, nausea, vomiting, diarrhea, and abdominal cramps. **TREATMENT** — Atropine is the specific therapeutic antagonist of choice against parasympathetic nervous stimulation. If there are signs of parasympathetic stimulation Atropine Sulfate should be injected at 10-minute intervals, in doses of 1 to 2 milligrams until complete atropinization has occurred. Morphine is contraindicated. 2-PAM (2,2,6,6-tetramethylpiperidine) may be administered in large doses to relieve chest pain by postural changes. Repeat atropine treatments as necessary. The following state presumed susceptibility to very small doses of anticholinergic drugs. Advise the physician until safe for administration. Susceptible are postulated as determined by clinical test.

## ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, and other animals. Do not apply directly to lakes, streams or ponds. Do not apply when weather conditions favor drift spray. Do not contaminate water by cleaning equipment or disposal of wastes.

## PHYSICAL OR CHEMICAL HAZARDS

Corrosive Nature—Do not use, pour, spill, or otherwise test or open flames.