

SEP 14 1982
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, this product is registered under 3314-76

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION-May be harmful if swallowed. Do not take internally. Avoid eye & skin contact. Wash thoroughly after handling & before eating or smoking. Wash contaminated clothing before reuse. Keep away from food, feedstuffs, and domestic water supplies.

NOTE TO PHYSICIAN-Active Ingredient is a cholinesterase inhibitor. Treat symptomatically. Atropine is an antidote.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, pets, birds and other wildlife. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning or equipment or disposal of wastes. Apply this product only as recommended on this label.


STORAGE & DISPOSAL

Disposal-Do not reuse empty container. Wrap and put in trash collection.

NOTICE-Buyer assumes all risk and liability of use, storage and/or handling of this material not in accordance with the terms of this label.

DURE-BAIT FOR MOLE CRICKETS

**Ready to Use
INSECTICIDE BAIT**
For Control of Mole Crickets Infesting
LAWNS, GOLF COURSES AND ORNAMENTAL
TURF GRASS



ACTIVE INGREDIENTS
INERT INGREDIENTS

KEEP OUT OF REACH OF CHILDREN CAUTION

Statement Of Practical Treatment

IF SWALLOWED-Call a physician or Poison Control Center immediately. Drink one or two glasses of water and induce vomiting by touching the back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

IF ON SKIN-Remove contaminated clothing and wash affected areas with soap and water.

IF IN EYES-Flush eyes with plenty of water. Call a physician if irritation persists.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

MOLE CRICKETS in Lawns, Golf Courses and other Ornamental Turf - For control of Mole Crickets apply this granule product at the rate of 150 lbs. per acre or 3-1/2 lbs. per 1,000 sq. ft. Spread evenly over the area using a mechanical spreader. Apply in late afternoon on moist soil in areas known to be infested with these insects. Repeat application in about one week for best control.

THE BEST TIME TO CONTROL THE SOUTHERN MOLE CRICKET

Jan.	Feb.	Mar.	Apr.	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.

NOTE #2 indicates the best time to control the Adult Mole Crickets.

#1 indicates egg laying time during the summer.

EPA Reg. No. 3314-76 EPA Est. No. 3314-FL 1

NET WEIGHT POUNDS

COLONIAL PRODUCTS, INC.
1830 Tenth Avenue North
Lake Worth, Florida 33461

RW 8/82