

**ACCEPTED**  
 12/14/70  
 UNDER THE FEDERAL INSECTICIDE  
 FUNGICIDE AND RODENTICIDE ACT  
 REGISTERED  
 3325-47 SUBJECT  
 TO ALL REGULATIONS AND ORDINANCES.

**NEW...**

10% more  
product

Puff Pack

**hide**®  
 KILLER FORMULA

**MICE and RATS**

NET WEIGHT  
5½ oz.

CHILDREN. AVOID  
STUFFS. KEEP  
BACK LABEL  
SAFE.

**TWIST THE LID ...  
 SQUEEZE THE PACK ... HIDE "PUFFS" OUT  
 THE BETTER YOU HIDE IT—THE BETTER IT WORKS!**

**FOR MICE AND RATS**—Do not place in areas visible or accessible to children, pets or domestic animals. Apply HIDE in hiding places, along walls, behind objects, in path between object and wall in attic, basement, garage and outside buildings. Lay down a paper-thin layer of dust along rat or mouse runways. Heavily dust rat burrows and under rat or mouse infested buildings. In some cases it may be necessary to repeat dusting of heavily used runways or renew dust applications that have been removed by rain, etc. After completion of control program, clean up remaining powder. Control should be achieved in 10-14 days. Treatment should continue until all signs of activity have stopped. One package of 5½ ounces should cover approximately 14 square feet of rat runway.

**WARNING:** The active ingredient of this product is an anticoagulant chemical. Keep HIDE out of reach of children and pets, wash hands after using and avoid breathing dust. Avoid contamination of food, food preparation surfaces and cooking ware. If taken internally, contact physician immediately and retain HIDE package for reference to ingredients. Follow directions and avoid all danger. Do not use in feed or food processing plants, or in any location where the possibility exists of contaminating feed or food items.

**ANTIDOTE:** In case of ingestion by humans, induce vomiting by giving a tablespoon of salt in a glass of warm water and repeating until vomit fluid is clear. Call a physician immediately. Treatment by physician should include the administration of oral or intravenous injection of Vitamin K-1. In critical situations, a transfusion with fresh whole blood should be given.

**NOTICE:** Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

USDA REG. NO. 3325-47

**CLARENCE BOORD & SONS, INC.**  
 LEON • IOWA

ACCEPTED  
 12/14/70  
 UNDER THE  
 FOOD  
 AND  
 DRUG  
 ADMINISTRATION  
 3325-47  
 DIST. R.

**NEW...**

10% more  
 product

**hide**®

PMP  
 FORMULA

**MICE and RATS**

**ACTIVE INGREDIENTS:**

Calcium Phosphate	1.2
Iron Oxide	0.1%
Other Ingredients	17.7%
Inert Ingredients	81.0%

**NET WEIGHT**  
 5 1/2 oz.

**WARNING: KEEP OUT OF REACH OF CHILDREN. AVOID  
 CONTAMINATION OF FEED AND FOODSTUFFS. KEEP  
 AWAY FROM DOMESTIC ANIMALS. SEE BACK LABEL  
 FOR ADDITIONAL CAUTIONS.**

**THE BETTER YOU HIDE IT—THE BETTER IT WORKS!**

**FOR MICE AND RATS**—Do not place in areas visible or accessible to children, pets or domestic animals. Apply HIDE in hiding places, along walls, behind objects, in path between object and wall in attic, basement, garage and outside buildings. Lay down a paper-thin layer of dust along rat or mouse runways. Heavily dust rat burrows and under rat or mouse infested buildings. In some cases it may be necessary to repeat dusting of heavily used runways or renew dust applications that have been removed by rain, etc. After completion of control program, clean up remaining powder. Control should be achieved in 10-14 days. Treatment should continue until all signs of activity have stopped. One package of 5 1/2 ounces should cover approximately 14 square feet of rat runway.

**WARNING:** The active ingredient of this product is an anticoagulant chemical. Keep HIDE out of reach of children and pets, wash hands after using and avoid breathing dust. Avoid contamination of food, food preparation surfaces and cooking ware. If taken internally, contact physician immediately and retain HIDE package for reference to ingredients. Follow directions and avoid all danger. Do not use in feed or food processing plants, or in any location where the possibility exists of contaminating feed or food items.

**ANTIDOTE:** In case of ingestion by humans, induce vomiting by giving a tablespoon of salt in a glass of warm water and repeating until vomit fluid is clear. Call a physician immediately. Treatment by physician should include the administration of oral or intravenous injection of Vitamin K-1. In critical situations, a transfusion with fresh whole blood should be given.

**NOTICE:** Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

USDA REG. NO. 3325-47

**CLARENCE BOORD & SONS, INC.**  
 LEON • IOWA