15% FERMATE* DUST

(FERBAM)

FOR PREVENTION OF TOBACCO BLUE MOLD, DOWNY MILDEN OF CABBAGE AND BROCCOLI PLANT BEDS, AND BLACK SPOT AND RUST ON ROSES.

ACTIVE INGREDIENT:

Ferbam (Ferric Dimethyl Dithiocarbamate) 11.4% INERT INGREDIENTS: 88.6%

TOTAL 100.00%

READ ENTIRE LABEL. USE IN STRICT ACCORDANCE WITH LABEL DIRECTIONS AND CAUTIONS.

DIRECTIONS FOR USE

For Tobacco Blue Mold control start dusting when plants are about the size of the or when Blue Mold first appears in the area, and dust regularly twice a week at the following rates per 100 square yards.

First to fourth dusting 1 to 1½ lbs., fifth to sixth 2 to 2½ lbs. seventh & superquent 3 to 3½ lbs. Repeat dusting immediately following rain.

For Downy Mildew of Cabbage and Broccoli Plant beds start dusting when first signs of Downy Mildew appears. Dust regularly 2 to 3 times a week, using sufficient material to thoroughly cover plants. Repeat as necessary.

For preventions of Black Spot and Rust on roses begin application when first leaves unfold. Dust regularly 2 to 3 times a week, using sufficient material to thoroughly cover plants. Repeat as necessary.

NOTICE: Our recommendation for use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice. The buyer must assume all responsibility, including injury or damage, resulting from its misuse as such, or in combination with other materials.

CAUTION KEEP OUT OF REACH OF CHILDREN

CAUTION: Harmful if swallowed or inhaled". Avoid breathing of dust. Do not get in eyes, on skin or clothing. May cause irritation of nose, throat and skin. In case of contact, wash with plenty of water. For eyes get medical attention.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes.

CONTAINER DISPOSAL

Destroy this Container: Do not Re-Use. Destroy it by burying with waste or burning. Stay away from smoke or tumes.

Apply this product only as specified on this label.

U.S.D.A. Reg. 3342-7

15% FERMATE DUST

(FERBAM)

FOR PREVENTION OF TOBACCO BLUE MOLD, DOWNY MILDEY OF CABBAGE AND BROCCOLI PLANT BEDS, AND BLACK SPOT AND RUST ON ROSES.

ACTIVE INGREDIENT:

TOTAL 100.00%

READ ENTIRE LABEL USE LASTRICA ACCORDA ACLASTH LABEL DIRECTIONS AND CALHONS

DIRECTIONS FOR USE

For Tobacco Blue Mold control start dusting when plants are about the size of a dime or when Blue Mold first appears in the area, and dust regularly twice a week at the following rates per 100 square yards.

First to fourth dusting 1 to 1½ lbs., fifth to sixth 2 to 2½ lbs. seventh & subsequent 3 to 3½ lbs. Repeat dusting immediately following rain.

For Downy Mildew of Cabbage and Broccoli Plant beds start dusting when first signs of Downy Mildew appears. Dust regularly 2 to 3 times a week, using sufficient material to thoroughly cover plants. Repeat as necessary.

For preventions of Black Spot and Rust on roses begin application when first leaves unfold. Dust regularly 2 to 3 times a week, using sufficient material to thoroughly cover plants. Repeat as necessary.

NOTICE: Our recommendation for use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice. The buyer must assume all responsibility, including injury or damage, resulting from its misuse as such, or in combination with other materials.

CAUTION KEEP OUT OF REACH OF CHILDREN

ANCHON CONTRACTOR AND ANCHOR AND ANCHOR OF THE CONTRACTOR AND ANCHOR OF TH

* ONIMINER DEPOSAL

And the state of the second of