Best Available Copy

4% MALATHION DUST

ACTIVE INGREDIENTS

Malathion *O. O-dimethyl dithlophosphate of dlethyl mercaptosuccinate

INERT INGREDIENTS (Pyrophyllite)

TOTAL 100%

DIRECTIONS

BEANS: For control of Mexican Bean Beetle, Bean Leaf Hopper, Spider Mites, Thrips, and Aphids, copply at the rate of 80-35 pounds per acre For best results apply to underside of leaves. Do not apply to beans within 3 days of harvest.

CABBAGE: For control of Aphids apply at the rate of 80-35 pounds per acre. Do not apply within 7 days of harvest.

NOTICE: Our recommendation for use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice. The buyer must assume all responsibility, including injury or damage, resulting from its misuse as such, or in combination with other materials.

"This product is highly toxic to bees exposed to direct treatment or residue on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service."

Harmful if swallowed! Avoid breathing of dust, may cause irritation of nose & throat. Do not set interest, on skin or clothing. Wash thoroughly after handling, change contaminated clothes daily. Avoid contamination of feed and food stuff. Do not feed treated forage to dairy animals or to livestock being finished for slaughter.

This product will kill Fish the

Keep out of Lakes, Streams and Ponds-Bestroy this Container. Do not ResuseDestroy it by Burying with waste or burning. Stay away from smoke or fumes.

Apply this product only as specified on this label.

U.S.D.A. Reg. No. 3342-14

JUL 2 2 1970

UNDER THE FOR ECONOMIC LISTON REGISTERED UNDER NOW AND STEEL BOTTON ATTACHED CONTENTS.

