

5% SEVIN DUST

ACTIVE INGREDIENTS:

Carbaryl (1-naphthyl N - methylcarbamate) 5.0%

INERT INGREDIENTS 95.0%

TOTAL 100.00%

Read Entire Label. Use in Strict Accordance With Label Directions and Cautions.

DIRECTIONS FOR USE

BEANS: (Green beans, lima beans, snap beans, cowpeas and black eyed peas). Use 10 lbs. per acre for Mexican Bean Beetle; use 20 lbs. per acre for leaf beetle, cucumber beetles, flea beetles, Japanese beetle, leafhoppers, velvet bean caterpillar and western bean cutworm. Use 20 to 30 lbs. per acre for armyworms, corn earworm, stink bugs and tarnished plant bug. Use up to the day of harvest. **CORN:** Use 20 to 40 lbs. per acre for corn earworm, corn rootworm adults, European corn borer, fall armyworm, flea beetles, Japanese beetle, sap beetles and leafhoppers. For larvae in whorl (budworm damage) and foliage feeders, apply to entire plant. Repeat as necessary. For insects attacking silks and ears, apply at 2 to 3 day intervals starting when silks appear and continuing until silks begin to dry. Three or more applications may be required depending on severity of infestation. Timing and good coverage are absolutely essential for effective control. Use up to the day of harvest.

CABBAGE, BROCCOLI, BRUSSEL SPROUTS, CAULIFLOWER AND KOHLRABI, CUCUMBERS, MELONS, PUMPKIN, SQUASH, HEAD LETTUCE, CARROTS, POTATO, TOMATO, EGGPLANT, PEPPER, AND OKRA: Use 10 to 20 lbs. per acre for pickleworm and melon worm, for flea beetles and harlequin bug. 20 to 30 lbs. per acre for cucumber beetles, leafhoppers, squash bugs, cabbage caterpillars, armyworms, & corn earworm. Avoid excessive applications. Some leaf injury may occur if treatments are made when tender foliage is wet or in the presence of prolonged high humidity.

CAUTION: Do not apply within three days of harvest on Cabbage, broccoli, Brussel sprouts, cauliflower, Kohlrabi and head lettuce. **CHICKEN MITES** (1 lb. 5 per cent Dust) per 100 birds. 1 lb. (5 per cent Dust) per 40 sq. ft. Litter. **POULTRY:** "Do not apply to poultry within 7 days of slaughter." Do not repeat application within 4 weeks. "Do not contaminate feed or drinking water."

TOBACCO: Flea Beetles & Japanese Beetle 16 to 20 lb. per acre. Good coverage on underside of leaves over entire plant, especially lower leaves, is essential. **HORNWORMS:** 16 to 20 lb. per acre. Apply insecticide especially to top 6 leaves and avoid unduly heavy application. Large worms cause more damage and are harder to kill than small worms. Treat infested fields before largest worms exceed 1½ inches in length.

GRASSHOPPERS: 16 lb. per acre. Apply dust evenly to tobacco foliage and to barrier strips 15 to 30 ft. wide around tobacco. Do not graze or feed animals on treated barrier strips for 7 days following treatment.

NOTE: If late applications are necessary, treat immediately after priming rather than before priming. Treated fields may be entered immediately after dust has settled. Some phytotoxicity may occur on tender foliage in the presence of rain or high humidity of several days duration following application.

NOTICE: Our recommendation for use of this product are based upon test believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice. The buyer must assume all responsibility, including injury or damage, resulting from its misuse as such or in combination with other materials.

CAUTION: This product is highly toxic to fish and aquatic life. Do not apply to water bodies. Do not apply to areas where water is present. Do not apply to areas where water is present.

CAUTION:

KEEP OUT OF REACH OF CHILDREN & ANIMALS

"Fatal if swallowed or inhaled." Avoid breathing of dust. Do not get in eyes, on skin or clothing. Wash thoroughly after handling. Change contaminated clothes daily. Avoid contamination of feed and feed stuff. Do not feed treated forage to dairy animals or to livestock being finished for slaughter.

Destroy this Container: Do not Re-Use. Destroy it by burying with waste or burning. Stay away from smoke or fumes. Apply this product only as directed on this label.

ACCEPTED

5% SEVIN DUST

ACTIVE INGREDIENTS:

Carbaryl (1-naphthyl N - methylcarbamate) 5.0%

INERT INGREDIENTS 95.0%

TOTAL 100.00%

Read Entire Label. Use in Strict Accordance With Label Directions and Cautions.

DIRECTIONS FOR USE

BEANS: (Green beans, lima beans, snap beans, cowpeas and black-eyed peas) Use 10 lbs. per acre for Mexican bean Beetle; use 20 lbs. per acre for leaf beetle, cucumber beetles, flea beetles, Japanese beetle, leafhoppers, velvet bean caterpillar and western bean cutworm; use 20 to 30 lbs. per acre for armyworms, corn earworm, stink bugs and tarnished plant bug. Use up to the day of harvest. **CORN:** Use 20 to 40 lbs. per acre for corn earworm, corn rootworm adults, European corn borer, fall armyworm, flea beetles, Japanese beetle, sap beetle and leafhoppers. For larvae in whorl (budworm damage) and foliage feeders, apply to entire plant. Repeat as necessary. For insects attacking silks and ears, apply at 2 to 3 day intervals starting when silks appear and continuing until silks begin to dry. Three or more applications may be required depending on severity of infestation. Timing and good coverage are absolutely essential for effective control. Use up to the day of harvest.

CABBAGE, BROCCOLI, BRUSSEL SPROUTS, CAULIFLOWER AND KOHLRABI, CUCUMBERS, MELONS, PUMPKIN, SQUASH, HEAD LETTUCE, CARROTS, POTATO, TOMATO, EGGPLANT, PEPPER AND OKRA: Use 10 to 20 lbs. per acre for pickleworm and melon worm; for flea beetles and harlequin bug, 20 to 30 lbs. per acre; for cucumber beetles, leafhoppers, squash bug, blight caterpillars, armyworms, & corn earworm. Avoid excessive applications. Some leaf injury may occur if treatments are made when the foliage is wet or in the presence of prolonged high humidity.

CAUTION: Do not apply within three days of harvest on Cabbage, broccoli, Brussels sprouts, cauliflower, Kohlrabi and head lettuce. **CHICKEN MITES:** Use 5 to 10 lbs. per acre on turkeys, ducks, geese, and guinea fow. Do not apply to poultry within 7 days of slaughter. Do not repeat application within 4 weeks. Do not contaminate feed or drinking water.

TOBACCO: Flea beetle, Japanese Beetle, and tobacco beetle. Use 20 to 30 lbs. per acre. Apply to entire plant. Repeat as necessary. Apply especially to top 6 leaves and to bottom 6 leaves. Apply to entire plant when damage is done. Do not apply to kill them, small worms. Treat entire field before worms exceed 1/2 inches in length.

HORNWORMS: Use 20 to 30 lbs. per acre. Apply to entire plant, especially to top 6 leaves and to bottom 6 leaves. Apply to entire plant when damage is done. Do not apply to kill them, small worms. Treat entire field before worms exceed 1/2 inches in length.

GRASSHOPPERS: Use 20 to 30 lbs. per acre. Apply to entire plant. Do not apply to beets. Treat 15 to 20 ft. wide around tobacco. Do not graze or feed animal on treated barrier strips for 7 days following treatment.

NOTE: If late applications are necessary, treat immediately after priming rather than before priming. Treated fields may be entered immediately after dust has settled. Some phytotoxicity may occur on tender foliage in the presence of rain or high humidity of several days duration following application.

NOTICE: This is a recommendation for use of this product. It is based on tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice. The buyer must assume all responsibility, including injury or damage, resulting from its misuse as such or in combination with other materials.

ACCEPTED