

5-13-71
3342-56

SEVIN VEGETABLE DUST

ACTIVE INGREDIENTS (Carbaryl (1-Naphthyl N-Methylcarbamate)	1.75%
INERT INGREDIENTS	98.25%
Total 100.00%	

Read Entire Label. Use in Strict Accordance With Label Directions and Cautions.

DIRECTIONS FOR USE

BEANS: To control Mexican Bean Beetle apply at the rate of 25 to 30 pounds per acre or 2 ounces per 50 feet of row. Make the first application when insects or egg masses on the underside of the leaves are first seen. Thorough application to the lower as well as the upper leaf surface is necessary for good results. Repeat applications at 7 to 10 day intervals as necessary for control.

NOTICE: Our recommendation for use of this product are based upon test believed to be reliable. The use of this product being beyond the control of manufacturer no gurantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice. The buyer must assume all responsibilty, including injury or damage, resulting from it misuse as such, or in combination with other materials.

"This product is highly toxic to bees exposed to direct treatment or residue on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service."

CAUTION: Keep Out Of Reach Of Children!

"Harmful if swallowed or inhaled." Avoid Breathing of dust. Do not get in eyes, on skin or clothing. Wash thoroughly after handling, change contaminated clothes daily. Avoid contamination of feed and food stuff. Do not feed treated forage to dairy animals or to livestock being finished for slaughter.

Destroy this Container Do not Re Use Destroy it by burying with waste or burning. Stay away from smoke or fumes. Apply this product only as specified on this label.

USDA Reg. 3342-56

5-13 41
3342-56

SEVIN VEGETABLE DUST

ACTIVE INGREDIENTS (Carbaryl (1-Naphthyl N-Methylcarbamate)	1.75%
INERT INGREDIENTS	98.25%
Total 100.00%	

Read Entire Label. Use in Strict Accordance With Label Directions and Cautions.

DIRECTIONS FOR USE

BEANS: To control Mexican Bean Beetle apply at the rate of 25 to 30 pounds per acre or 2 ounces per 50 feet of row. Make the first application when insects or egg masses on the underside of the leaves are first seen. Thorough application to the lower as well as the upper leaf surface is necessary for good results. Repeat applications at 7 to 10 day intervals as necessary for control.

NOTICE: Our recommendation for use of this product are based upon test believed to be reliable. The use of this product being beyond the control of manufacturer no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice. The buyer must assume all responsibility, including injury or damage, resulting from a misuse as such, or in combination with other materials.