DIRECTIONS FOR USE

This material is a concentrated liquid formulation containing 4 pounds of parathion per gallon. One pint of this material contains the same amount of parathion as 3-1/3 pounds of a 15% wettable formulation.

BEANS: For Aphids, Mexican Bean Beetle and Potato Leafhopper use 1 pint in sufficient water to treat one acre.

CABBAGE, BROCCOLI, BRUSSELS SPROUTS, COLLARDS, KALE MUSTARD For Aphids, Serpentine Leafminer. Caterpillars and Vegetable Weevil use 1 pint in sufficient water to treat one acre.

CELERY: For Aphids, Celery Leaf Tier and Spider Mites use 1 pint in sufficient water to treat one acre.

CUCUMBERS, MELONS AND SQUASH: For Aphids, Cucumber Beetle, Leafminers, Spider Mites, Melonworms, Pickleworms and Squash Bugs use 1/2 to 1 pint in sufficient water to treat one acre.


OKRA: For Aphids, Okra Caterpillar, Serpentine Leafminer, Blister Beetles and Stink Bugs use 1 pint in sufficient water to treat one acre.

PEPPER: For Aphids and Thrips use 1 pint in sufficient water to treat one acre POTATOES: For Aphids, Blister Beetles, Leafhoppers, Leafminers, Plant Bugs. Thrips and Vegetable Weevils use 1 pint in sufficient water to treat one acre TOMATOES: For Aphids, Leafminers, Spider Mites and Tomato Russet Mites use 1 pint in sufficient water to treat one acre.

COTTON: For Cotton Aphid, Spider Mites, Stink Bugs, cotton Leaf Perforator and Southern Garden Leafhopper use 1 pint in sufficient water to treat one acre. For Cotton Leafworm use 1/2 pint in sufficient water to treat one acre. Do not apply within 5 days of hand picking.

Observe all cautions listed under individual crops in addition to those listed below.

Do not apply within 7 days of harvest on broccoli, brussels sprouts and melons, within 10 days of harvest on cabbage, collards, kale, mustard and tomatoes; within 15 days of harvest on beans, cucumbers, peppers and squash; within 21 days of harvest on celery and okra; within 5 days of harvest on potatoes. Do not apply before vining on cucumbers.



TIGER & BRA PARATHION 4 EMULS

ACTIVE INGREDIENTS:

*Parathion (*O, O-diethyl O-p-nitrophenyl thiophosphate.)
Aromatic petroleum solvent
INERT INGREDIENTS

Contains 4.0 pounds Parathion per gallon.



DANGER KEEP OUT OF REACH OF CHIL

POISON

See side panel for Antidote Statement and additional caution FIRST AID TREATMENT

ANTIDOTE: If swallowed, Give a tablespoonful of salt in a glass of warm water and revictim lie down and keep quiet. Call a Physician Immediately! If on skin, In case of continumediately wash skin with soap and water.

WARNING: POISONOUS if Swallowed, Inhaled or Absorbed Through Skin! Rapidly Ah eyes, on skin or on clothing. Wear natural rubber gloves, protective clothing and goggles, with soap and water. Wear a mask or a respirator of a type passed by the U.S. Departn tection. Keep all unprotected persons out of operating areas or vicinity where there may h not be re-entered until drifting insecticide and volatile residues have dissipated. Do not a hands, arms and face thoroughly with soap and water before eating or smoking. Wash all hot water before re-use.

"THIS PRODUCT IS TOXIC TO FISH AND WILDLIFE." "KEEP OUT OF ANY BO ON TREATED AREAS MAY BE KILLED." DO NOT CONTAMINATE WATER BY CL POSAL OF WASTES," "APPLY THIS PRODUCT ONLY AS SPECIFIED ON THIS

MANUFACTURED IN U. S. A. BY CAPE FEAR CHEMICALS, INC.

NET CONTENTS 1 GALLON Manufacture/Seller makes no warranty express or implied covering this product. use whether in accordance with directions or not.

DIRECTIONS FOR USE

This material is a concentrated liquid formulation containing 4 pounds of parathlen per gallon. One pint of this material contains the same amount of parathlen as 3-1-3 pounds of a 15% wettable formulation.

BEANS: For Aphids, Mexican Bean Beetle and Potato Leafhopper use 1 pint in sufficient water to treat one acre.

CABBAGE, BROCCOLI, BRUSSELS SPROUTS, COLLARDS, KALE MUSTARD For Aphids, Serpentine Leafminer, Caterpillars and Vegetable Weevil use 1 pint in sufficient water to treat one acre.

CELERY: For Aphids, Celery Leaf Tier and Spider Mites use 1 pint in sufficicut water to treat one acre.

CUCUMBERS, MELONS AND SQUASH: For Aphids, Cucumber Beetle, Leafminers, Spider Mites, Melonworms, Pickleworms and Squash Bugs use 1/2 to 1 pint in sufficient water to treat one acre.

GERA: For Aphids, Okra Caterpillar, Serpentine Leafminer, Blister Beetles and Stink Bugs use 1 pint in sufficient water to treat one acre.

PEPPER: For Aphids and Thrips use 1 pint in sufficient water to treat one acre POTATOES: For Aphids, Blaster Beetles, Leafhoppers. Leafminers, Plant Bugs. Thrips and Vegetable Weevils use 1 pint in sufficient water to treat one acre TOMATOES: For Aphids, Leafminers, Spider Mites and Tomato Russet Mites use 1 pint in sufficient water to treat one acre.

COTTON: For Cotton Aphid, Spider Mites, Stink Bugs, cotton Leaf Perforator and Southern Garden Leafhopper use 1 pint in sufficient water to treat one acre. For Cotton Leafworm use 1 2 pint in sufficient water to treat one acre. Do not copyly within 5 days of hand picking.

Observe all cautions listed under individual crops in addition to those listed

Do not apply with a 7-days of harvest on broccoli, brussels sprouts and melons, within 10-days of harvest on cabbage, collards, kale, mustard and tomatoes; within 15-days of harvest on beans, cucumbers, peppers and squash; within 21-days of harvest on celery and okra; within 5-days of harvest on potatoes. Do not apply before vining on cucumbers.

TIGER BRA PARATHION 4 EMULS

ACTIVE INGREDIENTS:

*Parathion (*O, O-diethyl O-p-nitrophenyl thiophosphate.)

Aromatic petroleum solvent
INERT INGREDIENTS

Contains 4.0 pounds Parathion per gallon.

MANUFACTURED IN U. S. A. BY CAPE FEAR CHEMICALS, INC.

NET CONTENTS 1 GALLON

Manufacture Seler makes no warranty express or unblied covering this product, use whether in accordance with directions or not

IGER BRAND ATHION 4 EMULSIABLE

iitrophenyl	thiophosphate.)		47%
		****	46%
			7%
gallon.			100%

DANGER KEEP OUT OF REACH OF CHILDREN!



le panel for Antidote Statement and additional caution statements.
FIRST AID FREATMENT

ablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Have La Physician Immediately! If on skin. In case of contact, remove contaminated clothing and nd water.

wed, Inhaled or Absorbed Through Skin! Rapidly Absorbed. Through Skin! Do not get in natural rubber gloves, protective elething and goggles. In case of contact wash immediately or a respirator of a type passed by the U.S. Department of Agriculture for parathion proportion of operating areas or vicinity where there may be danger of drift. Vacated areas should rticide and volatile residues have dissipated. Do not contaminate feed and foodstuffs. Wash the soap and water before cating or smoking. Wash all contaminated clothing with soap and

TSH AND WILDLIFE." "KEEP OUT OF ANY BODY OF WATER." "BIRD FEEDING KILLED." DO NOT CONTAMINATE WATER BY CLEANING OF EQUIPMENT, OR DIS-THIS PRODUCT ONLY AS SPECIFIED ON THIS LABEL."

U. S. A. BY CAPE FEAR CHEMICALS, INC. Elizabethtown, N. C.

NET CONTENTS 1 GALLON

kes no warranty express or implied covering this product. User assumes all risk for
ince with directions or not.

USDA Reg. No. 3342

soil treatment: For the control of wireworms and other soil insects attacking roots of beans, carrots, corn, cabbage, cauliflower, lettuce, Irish potatoes, tobacco and ornamentals, use 1 to 2 quarts in 100 gallons of water per acre as a broadcast preplanting treatment. The lower rate of application should be used on mineral soils while much soils will require the higher dosage. After the material is evenly distributed over the soil surface, it should be thoroughly worked into the top six to nine inches of soil with a double disk harrow. Application should be made at least one to two weeks before planting. Keep all persons out of treated area for 48 hours.

ATROPINE IS ANTIDOTAL

Keep on hand an emergency supply of 20 tablets, grains 1 100. In case of Poisoning take 3 tablets (gr. 1 100.) and call physician at once.

PHYSICIAN: Enough atropine must be given in the shortest practical time to produce dilation of pupils. Patient must be under constant supervision and additional atropine may be necessary without delay. Clear chest by postural drainage. Use artificial respiration and oxygen if patient becomes comatose. Use of airway may be necessary.

USAGE CAUTIONS

USE OF THIS MATERIAL IN A MANNER OR AT A TIME OTHER THAN IN ACCORDANCE WITH DIRECTIONS MAY PRODUCE PLANT INJURY, EXCESSIVE RESIDUES OR OTHER UNDESIRABLE RESULTS.

DISPOSAL WARNING!

Do Not Re-Use The Container. Completely Empty The Contents and Bury the Unused Chemical at least 18 inches deep in an isolated location away from water supplies.

Break the container and bury the pieces at least 18 inches deep in an isolated location from water supplies.

BRAND IGER ATHION 4 EMULSIABLE

nitrophenyl thiophosphate.)

gallon.

DANI

CAPE FEAR CHEMICALS, INC. Elizabeth town, N. C.

kes no warranty express or implied covering this product. User assumes all risk for

USDA Reg. No. 3342

17

4677,

100%

USAGE CAUTIONS

USE OF THIS MATERIAL IN A MANNER OR AT A TIME OTHER THAN IN ACCORDANCE WITH DIRECTIONS MAY PRODUCE PLANT INJURY, EXCESSIVE RESIDUES OR OTHER UNDESIRABLE RESULTS.

DEALERS SHOULD SELL IN ORIGINAL PACKAGES ONLY.

DISPOSAL WARNING!

Do Not Re-Use The Container. Completely Empty The Contents and Bury the Unused Chemical at least 18 inches deep in an isolated location away from water supplies.

Break the container and bury the pieces at least 18 inches deep in an isolated location from water supplies.