

BEST AVAILABLE COPY

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS &
DOMESTIC ANIMALS
CAUTION**

Harmful if swallowed. Do not breathe dust. Do not get on skin. In case of contact, wash immediately with soap and water. Do not contaminate feed or foodstuffs. Keep out of reach of children.

In loading operations wear a mask or respirator jointly approved by the Mining Enforcement and Safety Administration (formerly U.S. Bureau of Mines) and by the National Institutes for Occupational Safety and Health under the provisions of 30 CER Part 1.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and wildlife. Birds feeding on treated areas may be harmed. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

**PHYSICAL OR CHEMICAL
HAZARDS**

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME

RE-ENTRY STATEMENT


Do not use pesticide in area of unprotected humans or domestic animals. Do not allow entry to area treated until the chemical has completely dried. Keep children and pets off treated area until chemical has been washed into the soil by watering or rain and area is completely dry.

STORAGE AND DISPOSAL

1. **PROHIBITIONS** Do Not Contaminate Water, Food, or Feed by Storage or Disposal. Open Dumping is Prohibited.
2. **PESTICIDE DISPOSAL** Pesticide, Spray Mixture, or Rinse that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies.
3. **CONTAINER DISPOSAL** Dispose of in an incinerator or landfill approved for pesticide containers, or bury in a safe place.
4. **GENERAL** Consult Federal, State or Local disposal authorities for approved alternative procedures such as limited open burning.

**10%
Toxaphene
Dust**

ACTIVE INGREDIENTS
Toxaphene
INERT INGREDIENTS
*TECHNICAL CHLORINATED CAMPHENE: CHLORINE
CONTENTS 67% to 69%



KEEP OUT OF REACH OF CHILDREN

DANGER

POISON



Statement For Practical Treatment

IF SWALLOWED Drink one or two glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention immediately.
IF INHALED Remove victim to fresh air and apply respiration if indicated.
IF ON SKIN Wash promptly with soap and water. Rinse thoroughly.
IN EYES Rinse eyes for at least 15 minutes with water and call a physician immediately.

Manufactured By
Cape Fear Chemicals Inc.
ELIZABETHTOWN, NORTH CAROLINA
NET CONTENTS — 25 LBS.

EPA EST NO. 3342-NC-1

EPA Reg. No. 3342-84

DIRECTIONS FOR USE

"It is a violation of Federal law to use this product in a manner inconsistent with its labeling."

WHEAT, BARLEY, OATS AND RYE: To control Climbing and Surface-Feeding Cutworms, Army Worms, Grasshoppers and Chinch Bugs, apply at the rate of 20 lbs. per acre. Do not apply more than once after heads have begun to form. Do not apply within one week of harvest of Wheat, Rye and Oats, within 2 weeks of harvest of Barley. Do not pasture dairy animals or meat animals in treated fields.

CABBAGE, BROCCOLI, LETTUCE AND BRUSSELS SPROUTS: To control Cabbage Loopers, apply at the rate of 20 lbs. per acre. For Cutworms (soil), apply 40 lbs. per acre to soil surface. For Cabbage and Head Lettuce. Do not apply after the Heads begin to form. For Leaf Lettuce, do not apply after seeding state. For Broccoli and Brussels Sprouts, do not apply after the edible parts begin to form. On all the above, do not apply this product if leaves, stems or heads are to be fed to dairy animals or meat animals. Do not apply within 30 days of harvest.

COLLARDS AND KALE: To control Army Worm, Cabbage Worm, Cabbage Looper, Climbing and Surface Feeding Cutworms, apply at the rate of 20 to 40 lbs. per acre. Do not apply more than 4 times nor more than 3 times after thinning. Do not apply within 21 days of harvest when 40 lbs. per acre is used. "Do not apply within 30 days of harvest."

SPINACH: To control Army Worm, Cabbage Worm, Cabbage Looper, Climbing and Surface Feeding Cutworms, apply only one time at the rate of 20 lbs. per acre. Do not apply within 21 days of harvest.

BEANS: To control Cutworms (soil) apply 20 lbs. per acre on soil in late afternoon. Do not apply after pods begin to form. 7 day Pre-Harvest Interval. "Do not feed treated vines to dairy animals or animals being finished for slaughter."

TOMATOES: To control Tomato Hornworm, Tomato Fruit Worm, Serpentine Leaf Miner, Flea Beetles, Cutworms, Cabbage Looper, Russet Mites and Colorado Potato Beetles, Army Worms and related insects, apply at the rate of 40 lbs. to the acres. Do not feed treated forage to dairy animals or animals being finished for slaughter. "Do not apply within 21 days of harvest."

PEANUTS: To control Fall Army Worm and Velvet Bean Caterpillar, apply at the rate of 15 to 20 lbs. per acre. "Do not feed foliage to dairy animals or animals being finished for slaughter."

CORN: To control Army Worm and Fall Army Worm, apply at the rate of 10 lbs. per acre. Do not feed treated plants to dairy or meat animals.

SORGHUM: To control Army worm and Fall Army Worm apply at the rate of 30 lbs. per acre. Do not feed treated plants or ensilage from treated plants to dairy or meat animals. Do not use on Sorghum to be sold commercially or to be shipped interstate. Do not apply within 40 days of harvest or apply after heads have begun to form.

WARRANTY STATEMENT

Our recommendation for use of this product are based on tests believed to be reliable. The use of this product is beyond the control of the manufacturer and no warranty is expressed or implied as to the effects or results of the results to be obtained if not used in accordance with the directions or established safe practice. The buyer must assume all responsibility, including injury or damage resulting from its misuse as such, or in combination with other materials.