

SUPER KILL

(Contains 4% Thiodan)

Oct 18, 1974

ACTIVE INGREDIENT
Endosulfan (Hexachlorohexahydromethano-2,4,3-benzodioxathiepin oxide).....4.00%
INERT INGREDIENTS:.....96.00%
100.00%

CAUTION: DEALERS MUST SEE IN ORIGINAL PACKAGES ONLY

DIRECTIONS FOR USE

CAUTION: READ ENTIRE LABEL USE IN STRICT ACCORDANCE WITH LABEL DIRECTIONS AND CAUTIONS.

3342-94
Comments

Apply when insects first appear and repeat as required to maintain effective control. Coverage of both upper and lower leaf surfaces is essential for good control. Observe use limitations. If insect control is required beyond the following use patterns, supplement the control program with other suitable pesticides. Observe days interval between last application and harvest indicated by number in () following crop.

COLLARDS (21), BROCCOLI (7), CABBAGE (7): Cabbage Aphid, Cabbage Looper, Cross-Striped Cabbageworm, Diamond Back Moth Larvae, Flea Beetle, Harlequin Bug, Imported Cabbageworm, Stink Bug - 3 1/2 ozs. per 500 sq. ft.

APPLES (30): Apple Aphid - 7 ozs. per 500 sq. ft. Unlimited number of applications. Do not feed pomace to livestock.

APPLES (21): Apple Aphid - 7 ozs. per 500 sq. ft. Do not make more than 3 applications during the fruiting period. Do not feed pomace to livestock.

APRICOTS (30), NECTARINES (30), PEACHES (30): Green Peach Aphid - 7 ozs. per 500 sq. ft. Do not make more than 2 applications during the fruiting period.

ARTICHOKES (7): Artichoke Aphid, Artichoke Plume Moth, Green Peach Aphid - 5 ozs. per 500 sq. ft.

BEANS (DRY BEANS ONLY) (3): Bean Aphid, Bean Leaf Skeletonizer, Mexican Bean Beetle - 4 ozs. per 500 sq. ft. Cucumber Beetle - 3 ozs. per 500 sq. ft. Do not apply more than 3 times per season. Do not apply within three days of harvest. Do not feed treated threshings to livestock or allow to graze in treated fields. Do not use on Lima Beans that are to be removed from the field for processing. Do not feed to livestock.

BEANS (SUCCULENT): Cucumber Beetle - 3 ozs. per 500 sq. ft. Mexican Bean Beetle - 3 1/2 ozs. per 500 sq. ft. Do not apply more than three times per season. Do not apply within three days of harvest. Do not feed treated threshings to livestock or allow to graze in treated fields. Do not use on Lima Beans that are to be removed from the field for processing. Do not feed to livestock.

CAULIFLOWER (14): Cabbage Looper, Cross-Striped Cabbageworm, Diamond Back Moth Larvae, Flea Beetle, Harlequin Bug, Imported Cabbageworm, Stink Bug - 4 ozs. per 500 sq. ft. Do not apply after edible parts begin to form. Close trim outer leaves prior to marketing.

CUCUMBERS (0), SQUASH (SUMMER AND WINTER): Aphids, Cucumber Beetles, Squash Vine Borer, Striped Cabbage Flea Beetle - 5 ozs. per 500 sq. ft. For Squash Vine Borer apply weekly to flower buds, stems and vines beginning when moths first appear.

EGGPLANTS (1): Blister Beetle, Colorado Potato Beetle, Green Peach Aphid, Green Stink Bug, Whitefly - 2 1/2 ozs. per 500 sq. ft.

FILBERTS (0): Filbert Aphid, Filbert Leaf Roller - 7 ozs. per 500 sq. ft. Do not graze livestock in treated groves.

GRAPES (7): Grape Leafhopper - 6 ozs. per 500 sq. ft. Do not apply to Concord variety as severe injury is likely to occur.

IRIS: Iris Borer - Dust soil surface of beds thoroughly in spring when growth starts. Three applications are needed at 2-week intervals.

LETTUCE HEAD (14): Cabbage Looper, Diamond Back Moth Larvae, Green Peach Aphid, Imported Cabbageworm - 5 ozs. per 500 sq. ft. Do not make more than 3 applications after thinning. Remove wrapper leaves at harvest. Do not feed crop refuse to livestock.

MELONS (0): Aphids, Cucumber Beetles, Striped Cabbage Flea Beetle - 4 ozs. per 500 sq. ft. Cabbage Looper, Omnivorous Leaf Roller - 6 ozs. per 500 sq. ft. Ringworm (on Watermelon only) - 4 ozs. per 500 sq. ft.

ORNAMENTALS (Greenhouse and Out-of-doors): Aphids, Whitefly - Dust lightly but evenly. On Chrysanthemums best results will be obtained if applied before plants flower. Do not use on Birch trees. In Greenhouse, wear mask or respirator approved by U. S. Bureau of Mines for pesticide protection.

PEAS (SEED PEAS ONLY): Pea Aphid, Pea Weevil - 5 ozs. per 500 sq. ft. Do not feed treated vines or threshings to livestock.

PEPPERS (1): Flea Beetle, Green Peach Aphid, Hornworms - 5 ozs. per 500 sq. ft. Pepper Maggot - 5 ozs. per 500 sq. ft.

POTATOES (0): Aphids, Armyworms, Colorado Potato Beetle, Green Stink Bug, Leaf-footed Bug, Plant Bugs, Potato Flea Beetle, Potato Leafhopper, Potato Tuberworm - 4 ozs. per 500 sq. ft. European Corn Borer, False Chinch Bug, Potato Psyllid - 4 ozs. per 500 sq. ft. Three-lined Potato Beetle - 2 1/2 ozs. per 500 sq. ft. Whitefly - 3 ozs. per 500 sq. ft. Do not plant root crops other than carrots, potatoes and sweet potatoes as follow-up crops.

PUMPKINS (0): Aphids, Cucumber Beetles, Squash Vine Borer - 4 ozs. per 500 sq. ft. For Vine Borer apply weekly to flower buds, stems and vines beginning when moths first appear.

SAFFLOWER: Green Peach Aphid - 5 ozs. per 500 sq. ft. Do not apply after flower heads open.

STRAWBERRY (4): Meadow Spittlebug, Strawberry Aphid - 4 ozs. per 500 sq. ft. Do not apply at intervals of less than 15 days or more than twice during any 35 day period when fruit is present.

SUGAR BEETS (30): Green Peach Aphid, Sugar Beet Webworm - 4 ozs. per 500 sq. ft. Meadow Spittlebug - 2 1/2 ozs. per 500 sq. ft. Do not feed sugar beet tops to livestock.

TOBACCO (5): Cabbage Looper, Green Peach Aphid, Tobacco Budworm, Tobacco Flea Beetle, Tobacco Hornworm - SEED BED 1 1/2 lbs. per 100 sq. yds. FIELD: 15 to 24 lbs. per acre. Cutworms, Pillbugs, Slugs (FLORIDA-GEORGIA SHADE GROWN TOBACCO AREA ONLY) - .375 to .56 lbs. per 100 sq. yds. plant bed. Direct dust at hiding places of these pests.

TOMATOES (1): Blister Beetles, Colorado Potato Beetle, Flea Beetle, Green Peach Aphid - 2 1/2 ozs. per 500 sq. ft. Cabbage Looper - 3 ozs. per 500 sq. ft. Stink Bugs - 4 ozs. per 500 sq. ft. Tomato Fruitworm, Yellow-striped Armyworm - 4 ozs. per 500 sq. ft. (For both field and greenhouse use.) Tomato Hornworm - 4 ozs. per 500 sq. ft. Tomato Russet Mite - 4 ozs. per 500 sq. ft. Whitefly - 2 1/2 ozs. per 500 sq. ft. (For both field and greenhouse use.)

NOTICE: Our recommendation for use of this product are based upon test believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice. The buyer must assume all responsibility, including injury or damage, resulting from its misuse as such or in combination with other materials.

CAUTION: This product is highly toxic to bees exposed to direct treatment or residue on crops. Protective information may be obtained from your County Agricultural Service.

CAUTION: KEEP OUT OF REACH OF CHILDREN

CAUTION: Harmful if swallowed, inhaled or absorbed through skin. Avoid contact with skin, eyes or clothing. Avoid breathing of dust. In case of contact with skin or eyes, flush with plenty of water, for eyes get medical attention. Wash with soap and water after handling and before eating or smoking, wear clean clothing. Avoid contamination of feed and foodstuffs. Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals. Workers entering treated areas within 24 hours of application should wear protective clothing.

CAUTION: This product is toxic to fish, shrimp, crabs, birds, and other wildlife. Bees and other wildlife in treated areas may be killed. Keep out of lakes, streams and ponds. Do not apply when runoff is likely to occur. Do not apply when weather conditions favor drift to areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

CONTAINER DISPOSAL

Destroy this container. Do not Re-Use. Destroy it by burning with water burning. If you do not burn it, crush it. Apply this product only as specified on this label.

EPA EST. NO. 3342-NC-1

ACCEPTED WITH COMMENTS