

DIRECTIONS

For use in Canteens, Restaurants, HOSPITALS, Schools, and Factories where POSITIVE DISINFECTANT and deodorizing is required on floors, walls, woodwork, desks and work tables. In food handling areas, remove food from area, disinfect and then rinse surface with potable water before reuse. Dilute this concentrate with water, using 4 oz. per gallon of water. Mop or wipe over areas to be disinfected and allow 10 minutes contact time to disinfect.

This disinfectant is non-selective and is effective against Salmonella Typhosa, Staphylococcus aureus, Salmonella choleraesuis, and Pseudomonas aeruginosa, organisms representative of gram positive and gram negative pathogens.

Where maximum deodorizing effect is to be obtained, leave solution for extended periods of contact time before mopping up.

EPA Reg. No. 330-33-100
EPA Reg. No. 330-33-100

LEMON DISINFECTANT No. 10

**FUNGICIDAL DISINFECTANT - DEODORIZER - CLEANER
FOR HOSPITALS AND INSTITUTIONS
PHENOL COEF. STAPH AUREUS - 10**

Active Ingredients: 1.7% p-AKVI-SO (C14-SO-3-12), 1.2% Benzyl Alcohol, 1.4% Essential Oils, 1.0% Ethanol, 1.0% Hydroxyethyl Cellulose, 1.0% Diamine Diacetate, 1.0% Diethylene Glycol, 1.0% Polyethylene Glycol, 1.0% Water, 1.0% Hydroxyethyl Cellulose, 1.0% Polyethylene Glycol, 1.0% Water, 1.0% Hydroxyethyl Cellulose, 1.0% Polyethylene Glycol, 1.0% Water.

WARNING - KEEP OUT OF REACH OF CHILDREN

Irritant if swallowed. Causes eye and skin irritation. Do not get into eyes. Avoid contact with skin. If contact occurs flush with plenty of water. If eye contact, immediately consult physician. Avoid contact with children. Rinse mouth with water after use. Wash hands and discard.

Manufactured by
REILY CHEMICAL CO.
New Orleans, La. 70111
FOR

ACCEPTED
MAR 10 1974
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REG. P.
NO. 22072 SUBJECT
TO ATTACHED COMMENTS.

**STATEMENT OF QUALITY
WARRANTY**

This product is guaranteed to meet AOAC (Association of Official Agricultural Chemists) Disinfection Confirmation Test vs. Salm. Choleraesuis, Staph. Aureus, Pseudomonas aeruginosa when tested in accordance with AOAC official methods in stated use dilution.

This product is effective against Trichophyton interdigitale in stated use dilution.

When tested in accordance with official AOAC methods for Phenol Coefficients, this product is equally non-selective and has shown the following phenol coefficients:

Staph. Aureus 10
Salm. Typhosa 7

CONTAINER DISPOSAL

Rinse empty container with water and discard it.

Net Contents 1 Gal.