

NON-SELECTIVE WEED CONTROL on NON-CROP LAND

Apply SIMAZINE 80W at rates recommended below. Use the higher rates on heavy clay and muck soils. Best results are obtained when SIMAZINE 80W is applied before weeds emerge. If weed growth has begun and removal is not practicable, include a contact or translocated herbicide in the spray.

Do not contaminate domestic or irrigation water supplies or lakes, streams or ponds.

SIMAZINE 80W can be used to provide long-term weed control on industrial sites, highway medians and shoulders, railroad rights-of-way, lumberyards, petroleum tank farms, and in non-crop areas on farms, such as around buildings, equipment and fuel stops, along fences, roadsides, and lanes. Aerial applications may be made where it is practicable.

Use sufficient water to assure thorough coverage. Use at least 1 gal. of water for each 1 lb. of SIMAZINE 80W; more if practical for both ground and aerial applications. Mechanical by-pass, or jet agitation is necessary to keep SIMAZINE in suspension during application.

To control annual broadleaf and grass weeds (such as barnyard and grass weeds, cheat, crabgrass, lambsquarters, foxtail, ragweed, puncturevine and turkey mullein): Broadcast 6-12½ lbs. per acre.

control most annual and many perennial broadleaf and grass weeds (including quackgrass, bluegrass, redtop, burdock, Canada thistle, orchardgrass, purple top, smooth brome, dog fennel, and plantain): Broadcast 12½ - 25 lbs. per acre.

To control hard-to-kill perennial weeds (such as bull thistle and sow thistle): Broadcast 25 - 50 lbs. per acre. For longer residual control in regions of high rainfall at a long growing season, broadcast 25 - 50 lbs. per acre.

For small areas, 4½ ozs. per 1,000 sq. ft. is equivalent to 12½ lbs. per acre.