

SANITIZING AGENT

Mito dilutions do not lose their effectiveness on standing. They are colorless, odorless, and non-corrosive to metals and plastics.

CAUTION: Mito should not be used with soap.

Mito is compatible with non-ionic detergents such as TRI-CLEAN, MISSION LIQUID DETERGENT, and GLASS-NU.

EPA REG. N. 3417-20-AA

Mito is recommended for use in bars, restaurants, hotels, motels, schools, for the sanitation of glassware, silverware, dishes, cooking utensils, etc.

DIRECTIONS

SANITIZATION OF GLASSWARE, SILVERWARE, DISHES, COOKING UTENSILS, ETC. — After washing with suitable detergent, such as Brito or Glass-Nu, rinse thoroughly, then immerse in a sanitizing rinse made with one ounce of Mito in four gallons of water for thirty seconds to two minutes in accordance with local health Department requirements. Then rinse with fresh potable water. Allow to air dry — do not use a towel.

MITO

Phenol Coefficients (A.O.A.C. Method 20 C.):

Salmonella typhosa — 50 Staphylococcus aureus — 75

ACTIVE INGREDIENTS:

ethyl alcohol 2.5%

n-Alkyl (50% C₁₄ 40% C₁₂ 10% C₁₆)

dimethyl benzyl ammonium chlorides10%

INERT INGREDIENT: Water87.5%

DANGER: KEEP OUT OF REACH OF CHILDREN.

See side panel for first aid and additional precautions.



Net Contents ONE GAL. (3.786 Liters)

MISSION CHEMICAL CO.
JANITOR SUPPLIES

4990 NAPLES ST.
SAN DIEGO, CALIF.
PHONE 276-5350

MAILING ADDRESS:
P.O. BOX 80608
SAN DIEGO, CA 92138

ADDITIONAL PRECAUTIONS: Avoid contact with eyes, eye damage and skin irritation. Do not get in eyes, on skin or on clothing. Protect eyes and skin when handling. Harmful if swallowed. Avoid contamination of food.

FIRST AID: In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Call a physician immediately.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock respiratory depression and convulsion may be needed.

This product fulfills the criteria of appendix F of the Grade A Pasteurized Milk Ordinance 1965 Recommendations of the U.S. Public Health Service in water up to 23 PPM of hardness calculated as CaCO when tested by the A.O.A.C. Germicidal and Detergent Sanitizers Official Method.

Rinse empty containers thoroughly with water and discard it.

