

## MAXI-CLOR® Chlorinating Tablets Concentrate State SLOW DISSO

COMPLETELY SOLUBLE NO MINERAL SCALE BUILD-UP SUPER CHLORINATING TABLETS FOR WATER DISINFECTION

24-HR. CHLORINE RESIDUAL FOR ROUND THE CLOCK ACTIVITY

KEEP OUT OF REACH OF CHILDREN DANGER See side panel for precautions in handling.

NET WT. 5 LBS.

Mfg by: N. Jonas & Co., Inc. • Bensalem, PA 19020

ACCEPTED JAN 10 2002

and Act, as exerciced, for the three under a g

## PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER, CORROSIVE: Causes irreversible eye damage and skin burns. May be fatal if absorbed through skin. May be fatal if inhaled. Do not breath dust or spray mist irritating to nose and throat. Harmful if swallowed. Do not get in eyes, on skin, or on ciciting. Wear goggles or face shield, protective clothing and rubber gloves when handling this product. Wash thoroughly with scap and water after handling and before eating, drinking, or using robacco. Remove contaminated cluthing and wash before reuse.

	FIRST AID
IF SWALLOWED	<ul> <li>Cell a poison control center or coctor immediately for treament advice.</li> <li>Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>Do not joint anything by month to an unconscious person</li> </ul>
IF IN EYES	<ul> <li>Hold eye open and rinse slowly and gently with water tor 15-20 minutes.</li> <li>Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>Call a poison control center or doctor center for treatment auticity.</li> </ul>
IF ON SKIN OR CLOTHING	Take off contaminated clothing.     Rinse skin immediately with pienty of water for 15-20 minutes.     Call a noison conting center or doctor for treatment advice
IF INHALED	<ul> <li>Move person to fresh air.</li> <li>If person is not breathing, cali 911 or an ambulance, then give artificial respiration, preterably mouth-to-mouth, if possible.</li> <li>Cali a polson control center or doctor for further treatment_advice.</li> </ul>
	Have the product container or taber with you when calling a polson control center or doctor, or going for treatment.
	NOTE TO PHYSICIAN
P	robable mucosal damage may contraindicate the use of gastric lavage.

PHYSICAL AND CHEMICAL HAZARDS: Strong Oxidizing Agent. Mix only with water, Use clean dry utensits. Do not add this product to any dispensing device containing remnants of any other product. Such use may cause a violent reaction leading to fire or explosion. Contamination with moisture, organic matter, or other chemicals may start a chemical reaction, with generation of heat, liberation of hazardous gases, and possible generation of fire and explosion. In case of contamination or decomposition, do not reseal container. If possible isolate container in open air or well ventilated area. Flood with large volumes of water, if necessary.

This product is meant to help keep bacteria and algae under control in swimming pools.

DIRECTIONS FOR USE:

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- Before using Chlorinating Tablets make sure that there is a sufficient chlorine residual (3.5-5.0 ppm) in the swimming pool by superchlorination. Next day add 1 lb. Chlorine Stabilizer (Booster) for each 4,000 gallons of water.
- Place approximately 7 tablets for 2500 gals. of pool water either into a tablet dispenser and float it in the pool or directly into skimmer basket if one is available. In no case should tablets be thrown directly into or on a vinyl liner as bleaching will result.
- 3. A good pool water test kit should be used at least once a day to check both pH and chlorine levels. pH should be between 7.2-7.6 and adjusted as necessary. Chlorine should be between 1-2 ppm. If chlorine level is too low, add more tablets until there is enough to maintain the 1-2 ppm.

This product can be used in any dispenser that uses Chlorine Concentrate tablets or sticks. Should not be mixed with any other kind of chlorine as an explosion could result. NOTE: During heavy bathing loads, very hot weather, rain, etc. and at least once a week, chlorine residual should be raised to 3.5-5.0 ppm with granular chlorine (superchlorinated). It is preterable to do this after swimming day so that pool will have a chince to come back to the 1-2 ppm before swimmers go back in next day Reentry links treated swimming pools is prohibited above levels of 3 ppm of chlorine.

Tablets will last approximately 5-7 days depending upon temperature and amount of water flow. Since there are approximately 48 tablets to the pound, a 2 fb. Jar will last a 2,500 gal. pool 7 weeks; a 5,000 gal. pool approximately 3 1/2 weeks; and a 10,000 gal. pool a little less than 2 weeks. A 4 fb. jar will last a 2,500 gal. pool 14 weeks; a 5,000 gal. pool 7 weeks; and a 10,000 gal, pool approximately 4 weeks. etc.

STORAGE AND DISPOSAL: Keep this product dry in a tightly closed container, when not in use. Store in a cool, dry, well ventilated area away from heat or open flame, in case of decomposition, isolate container (if possible) and flood area with large amounts of water to dissolve all material before discarding this container. Do not reuse empty container but place in trash collection. Do not contaminate lood or feed by storage, disposal, or cleaning of equipment.

0 ると

432