

3432-61

1-15-2002

1/2

J-111199-OnGuard Super 3" 12/13/01 8:40 AM Page 1

# OnGuard™

MAXI-CLOR®

## Super 3" Chlorinating Tablets

CONCENTRATE  
STABILIZED  
SLOW DISSOLVE

COMPLETELY SOLUBLE, NO MINERAL SCALE BUILDUP  
SUPER CHLORINATING TABLETS FOR WATER DISINFECTION

24-HR. CHLORINE RESIDUAL FOR  
ROUND THE CLOCK ACTIVITY

**ACTIVE INGREDIENTS:**

Trichloro-s-Triazinetrione ..... 99%

OTHER INGREDIENT ..... 1%

TOTAL ..... 100%

Approximately 90% available chlorine

EPA Reg. No. 324-324  
EPA Est. No. 324-324

KEEP OUT OF REACH OF CHILDREN

**DANGER**

See side panel for precautions in handling.

**NET WT. 5 LBS.**

Mfg by: N. Jonas & Co., Inc. • Bensalem, PA 19020

ACCEPTED

JAN 15 2002

For The Federal Insecticide, Fungicide & Rodenticide Act, 1902, as amended, for the

3432-61

**PRECAUTIONARY STATEMENTS**

**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**DANGER. CORROSIVE.** Causes irreversible eye damage and skin burns. May be fatal if absorbed through skin. May be fatal if inhaled. Do not breathe dust or spray mists. Irritating to nose and throat. Harmful if swallowed. Do not get in eyes, on skin, or on clothing. Wear goggles or face shield, protective clothing and rubber gloves when handling this product. Wash thoroughly with soap and water after handling and before eating, drinking or using tobacco. Remove contaminated clothing and wash before reuse.

FIRST AID	
IF SWALLOWED	<ul style="list-style-type: none"><li>• Call a poison control center or doctor immediately for treatment advice.</li><li>• Have a person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor.</li><li>• Do not give anything by mouth to an unconscious person.</li></ul>
IF IN EYES	<ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li><li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li><li>• Call a poison control center or doctor center for treatment advice.</li></ul>
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
IF INHALED	<ul style="list-style-type: none"><li>• Move person to fresh air.</li><li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.</li><li>• Call a poison control center or doctor for further treatment advice.</li></ul>
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
NOTE TO PHYSICIAN	
Probable mucosal damage may contraindicate the use of gastric lavage.	

**PHYSICAL AND CHEMICAL HAZARDS:** STRONG OXIDIZING AGENT. Mix only with water. Use clean dry utensils. Do not add this product to any dispensing device containing remnants of any other product. Such use may cause a violent reaction leading to fire or explosion. Contamination with moisture, organic matter, or other chemicals may start a chemical reaction, with generation of heat, liberation of hazardous gases, and possible generation of fire and explosion. In case of contamination or decomposition, do not reseal container. If possible isolate container in open air or well ventilated area. Flood with large volumes of water, if necessary.

**This product is meant to help keep bacteria and algae under control in swimming pools.**

**DIRECTIONS FOR USE**

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

1. Before using Chlorinating Tablets make sure there is sufficient chlorine residual (3.5-5.0 ppm) in the swimming pool by superchlorination. Next day add 1 lb. Chlorine Stabilizer (Booster) for each 4,000 gallons of water.
2. Place approximately 1 super tablet for 6,800 gallons of water per week either into a tablet dispenser and float it in the pool or directly into a skimmer basket if one is available. In no case should tablets be thrown directly into or on a vinyl liner as bleaching will result.
3. A good pool water test kit should be used at least once a day to check both pH and chlorine levels. pH should be between 7.2-7.6 and adjusted as necessary. Chlorine should be between 1-2 ppm. If chlorine level is too low, add more tablets until there is enough to maintain the 1-2 ppm. This product can be used in any dispenser that uses Chlorine Concentrate tablets or sticks. Should not be mixed with any other kind of chlorine as an explosion could result.

**NOTE:** During heavy bathing loads, very hot weather, rain, etc., and at least once a week, chlorine residual should be raised to 3.5-5.0 ppm with granular chlorine (superchlorinated). It is preferable to do this after swimming day so that pool will have a chance to come back to the 1-2 ppm before swimmers go back in next day.

Reentry into treated swimming pools is prohibited above levels of 3 ppm of chlorine. Super tablets will last approximately 5-7 days depending upon temperature and amount of water flow. A 20 lb. pail will last a 10,000 gal. pool approximately 30 weeks; 30,000 gal. pool approximately 10 weeks. Each tablet weighs approximately 7.3 oz.

**STORAGE AND DISPOSAL:** Keep this product dry in a tightly closed container, when not in use. Store in a cool, dry, well ventilated area away from heat or open flame. In case of decomposition, isolate container (if possible) and flood area with large amounts of water to dissolve all material before discarding this container. Do not reuse empty container but place in trash collection. Do not contaminate food or feed by storage, disposal, or cleaning of equipment.