# PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS CAUTION

Harmful if swallowed. Do not preathe dust. Do not get on skin. In case of contact, wash immediately with soap and water. Do not contaminate feed or foodstuffs. Keep out of reach of children."

in loading operations wear a mask or respirator jointly approved by the mining Enforcement and Safety Administration (formerly U.S. Bureau of Mines) and by the National Institutes for Occupational Safety and Health under the provisions of 30 CER Part 1."

### **ENVIRONMENTAL HAZARDS**

This product is toxic to fish and wildlife. Birds feeding on treated areas may be harmed. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drill from treated areas. Do not confarminate water by cleaning of equipment or disposal of wastes. Apply this productionly as specified on this label.

### PHYSICAL OR CHEMICAL HAZARDS

DO NOT JOE P. JR. SPILL OR STORE NEAR HEAT OR OPEN FLAME.

#### **RE-ENTRY STATEMENT**

Do not use pesticide in area of unprotected humans or domestic animals. Do not allow entry to area treated until the "hemical has completely dried. Keep children and pets off treated area until chemical has been washed into the soil by watering or rain and area is completely dry.

#### STORAGE AND DISPOSAL

- 1 PROHIBITIONS Do Not Contaminate Water, Food, or Feed by Storage or Disposal Open Dumping is Prohibited
- PESTICIDE DISPOSAL Pesticide, Spray Mixture, or Rin sate that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies.
- CONTAINER DISPOSAL Dispose of in an incinerator or landful approved for pesticide containers, or bury in a safe place
- 4. GENERAL Consult Federal State or Local disposal authorities for approved alternative procedures such as limited open burning.

## 10% Toxaphene Dust

ACTIVE INGREDIENTS APR 16 1980
Toxaphene INERT INGREDIENTS APR 16 1980
TECHNICAL CHLORINATED CAMPHENE: ACHLORINE CONTENTS 67% to 69%)
The first ingredient in the contents of the contents of

# KEEP OUT OF REACH OF CHILDREN DANGER POISON

### Statement For Practical Treatment

IF SWALLOWED Drink one or two glasses of water and induce vomiting by touching back of throat with finger. Do not induce vom finger give anything is mouth to arm is onscious person. Get medical attention immediately.

- F. INHALE () Remove victim to fresh air and apply respiration if indicated
- IF ON SKIN Wash promptly with soap and water. Rinse thoroughts.
  IN EXES Rinse exes for at least 15 minutes with water and call a physic ari in-
- TN EYES Ringe eyes for at least 15 minutes with water and call a physic ari inmediately.

### Manufactured By Cape Fear Chemicals Inc.

ELIZABETHTOWN NORTH CAROLINA NET CONTENTS — 25 LBS.

EPA EST NO. 3342-NC-L

EPA Reg. No. 3342 84

### DIRECTIONS FOR USE

"It is a violation of Federal law to use this product in a man ner inconsistent with its tabelling."

WHEAT, BARLEY, OATS AND RYE: To control Climbing and Surface-Feeding Cutworms, Army Worms, Grasshop pers and Chinch Bugs, apply at the rate of 20 lbs. per acre. So not apply more than once after heads have begun to form. Do not apply within one week of harvest of Wheat, Rye and Oats, within 2 weeks of harvest of Barley. Do not pasture dairy animals or meat animals in treated fields.

CABBAGE, BROCCOLI, LETTUCE AND BRUSSELS SPROUTS: To control Cabbage Loopers, apply at the rate of 20 lbs. per acre. For Cutworms (soil), apply 40 lbs per acre to soil surface For Cabbage and Head Lettuce. Do not apply after the Heads begin to form. For Leaf Lettuce, do not apply after seeding state. For Broccoil and Brussels Sprouts, do not apply after the edible parts begin to form. On all the above, do not apply this product if leaves, stems or heads are to be fed to dairy animals or meat animals. "Do not apply within 30 days of harvest."

COLLARDS AND KALE: To control Army Worm, Cabbage Worm, Cabbage Looper, Climbing and Surface Feeding Cut worms, apply at the rate of 20 to 40 lbs, per acre. Do not apply more than 4 times nor more than 1 times after thinning. Do not apply within 21 days of harvest when 40 lbs, per acre is us. ed. "Do not apply within 30 days of harvest."

\$PINACH: To control Army Worm, Cabbage Worm, Cabbage Looper, Climbing and Surface Feeding Cutworms, apply only one time at the rate of 20 lbs. per acre. Do not apply within 21 days of harvess.

BEANS: To control Curworms (soil) apply 20 lbs per acre on soil in late afternoon. Do not apply after pods begin to form. 7 day. Pre-Harvest Interval. "Do not feed treated vines to dairy animals or animals being finished for slaughter."

TOMATORS: To central Temato Hornworm, Tomato Fruit Werm, Serpentine Leef Miner, Fiea Beetles, Cutworms, Cabbase Leeper, Russet Miles and Colorado Potato Beetles, Army Worms and related insects, apply at the rate of 40 lbs. to the acres. Do not feed treated forage to dairy animals or animals being finished for slaughter. "Do not apply within 21 days of harvest."

PEANUTS: To control Fall Army Worm and Velvet Bean Caterpillar, apply at the rate of 15 to 20 lbs per acre. "Do not feed follage to diary animals or animals being finished for slaughter."

CORN: Te control Army Worm and Fall Army Worm, apply at the rate of 10 lbs. per acre. Do not feed treated plants to dairy or meat animals.

sORGHUM: To control Army worm and Fall Army Worm apply at the rate of 30 tos per acre. Do not feed treated plants of ensilage from treated plants to diarry or meat animals. Do not use on Sorghum to be sold commercially or to be shipped interstate. Do not apply within 40 days of harvest or apply after heads have begun to form.

### **WARRANTY STATEMENT**

In the removemental rich exists of this principlate missing partitions. It is the even to prove a principle the use of this grown is required to exempt the manufacturer of the provention of the manufacturer of the provention of the manufacturer of the effect of the motion of the manufacturer of the effect of the motion of