



PRICE
\$7.95
EXTRACTION GUARANTEED

Vertagreen

TRADEMARK

Triple Duty


Fertilizer, Insecticide,
Crabgrass Preventer
10-6-4

ACTIVE INGREDIENT	By Weight
Diethyl ester of Tetrahydro- pyrophosphate acid	2.88%
Tetrahydro Chloride	1.00%
Equivalent to carbaryl-A, 7-methyl hexahydrophthalide Related Compounds	0.72% 0.40%
INERT INGREDIENTS	95.00%
	Total 100.00%

CAUTION: KEEP OUT OF REACH OF CHILDREN
ADDITIONAL CAUTIONS: SEE REVERSE SIDE

Treats 2500 Sq. Ft. (50X50)
25 Lbs. Net Wt.

Triple Duty



25 Lbs. Net Wt.

Vertagreen Triple Duty Fertilizer, Insecticide, Crabgrass Preventer 10-6-4

WITH DIMETHYL ESTER OF TETRAHYDRO-PYROPHOSPHATE ACID, CARBARYL-A, AND CHLORIDE

MINIMUM GUARANTEED ANALYSIS

Diethyl ester of tetrahydro- pyrophosphate acid	2.88%
Tetrahydrochloride	1.00%
Equivalent to carbaryl-A, 7-methyl hexahydrophthalide Related compounds	0.72% 0.40%
INERT INGREDIENTS	95.00%
	Total 100.00%



READ FOLLOWING CAREFULLY BEFORE APPLICATION

USE THIS PRODUCT ONLY FOR RECOMMENDED PURPOSES AND AT RECOMMENDED DOSE RATE. Do not use for the control of certain soil insects on grass, such as grubs, wireworms, and cutworms, plus the prevention control of annual weeds, such as pigweed, and yellow foxtail. Do not use for crabgrass control on lawns, pastures, and other grassy areas. When used as recommended, it will not harm desirable plants or animals.

DIRECTIONS FOR APPLYING

WHEN TO APPLY: This product must be applied in early spring before above ground growth begins. At the cessation of growing season, these weeds may germinate any time there is a warm rainy spell with night temperatures above 50° F. for 3 to 5 days.

SEEDING: The ideal time to sow lawn seeds is at the fall or spring seedings. It is necessary to sow seeds sufficiently early in spring so as to get a good stand of grass before desiccating by applying the pre-emergent control. An application should be made at the time of the first cutting of the grass. Do not use this product on lawns that have been mowed for more than 10 days in the recommended season.

COMMON CHICKWEED: This weed thrives in cool weather. In some areas it is a problem in the fall and spring months. Depending on the weather it grows anytime. To control this weed it is suggested that you use this product to prevent new plants from overwintering for a season.

HOW TO APPLY: Use a fertilizer spreader. Following settings are approximate for applying contents of this package at suggested rate of 10 lbs. per 1000 sq. ft. Settings are based on an average walking speed. Reduce settings for faster speed, raise settings for higher speed.

SPREADER SETTINGS CHART

Model	Setting
Woods 1091, 1092, 1094	7 1/2
Woods 1250, 1252	6 1/2
Control Spreaders Model	7 1/2
Jackson & Sons	6 1/2
Leam Spray	8
Scott	7 1/2

CAUTION

Do not use internally. Avoid inhalation or contact with eyes. Do not inhale dust. Do not use on lawns that have been treated with other herbicides and insecticides. Do not use on lawns that have been treated with other herbicides and insecticides.


This product is toxic to fish and wildlife. Keep out of lakes, streams, and ponds. Birds feeding on treated areas may be killed.

NOTICE


Agri-Chemicals, Inc. can accept no responsibility for injury to persons or property, or damage to crops, lawns, or other property resulting from failure to follow directions for use or to observe cautions printed on this package.

Patented for use under U.S. Patent Number 2,922,824

USDA Reg. No. 2442-556



Triple Duty



UNDER THE FEDERAL INSECTICIDE
ACT AND THE FEDERAL FERTILIZER ACT
FOR THE UNITED STATES DEPARTMENT OF AGRICULTURE
ED. JENCKE 3449-536

Vertagreen

TRADEMARKS

Triple Duty

Fertilizer, Insecticide, Crabgrass Preventer

10-6-4

ACTIVE INGREDIENT

By Weight

Dimethyl ester of Terephthalic acid	2.50%
Technical Chloranil	1.50%
Equivalent to cyfluthrin-4,	
7-methoxy tetrahydroimidazole	0.75%
Related Compounds	0.40%

INERT INGREDIENTS

91.20%
Total 100.00%

CAUTION: KEEP OUT OF REACH OF CHILDREN
ADDITIONAL CAUTIONS: SEE REVERSE SIDE

Treats 2500 Sq. Ft. (50X50)

25 Lbs. Net Wt.

10-6-4

WITH DIMETHYL ESTER OF T

MINIMUM GUARANTEED ANALYSIS
Total Nitrogen (N)
Available Phosphoric Acid (P₂O₅)
Soluble Potash (K₂O)
Where required by state law
Potential Acidity 500 lbs Calcium Carbonate Equivalent per ton



READ FOLLOWING

USE THIS PRODUCT ONLY FOR RE
Use this product as follows for
cutworms, grubs, mole crickets and
such as annual and perennial crab
grass, pursley, lambsquarters, purslane and
not harm established grass and will
as seeds germinate

DIREC
When to apply: This
germinate. With the
is damp rainy weather with night

SEEDING: The ideal time to
sow seed sufficiently early so that
applying the pre-emergence control
is out of the ground and delay se
period, the more grass seed must
days is not recommended

COMMON CHICKWEED: T
lem in the fall and spring, in othe
this weed, it is suggested that you
a season.

HOW TO APPLY: Use a ferti
contents of this package at sugges
an average walking speed. Reduce

SPEE
Wards 1001, 10
Ward, 1000, 10
Central (Success
Jackson & Beas
Lawn Beauty
Scott

Do not take internally. Avoid inhaling
stuff. Wash after handling or using.
are well watered and completely d

ACTION GUAR

tagreen

le Duty

Insecticide, s Preventer

By Weight

2.825
1.995

Vertagreen Triple Duty Fertilizer, Insecticide Crabgrass Preventer 10-6-4

WITH DIMETHYL ESTER OF TETRACHLOROTEREPHTHALIC ACID & CHLORDANE

MINIMUM GUARANTEED ANALYSIS
Total Nitrogen (N) 10.00%
Available Phosphoric Acid P_2O_5 6.00%
Soluble Potash K_2O 4.00%
Where required by state law
Potential Acidity 500 lbs Calcium Carbonate Equivalent per ton

ACTIVE INGREDIENTS	By Weight
Dimethyl ester of tetrachloro-terephthalic acid	2.825
Technical Chlordane Equivalent to octachloro-4,7-methanotetrahydroindane Related Compounds	1.995
INERT INGREDIENTS	9.180
Total	10.000

(This product contains 2.52 lbs. Dimethyl ester of tetrachloro-terephthalic acid and 1.20 lbs. Chlordane per 100 lbs.)

Distributed by
USS Agri-Chemicals, Inc.
a Subsidiary of United States Steel Corporation
P. O. Box 1635 Atlanta, Georgia

READ FOLLOWING CAREFULLY BEFORE APPLICATION

USE THIS PRODUCT ONLY FOR RECOMMENDED PURPOSES AND AT RECOMMENDED RATES.
Use this product as follows for the control of certain soil insects on lawns, such as cutworms, grubs, mole crickets and wireworms, plus the pre-emergence control of annual weeds such as smooth and hairy crab grass, green and yellow foxtail, common chickweed, pursley, lambsquarters, purslane and witch grass in lawns. When used as recommended, it will not harm established grass and will control above weeds for an entire season by killing their seeds as they germinate.

DIRECTIONS FOR APPLYING

WEED PREVENTION: This product must be applied in early spring before above ground weeds begin to germinate. With the exception of smooth and hairy crab grass, these weeds may germinate any time of the year. For best results, apply this product during a period of damp, rainy weather with night temperatures above 50° F. for 3 to 5 days.

SEEDING: The ideal time to sow lawn seed is in the fall. If spring seeding is necessary, sow seed sufficiently early so there is a noticeable green stand of grass before applying the pre-emergence control. As an alternative, apply this product as soon as the grass is out of the ground and delay seeding for approximately 60 days. The shorter the period, the more grass seed must be sown to insure a desirable stand. Delay of more than 60 days is not recommended.

COMMON CHICKWEED: This weed thrives in cool weather. In some areas it is common in the fall and spring, in others, depending on the weather, it grows anytime of the year. To prevent this weed, it is suggested that you use this product to prevent new plants from developing.

ity

Total 100.00%

(This product contains 2.52 lbs. Dimethyl ester of tetrachloro terephthalic acid and 1.20 lbs. Chlordane per 100 lbs.)



Distributed by Agri-Chemicals, Inc. Subsidiary of United States Steel Corporation P. O. Box 1225 Atlanta, Georgia

READ FOLLOWING CAREFULLY BEFORE APPLICATION

USE THIS PRODUCT ONLY FOR RECOMMENDED PURPOSES AND AT RECOMMENDED DOSAGE.

Use this product as follows for the control of certain soil insects on lawns, such as ants, cutworms, grubs, mole crickets and wireworms, plus the pre-emergence control of annual weeds, such as smooth and hairy crab grass, green and yellow foxtail, common chickweed, Florida pusley, lambsquarters, purslane and witch grass in lawns. When used as recommended, it will not harm established grass and will control above weeds for an entire season by killing plants as seeds germinate.

DIRECTIONS FOR APPLYING

When used as a pre-emergence control, this product must be applied in early spring before above weeds germinate. With the exception of common chickweed, these weeds may germinate any time there is damp rainy weather with night temperatures above 50° F. for 3 to 6 days.

SEEDING: The ideal time to sow lawn seed is in the fall. If spring seeding is necessary, sow seed sufficiently early so there is a noticeable green stand of grass before deadline for applying the pre-emergence control. As an alternative, apply this product as soon as the frost is out of the ground and delay seeding for approximately 60 days. The shorter the waiting period, the more grass seed must be sown to insure a desirable stand. Delay of less than 30 days is not recommended.

COMMON CHICKWEED: This weed thrives in cool weather. In some areas it is a problem in the fall and spring, in others, depending on the weather, it grows anytime. To control this weed, it is suggested that you use this product to prevent new plants from developing for a season.

HOW TO APPLY: Use a fertilizer spreader. Following settings are approximate for applying contents of this package at suggested rate of 10 lbs. per 1000 sq. ft. Settings are based on an average walking speed. Reduce setting for slower speed; raise setting for higher speed.

SPREADER SETTING GUIDE

Model	Setting
Wards 1091, 1092, 1095	7 1/2
Wards 1090, 1091	6 1/2
Central (Recessed Wheel)	7 1/2
Jackson & Sears	6 1/2
Lawn Beauty	6
Scott	7 1/2

CAUTION

Do not take internally. Avoid inhalation or contact with eyes. Do not contaminate feed or food stuffs. Wash after handling or using. Keep children and pets off treated areas until treated areas are well watered and completely dried.


This product is toxic to fish and wildlife. Keep out of lakes, streams, and ponds. Birds feeding on treated areas may be killed.

NOTICE

USS Agri-Chemicals, Inc. can accept no responsibility for injury to persons or for loss or damage to lawns, soil, or other property resulting from failure to follow directions for use or to observe cautions printed on this package.

Licensed for use under U. S. Patent Number 2,923,634

USDA Reg. No. 3442-556



ACC 11-14

UNDER THE FEDERAL FUNGICIDE AND RO FOR LHO...

Triple Duty