

ACCEPTED

3442-623

OCT 7, 1969

# 3442-623



UNDER THE FEDERAL FUNGICIDE AND FERTILIZER ACT FOR ECONOMIC UNDER NO.

INSECTICIDE FERTILIZER ACT REGISTERED

READ FOLLOWING CAREFULLY BEFORE APPLICATION

USE THIS PRODUCT ONLY FOR RECOMMENDED PURPOSES AND AT RECOMMENDED DOSAGE

# Thiram 75 WP

## Turf Fungicide & Animal Repellent

### For Golf Course Use

(Contains 75 lbs. Thiram per 100 lbs.)

ACTIVE INGREDIENT	By Weight
Thiram* . . . . .	75%
INERT INGREDIENTS . . . . .	25%
Total . . . . .	100%

\*Tetramethylthiuram disulfide


USDA Reg. No. 3442-623

## CAUTION

### Keep Out of Reach of Children

See other cautions on back panel

Net Wt. 3 Lbs.

 **Manufactured for**  
**Agri-Chemicals**  
 Division of United States Steel Corporation  
 P. O. Box 1685 Atlanta, Georgia

### DIRECTIONS FOR USE

Use this product for the control of brown patch, dollar spot and snow mold in lawns and as an animal repellent. Thiram is a gas repellent that is effective only on animals attempting to feed on the treated foliage. Use in sufficient water for good coverage.

**ANIMAL REPELLENT:** To protect trees, shrubs, nursery stock and ornamentals for 6 months or longer (depending on weather and other conditions) from rabbits, meadow mice and deer, use a mixture of 1 lb. of this product, 3 qts. of water and 2 lbs. of a weather resistant sticker. Do not mix more repellent than can be used in a short time. Apply when plants are dormant or growing but before expected time of animal damage. Use only when surfaces of plants to be treated are dry and under favorable drying conditions.

**SPRAY APPLICATION:** Apply mixture as a full-coverage spray with a hand or mechanical sprayer. For tree seedlings, do not treat more than 3,000 seedlings per gallon of finished spray.

**DIP APPLICATION:** Apply mixture to bundles of tree planting stock by dipping to thoroughly wet foliage and other parts to be protected. Permit excess repellent to run off treated plants without coming in contact with roots.

**BUNDLES OF NURSERY PLANTING STOCK SHOULD BE PERMITTED TO DRY THOROUGHLY BEFORE PLANTING.**

**BRUSH APPLICATION:** Apply on lower trunk of fruit trees or ornamental plantings from ground level to a height of 2 1/2 ft. to protect plants from trunk injury.

**DO NOT USE ON PLANT PARTS THAT ARE TO BE USED FOR FOOD OR FEED.**

**BROWN PATCH AND DOLLAR SPOT:** Use 6 oz. per 1,000 sq. ft. Apply early before diseases appear. During summer when periods of high temperature and humidity are favorable for disease development, apply at 10 day intervals.

**SNOW MOLD:** Use 8 oz. per 1,000 sq. ft. in late fall before snow falls and 4 oz. per 1,000 sq. ft. in the spring just after snow has melted. If repeated rains or periods of thawing occur, treatment at lower rate during open period in mid-winter will provide added protection.

NOTE: 1 oz. equals 6 level tablespoonfuls.

### CAUTION

- May cause skin irritation.
- Do not inhale dust.
- Do not get in eyes or on skin.
- Wash thoroughly after handling.
- Do not use grass clippings for food or feed purposes.

Avoid contaminating ponds or streams as fish may be harmed.

### NOTICE

USS Agri-Chemicals can accept no responsibility for injury to persons or animals or for loss or damage to turf, soil, or other property resulting from failure to follow directions for use or to observe cautions printed on this label.