

READ FOLLOWING CAREFULLY BEFORE APPLICATION

USE THIS PRODUCT ONLY FOR RECOMMENDED PURPOSES AND AT RECOMMENDED DOSAGE

DIRECTIONS FOR USE

For use on bluegrass, fescue, Bermudagrass and bentgrass in fairways. Do not use this product on golf greens or tees. Weeds affected are as follows:

| | |
|---|---|
| <p>SUSCEPTIBLE</p> <p>Dandelions (Taraxacum spp.) Clover (Trifolium spp.) Common Chickweed (Stellaria media) Mouse-ear Chickweed (Cerastium vulgatum) Plantain (Plantago spp.) Pigweed (Amaranthus spp.) Ragweed (Ambrosia spp.) Lambsquarters (Chenopodium album)</p> | <p>MODERATELY SUSCEPTIBLE</p> <p>Knotweed (Polygonum spp.) Black medic (Medicago lupulina) English daisy (Bellis perennis) Dock (Rumex spp.) Purslane (Portulaca oleracea) Wood sorrel (Oxalis spp.) Mallow (Malva spp.)</p> |
|---|---|

APPLICATION RATE: For established turf, use 1 1/2 fluid oz. in 1 to 2 gal of water per 1000 sq. ft. or 4 pints in 30 to 40 gals. per acre.

MIXING: Use clean spray equipment. Determine amount of water and MCP-2,4-D to be placed in tank. Partially fill tank with clean water, add herbicide, mix thoroughly, and add balance of water.

WHEN TO APPLY: Apply anytime weeds are young and actively growing. It is important that application be made when there is plenty of soil moisture, following a good rain or thorough watering. Do not water for at least 24 hours after application. If rain occurs within this period, effectiveness may be reduced, but wait at least a week to determine whether another application is necessary.

HOW TO APPLY: Use equipment which applies product uniformly. A boom type sprayer is recommended.

MOWING: Since action is through leaves, to provide adequate leaf area for best absorption of chemical, avoid cutting or mowing turf for 1 to 2 days before or after spraying.

RESPONSE: Full effects of application may not be visible for about three weeks.



ACCEPTED

JUN 2 0 1969

MCP-2,4-D

(Amine Salt)
Broadleaf Weed Killer

[Contains 2 lbs. 2-(2-Methyl-4-chlorophenoxy) propionic acid (MCP) and 1 lb. 2,4-Dichlorophenoxyacetic acid (2,4-D) per gal.]

UNDER THE FEDERAL INSECTICIDE, FUNGICIDE AND RODENTICIDE ACT FOR ECONOMIC POISON REGISTRATION

ACTIVE INGREDIENTS

| | |
|---|---------------------|
| Diethanolamine salt of 2-(2-Methyl-4-chlorophenoxy) propionic acid* | By Weight 31.00% |
| Diethanolamine salt of 2,4-Dichlorophenoxyacetic acid** | 15.35% |
| INERT INGREDIENTS | 53.65% |
| TOTAL | 100.00% |
| *2-(2-Methyl-4-chlorophenoxy) propionic acid (MCP) equivalent | 20.90% |
| **2,4-Dichlorophenoxyacetic acid (2,4-D) equivalent | 10.45% |

A Stable, Non-flammable, Non-corrosive Aqueous Solution

NOT FOR USE ON HOME LAWNS

USDA Reg. No. 3442-624

CAUTION Keep Out of Reach of Children

See other cautions on side panel

Net ContentsGal.

USS Agri-Chemicals, Inc.
Manufactured for
a Subsidiary of United States Steel Corporation
P. O. Box 1685 Atlanta, Georgia

Harmful to animals
Avoid contact with
After treatment, do not
DO NOT apply to any
a plant
DO NOT use first mow
DO NOT use for flower
DO NOT use for discolor
90° F.
DO NOT concentrate
DO NOT use in a suite
STORE in a cool, dry place
fungicides
PROTECT from children
DO NOT use on
Discard

USS Agri-Chemicals failure label.



ACCEPTED

JUN 2 0 1969

M CPP - 2,4-D

(Amine Salt)
Broadleaf Weed Killer

[Contains 2 lbs. 2-(2-Methyl-4-chlorophenoxy) propionic acid (M CPP) and 1 lb. 2,4-Dichlorophenoxyacetic acid (2,4-D) per gal.]

UNDER THE FEDERAL INSECTICIDE, FUNGICIDE AND RODENTICIDE ACT FOR ECONOMIC POISON REGISTERED

ACTIVE INGREDIENTS

| | |
|---|----------------|
| Diethanolamine salt of 2-(2-Methyl-4-chlorophenoxy) propionic acid* | By Weight |
| Diethanolamine salt of 2,4-Dichlorophenoxyacetic acid** | 31.00% |
| INERT INGREDIENTS | 53.65% |
| TOTAL | 100.00% |

| | |
|--|--------|
| * 2-(2-Methyl-4-chlorophenoxy) propionic acid (M CPP) equivalent | 20.90% |
| ** 2,4-Dichlorophenoxyacetic acid (2,4-D) equivalent | 10.45% |

A Stable, Non-flammable, Non-corrosive Aqueous Solution

NOT FOR USE ON HOME LAWNS

USDA Reg. No. 3442-624

CAUTION Keep Out of Reach of Children

See other cautions on side panel

Net ContentsGal.



Manufactured for
Agri-Chemicals, Inc.
a Subsidiary of United States Steel Corporation
P. O. Box 1685 Atlanta, Georgia

CAUTION

Harmful if swallowed.

Avoid contact with skin, eyes or clothing.

After use, wash hands and other parts of body contaminated with concentrate, spray or washings.

DO NOT apply when wind is blowing and DO NOT allow spray to drift to any other plants such as flowers, shrubs or vegetables as M CPP-2,4-D is a plant killer. Coarse sprays are less likely to drift.

DO NOT use on new grass seedlings until they have developed to stage of first mowing.

DO NOT use grass clippings from first mowing after treatment as a mulch for flowers or shrubs.

DO NOT apply during period of drought without first watering thoroughly.

DO NOT apply when temperatures are extremely high. Grass may become discolored or injured if this product is applied in very hot weather (over 90° F.).

DO NOT contaminate irrigation ditches, lakes, ponds or streams with concentrated solution, spray or spray washings.

DO NOT use same spray equipment for other purposes unless cleaned with a suitable chemical cleaner.

STORE in tightly closed containers, away from fertilizers, insecticides, fungicides, seeds, plants and water supplies.

PROTECT from freezing.

DO NOT reuse empty container. Wash thoroughly with water and detergent. Discard in a safe place.

NOTICE

USS Agri-Chemicals, Inc. can accept no responsibility for injury to persons or animals or for loss or damage to turf, soil or other property resulting from failure to follow directions for use or to observe cautions printed on this label.

READ FOLLOWING CAREFULLY BEFORE APPLICATION

USE THIS PRODUCT ONLY FOR RECOMMENDED PURPOSES AND AT RECOMMENDED DOSAGE

DIRECTIONS FOR USE

For use on bluegrass, fescue, Bermudagrass and bentgrass in fairways. Do not use this product on golf greens or tees. Weeds affected are as follows:

| SUSCEPTIBLE | MODERATELY SUSCEPTIBLE |
|--|---------------------------------|
| Dandelion (Taraxacum spp.) | Knotweed (Polygonum spp.) |
| Clover (Trifolium spp.) | Black medic (Medicago lupulina) |
| Common Chickweed (Stellaria media) | English daisy (Bellis perennis) |
| Moose ear Chickweed (Cirsium vulgatum) | Dock (Rumex spp.) |
| Plantain (Plantago spp.) | Purslane (Portulaca oleracea) |
| Pigweed (Amaranthus spp.) | Wood sorrel (Oxalis spp.) |
| Ragweed (Ambrosia spp.) | Melaleuca (Melaleuca spp.) |
| Lambquarters (Chenopodium album) | |

APPLICATION RATE: For established turf, use 1 1/2 fluid oz. in 1 to 2 gals. of water per 1000 sq. ft. or 4 pints in 30 to 40 gals. per acre.

MIXING: Use clean spray equipment. Determine amount of water and MCPP-2,4-D to be placed in tank. Partially fill tank with clean water, add herbicide, mix thoroughly, and add balance of water.

WHEN TO APPLY: Apply anytime weeds are young and actively growing. It is important that application be made when there is plenty of soil moisture, following a good rain or thorough watering. Do not water for at least 24 hours after application. If rain occurs within this period, effectiveness may be reduced, but wait at least a week to determine whether another application is necessary.

HOW TO APPLY: Use equipment which applies product uniformly. A boom type sprayer is recommended.

MOWING: Since action is through leaves, to provide adequate leaf area for best absorption of chemical, avoid cutting or mowing turf for 1 to 2 days before or after spraying.

RESPONSE: Full effects of application may not be visible for about three weeks.



ACCEPTED

JUN 2 0 1964

MCPP-2,4-D

(Amine Salt)
Broadleaf Weed Killer
 [Contains 2 lbs. 2-(2-Methyl-4-chlorophenoxy)propionic acid (MCPP) and 1 lb. 2,4-Dichlorophenoxyacetic acid (2,4-D) per gal.]

| ACTIVE INGREDIENTS | By Weight |
|--|-----------|
| Diethanolamine salt of 2 (2-Methyl-4-chlorophenoxy)propionic acid* | 31.00% |
| Diethanolamine salt of 2,4-Dichlorophenoxyacetic acid** | 15.35% |
| INERT INGREDIENTS | 53.65% |
| TOTAL | 100.00% |
| *2 (2-Methyl-4-chlorophenoxy)propionic acid (MCPP) equivalent | 20.90% |
| **2,4-Dichlorophenoxyacetic acid (2,4-D) equivalent | 10.45% |

A Stable, Non-flammable, Non-corrosive Aqueous Solution

NOT FOR USE ON HOME LAWNS

USDA Reg. No. 3442-624

CAUTION Keep Out of Reach of Children

See other cautions on side panel

Net Contents Gal.

Manufactured for
USS Agri-Chemicals, Inc.
 a Subsidiary of United States Steel Corporation
 P. O. Box 1685 Atlanta, Georgia

Harmful if swallowed

Avoid contact with skin

After use, wash hands

thoroughly, spray or wash

DO NOT apply when

in any other plants or

a plant killer. Coarse

DO NOT use on new

grass until after

first mowing.

DO NOT use grass clippings

for flowers or shrubs.

DO NOT apply during

heavy rain or

DO NOT apply when

temperature is

discoloring or injured

(90° F.).

DO NOT contaminate

water with

concentrated solution, spray

DO NOT use same sprayer

for a suitable chemical

DO NOT use in

STORE in tightly closed

containers, seeds, plants

PROTECT from freezing

DO NOT reuse empty

containers. Discard in a safe place

USS Agri-Chemicals, Inc.

Atlanta, Georgia

failure to follow directions on

label.



ACCEPTED

JUN 2 0 1969

MCPP - 2,4-D

(Amine Salt)
Broadleaf Weed Killer

[Contains 2 lbs. 2-(2-Methyl-4-chlorophenoxy)propionic acid (MCPP) and 1 lb. 2,4-Dichlorophenoxyacetic acid (2,4-D) per gal.]

| ACTIVE INGREDIENTS | By Weight |
|--|----------------|
| Diethanolamine salt of 2-(2-Methyl-4-chlorophenoxy)propionic acid* | 31.00% |
| Diethanolamine salt of 2,4-Dichlorophenoxyacetic acid** | 15.35% |
| INERT INGREDIENTS | 53.65% |
| TOTAL | 100.00% |

* 2-(2-Methyl-4-chlorophenoxy)propionic acid (MCPP) equivalent 20.50%

** 2,4-Dichlorophenoxyacetic acid (2,4-D) equivalent 10.45%

A Stable, Non-flammable, Non-corrosive Aqueous Solution

NOT FOR USE ON HOME LAWNS

USDA Reg. No. 3442-624

CAUTION Keep Out of Reach of Children

See other cautions on side panel

Net ContentsGal.

Manufactured for
USS Agri-Chemicals, Inc.
a Subsidiary of United States Steel Corporation
P. O. Box 1685 Atlanta, Georgia

CAUTION

Harmful if swallowed

Avoid contact with skin, eyes or clothing.

After use, wash hands and other parts of body contaminated with concentrate, spray or washings.

DO NOT apply when wind is blowing and DO NOT allow spray to drift to any other plants such as flowers, shrubs or vegetables as MCPP 2,4-D is a plant killer. Coarse sprays are less likely to drift.

DO NOT use on new grass seedlings until they have developed to stage of first mowing.

DO NOT use grass clippings from first mowing after treatment as a mulch for flowers or shrubs.

DO NOT apply during period of drought without first watering thoroughly.

DO NOT apply when temperatures are extremely high. Grass may become discolored or injured if this product is applied in very hot weather (over 90° F.).

DO NOT contaminate irrigation ditches, lakes, ponds or streams with concentrated solution, spray or spray washings.

DO NOT use same spray equipment for other purposes unless cleaned with a suitable chemical cleaner.

STORE in tightly closed containers, away from fertilizers, insecticides, fungicides, seeds, plants and water supplies.

PROTECT from freezing.

DO NOT reuse empty container. Wash thoroughly with water and detergent. Discard in a safe place.

NOTICE

USS Agri-Chemicals, Inc. can accept no responsibility for injury to persons or animals or for loss or damage to turf, soil or other property resulting from failure to follow directions for use or to observe cautions printed on this label.

