



VERTAGARD

CHLORDANE 8 EC

FOR PROFESSIONAL TURF USE

(Contains 8 lbs. Chlordane per gal.)


ACTIVE INGREDIENTS	By Weight
Chlordane, technical	72%
Petroleum Distillate	22%
INERT INGREDIENTS	6%
TOTAL	100%

13.2% octachloro-1,7-methanotetrahydroindane and 28.8% related compounds

USDA Reg. No. 3112-625

CAUTION
Keep Out of Reach of Children

See other cautions on side panel



Net Contents Gal.

ARMOUR AGRICULTURAL CHEMICAL CO.

P. O. BOX 1685 • ATLANTA, GA.

READ FOLLOWING CAREFULLY BEFORE APPLICATION

USE THIS PRODUCT ONLY FOR RECOMMENDED PURPOSES AND AT RECOMMENDED DOSAGE

DIRECTIONS FOR USE

This product is an emulsifiable concentrate and the amounts shown below should be mixed with sufficient water to provide adequate coverage.

SURFACE PESTS: For control of ants, armyworms, chiggers, crickets, earwigs, fleas, grasshoppers, lawn moths or sod webworms, mole crickets, spiders and ticks, use 1 tablespoonful (1/2 fluid oz.) of concentrate in sufficient water to spray 1000 sq. ft. area of turf or ornamental plants. (Approximately 1.1-3 lbs. Chlordane per acre). Repeat application when necessary.

SOIL INSECTS: For control of Asiatic garden beetle grubs, European chafers, grubs, green June beetle grubs, Japanese beetle grubs, Northern masked chafers, grubs, Oriental beetle grubs, white fringed beetle grubs, other white grubs and wireworms, use 4 tablespoonfuls (1/2 fluid oz.) of concentrate in sufficient water to spray 1000 sq. ft. of turf area. (Approximately 5 1/2 lbs. Chlordane per acre).

MOSQUITOES: Use 1 to 2 teaspoonfuls (1/6 to 1/3 fluid oz.) of concentrate in sufficient water to spray 1000 sq. ft. area of turf or shrubbery. (Approximately 1/2 to 1 lb. Chlordane per acre). Repeat application when necessary.

CRABGRASS: For the suppression or control of crabgrass and silver crabgrass (poison-grass), use 1 1/2 to 2 pints of concentrate in sufficient water to treat 1000 sq. ft. of turf area. (Approximately 15 to 25 lbs. Chlordane per acre). Make application in the spring 2 to 4 weeks before crabgrass seeds germinate. Do not apply within 30 days before or after seeding. This is an early season preventative treatment and is not effective as a post-emergence control of crabgrass plants. This application will also provide extended control of the above soil insects, plus temporary control of surface pests.

CAUTION

Harmful if swallowed.

Contact with skin can cause toxic symptoms.

Avoid breathing spray mist.

In case of contact with skin, wash with soap and water.

Avoid contamination of feed and foodstuffs.

Keep children and pets off treated turf areas until treated areas are well watered and completely dried.

Do not apply when weather conditions favor drift of spray from turf or shrubbery areas treated.

This product is toxic to fish and wildlife. Birds feeding on treated areas may be harmed. Keep out of lakes, ponds, and streams. Do not apply to any area not specified on the label.

NOTICE

Armour Agricultural Chemical Company can accept no responsibility for injury to persons or animals or for loss or damage to turf, soil, or other property resulting from failure to follow directions for use or to observe cautions printed on this package.