

Code: 13-F

5-6-69

USS Vertagreen Weed & Feed for Professional Turf

Copy for upper 1/3 of back:

USS

Vertagreen
Weed & Feed
for Professional Turf

12-4-8 plus herbicide

MINIMUM GUARANTEED ANALYSIS

Total Nitrogen (N)	12%
Available Phosphoric Acid (P ₂ O ₅)	4%
Soluble Potash (K ₂ O)	8%

ACTIVE INGREDIENT	by Weight
N-butyl-N-ethyl-2,2,4-trifluoro- 2,6-dinitro-p-toluidine	.55%
INERT INGREDIENTS	99.45%
TOTAL	100.00%

Contains .55 lb. benefin per 100 lbs.

USS Agri-Chemicals, Inc.
a Subsidiary of United States Steel Corporation
P. O. Box 1685 Atlanta, Georgia

Code: 13-F

5-6-69

USS Vertagreen Weed & Feed for Professional Turf

Copy for Face:

USS

Vertagreen
Trademarks

Weed & Feed
For Professional Turf

ACTIVE INGREDIENT

N-butyl-N-ethyl-2,2,4-trifluoro- 2,6-dinitro-p-toluidine	.55%
INERT INGREDIENTS	99.45%
TOTAL	100.00%


By Weight

CAUTION KEEP OUT OF REACH OF CHILDREN
ADDITIONAL CAUTIONS: SEE REVERSE SIDE

See reverse side for weeds controlled and tolerant grasses

50 Lbs. Net Wt.

Weed & Feed



Code: 18-7

5-6-59

USS Verticuter on Hand & Food for Professional Turf

Copy for lower 2/3 of 1 ch.

READ FOR INFORMATION ONLY. THIS IS A PRELIMINARY REPORT. THE DATA IS NOT FINAL. DO NOT USE THIS PRODUCT FOR ANY OTHER PURPOSES. THE DATA IS NOT FINAL. DO NOT USE THIS PRODUCT FOR ANY OTHER PURPOSES.

This product contains a combination of quality fertilizer and selective pre-emergence herbicide which controls annual bluegrass (*Poa annua*), smooth and hairy crabgrass, green and yellow foxtail, barnyardgrass, and goosegrass (silver crabgrass or crabfoot) by killing plants as seeds germinate. It does not control established weeds.

Use only on established lawns listed below which are tolerant to the herbicide in this product when applied as directed. See PRECAUTIONS.

Bluegrass (perennial)	Bermudagrass
Ryegrass (perennial)	Centipedegrass
Fescue (except fine-leaved)	St. Augustinegrass
Bahiagrass	Zoysiagrass

DIRECTIONS FOR APPLYING

HOW TO APPLY: Apply to established turf before the germination of annual weeds listed above. The general times of application are covered below. The precise time of application for a specific area will vary significantly because of climate and individual cultural practices. Consult State Agricultural Experiment Station Weed Specialists or Local Turf Authorities for any additional information regarding specific weed germination dates in your area.

SOUTHERN (warm season) GRASSES:

1. For pre-emergence control of annual bluegrass (*Poa annua*), apply in late summer to early fall, prior to annual bluegrass seed germination. Additional applications at the same rate can be made 4 to 6 months later in early spring, prior to soil warming and subsequent germination. In areas heavily infested with annual bluegrass, its elimination will temporarily result in thin turf areas. The fertilizer in this product plus proper irrigation will encourage the existing turfgrasses to fill in the thin spots previously occupied by the annual bluegrass.
2. For pre-emergence control of crabgrass, foxtail, barnyardgrass and goosegrass, apply in late winter or early spring prior to annual weed grass germination. A second application may be made 4 to 6 months after initial treatment for continued control of annual weeds which may germinate into the summer or early fall.

HOW TO APPLY: Use 12.5 lbs. per 1000 sq. ft. or 550 lbs. per acre (equivalent to 5 lb. of active ingredient per acre).

NORTHERN (cool season) GRASSES:

1. For pre-emergence control of annual bluegrass (*Poa annua*), apply in late summer or early fall (prior to soil warming), or in early spring prior to annual bluegrass seed germination.

For pre-emergence control of crabgrass, foxtail, barnyardgrass and goosegrass, apply in late winter or early spring prior to annual weed grass germination.

Use 12.5 lbs. per 1000 sq. ft. or 550 lbs. per acre (equivalent to 5 lb. of active ingredient per acre).

Code: 18-7

5-6-59

USS Verticuter on Hand & Food for Professional Turf

Copy for lower 2/3 of 1 ch.

HOW TO APPLY: Use a fertilizer applicator, avoiding types which would apply product in narrow concentrated bands. Apply evenly over the turf without skips or misses. A more uniform application can be made by spreading half the required amount of product over the area to be treated and then applying the remaining half at right angles or crosswise.

RESEEDING: Chemicals which effectively control annual weed seeds may also affect new desirable turfgrass seedlings. Reseeding should be delayed for at least 12 to 16 weeks after application. When reseeding, it is necessary that proper irrigation be maintained. To insure satisfactory reseeding results after using this product, the amount of seed used should be increased.

PRECAUTIONS: Do not apply in the spring to turfgrasses planted the previous fall. Do not apply to bentgrass putting greens or dichondra turf. Do not apply to newly sprigged areas of Bermudagrass, St. Augustinegrass, or Centipedegrass until these turfs are well established. If verticutting is done, do not apply for 2 to 3 weeks. When applied as recommended, this product will thin fine-leaved fescues.

CAUTION KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed or absorbed through the skin. Avoid contact with skin, eyes, or clothing. In case of contact, flush with water. Do not contaminate food or foodstuffs. Do not contaminate any body of water by direct application, cleaning of equipment, or disposal of wastes. Do not let livestock graze treated areas. Do not feed forage from treated areas to livestock. Destroy bags when empty. Do not reuse.

NOTICE

USS Agri-Chemicals, Inc. can accept no responsibility for injury to persons or animals or for loss or damage to turf, soil, or other property resulting from failure to follow directions for use or to observe cautions printed on this package.

USDA Reg. No. 5342-