

Way Green

MINIMUM GUARANTEED ANALYSIS

Total Nitrogen (N) 10%
Available Phosphoric Acid (P₂O₅) 6%
Soluble Potash (K₂O) 4%
Where required by State law:
Potential acidity 500 lbs. Calcium
Carbonate Equivalent per ton

ACTIVE INGREDIENTS

Technical Chlordane 1.2%
Equivalent to octachloro-4,
7-methanotetrahydroindane 0.72%
Related Compounds 0.48%
Siduron [1-(2-methylcyclohexyl)-3-phenylurea] 2.3%
INERT INGREDIENTS 96.5%
TOTAL 100.0%

This product contains 1.2 lbs. Chlordane and
2.3 lbs. Siduron per 100 lbs.



Distributed By
Agri-Chemicals
Division of United States Steel Corporation
P.O. Box 1685 Atlanta, Georgia

Handwritten notes:
05/17/72
2/1/72
44272

READ FOLLOWING CAREFULLY BEFORE APPLICATION

USE THIS PRODUCT ONLY FOR RECOMMENDED PURPOSES AND AT RECOMMENDED DOSAGE
Use this product on lawns to control soil insects and for the pre-emergence control of annual weeds as follows:


SOIL INSECTS: Ants, cutworms, grubs, mole crickets and wireworms.

ANNUAL WEEDS. Smooth and hairy crabgrass, foxtail, and barmyardgrass, in newly seeded or established plantings of the following grasses only: bluegrass, fescue, redtop, smooth brome, perennial ryegrass, orchardgrass and the following strains of bentgrass: Penncross, Seaside, Highland, Astoria, Nimisila, C-1, C-7 and C-19. Do not use on other bentgrass strains nor on Bermudagrass, as injury to turf may result.

Effect on golf course greens may vary with turf management practices and environmental conditions; there-



Agri-Chemicals
 Division of United States Steel Corporation
 P.O. Box 1685 Atlanta, Georgia



READ FOLLOWING CAREFULLY BEFORE APPLICATION

USE THIS PRODUCT ONLY FOR RECOMMENDED PURPOSES AND AT RECOMMENDED DOSAGE
 Use this product on lawns to control soil insects and for the pre-emergence control of annual weeds as follows:

SOIL INSECTS: Ants, cutworms, grubs, mole crickets and wireworms.

ANNUAL WEEDS. Smooth and hairy crabgrass, foxtail, and barnyardgrass, in newly seeded or established plantings of the following grasses only: bluegrass, fescue, reedtop, smooth brome, perennial ryegrass, orchardgrass and the following strains of bentgrass: Penncross, Seaside, Highland, Astoria, Nimisila, C-1, C-7 and C-19. Do not use on other bentgrass strains nor on Bermudagrass, as injury to turf may result.

Effect on golf course greens may vary with turf management practices and environmental conditions; therefore, its use on golf greens is not recommended, as injury to turf may result. Treatment will not control annual bluegrass (*Poa annua*), Clover, or most broadleaf weeds.

This product may be applied at any stage of turf development, including new seedings.

At least 1/2 inch of water must be provided by rainfall or watering within 3 days after treatment in order to carry the chemical into the root zone of germinating weeds.

DIRECTIONS FOR APPLYING

WHEN TO APPLY: Use in the spring before crabgrass germinates.

AMOUNT TO APPLY:

BAREGROUND SEEDINGS: In starting a new lawn or filling in open areas in established lawns in the spring, apply 5 lbs. per 1000 sq. ft. as the last operation over newly seeded areas. When over-seeding to thicken an existing lawn, follow directions for established lawns. Follow regular watering practices for good seed germination.

ESTABLISHED LAWNS: Apply 10 lbs. per 1000 sq. ft.

HOW TO APPLY: Use a fertilizer spreader. Following settings are approximate for applying contents of this package at rates indicated above. Settings are based on an average walking speed. Reduce for slower speed; raise setting for higher speed.

SPREADER SETTING GUIDE

<u>Make</u>	<u>New Seeded Area Setting</u>	<u>Established Lawn Setting</u>
Wards 1080, 1081	4 3/4	6 1/4
Central (Recessed Wheel)	6 3/4	7 1/2
Jackson & Sears	5	6 1/2
Lawn Beauty	6 3/4	8
Scott	5 1/2	7 1/8

CAUTION

Keep Out of Reach of Children

Do not take internally. May irritate eyes, nose, throat, and skin. Avoid inhalation or contact with skin, eyes, or clothing. Do not contaminate feed or foodstuffs. Wash after handling or using. Keep children and pets off treated areas until treated areas are well watered and completely dried.

This product is toxic to fish and wildlife. Keep out of lakes, streams and ponds. Birds feeding on treated areas may be killed.

NOTICE

USS Agri-Chemicals can accept no responsibility for injury to persons or animals or for loss or damage to lawns, soil or other property resulting from failure to follow directions for use or to observe cautions printed

...to use on grass is not recommended, as injury to lawn may result. Treatment will not control annual

bluegrass (Poa annual), Clover, or most broadleaf weeds.

This product may be applied at any stage of turf development, including new seedings.

At least 1/2 inch of water must be provided by rainfall or watering within 3 days after treatment in order to carry the chemical into the root zone of germinating weeds.

DIRECTIONS FOR APPLYING

WHEN TO APPLY: Use in the spring before crabgrass germinates.

AMOUNT TO APPLY:

BAREGROUND SEEDINGS: In starting a new lawn or filling in open areas in established lawns in the spring, apply 5 lbs. per 1000 sq. ft. as the last operation over newly seeded areas. When over-seeding to thicken an existing lawn, follow directions for established lawns. Follow regular watering practices for good seed germination.

ESTABLISHED LAWNS: Apply 10 lbs. per 1000 sq. ft.

HOW TO APPLY: Use a fertilizer spreader. Following settings are approximate for applying contents of this package at rates indicated above. Settings are based on an average walking speed. Reduce for slower speed; raise setting for higher speed.

SPREADER SETTING GUIDE

<u>Make</u>	<u>New Seeded Area Setting</u>	<u>Established Lawn Setting</u>
Wards 1080, 1081	4 3/4	6 1/4
Central (Recessed Wheel)	6 3/4	7 1/2
Jackson & Sears	5	6 1/2
Lawn Beauty	6 3/4	8
Scott	5 1/2	7 1/8

CAUTION

Keep Out of Reach of Children

Do not take internally. May irritate eyes, nose, throat, and skin. Avoid inhalation or contact with skin, eyes, or clothing. Do not contaminate feed or foodstuffs. Wash after handling or using. Keep children and pets off treated areas until treated areas are well watered and completely dried.

This product is toxic to fish and wildlife. Keep out of lakes, streams and ponds. Birds feeding on treated areas may be killed.

NOTICE

USS Agri-Chemicals can accept no responsibility for injury to persons or animals or for loss or damage to lawns, soil or other property resulting from failure to follow directions for use or to observe cautions printed on this package.

USDA Reg. No. 3442-672

