

**WEED AND FEED**

**14-4-4**

**KILLS WEEDS - FEEDS TURF**

**WEED AND FEED  
KILLS WEEDS - FEEDS TU**

# DACO

40 LBS. NET WT.

# WEED AND FEED

## 14-4-4

### MINIMUM GUARANTEED ANALYSIS

Total Nitrogen (N).....	14
Water Insoluble Nitrogen.....	4
Water Soluble Nitrogen (N).....	10
Water Soluble Phosphorus (P <sub>2</sub> O <sub>5</sub> ).....	4
Water Soluble Potash (K <sub>2</sub> O).....	4
2,4-Dichlorophenoxyacetic Acid (2,4-D).....	By Weight
.....	14
.....	512
.....	12.952
.....	100.000
.....	250
.....	100

PREPARED BY  
**GEORGE A. DAVIS, INC.**  
 5430 Northway Highway  
 Chicago, Illinois 60630

DACO Weed and Feed can be used two ways at one time. Kills weeds and feeds the turf in the same application. No more back-breaking digging of dandelion or plantain when DACO WEED and FEED is used. The high quality of turf food gives outstanding results in a thicker, greener turf that stays green too for weeks longer. DACO Weed and Feed can be applied any time the weeds are in a rapid state of growth. DO NOT APPLY WITH INTELLECTUALS ARE BELOW AGE except for the use of DACO WEED and FEED when MOISTURE IS LACKING in the soil.

### DIRECTIONS

Chicago, Illinois 60630

DACO WEED and FEED does two jobs at one time. Kills weeds and feeds the turf in the same application. No more back-breaking digging of dandelion or plantain when DACO WEED and FEED is used. The high quality of turf food gives outstanding results in a thicker, greener turf. Stays green too for weeks longer. DACO WEED and FEED can be applied any time the weeds are in a rapid state of growth. DO NOT APPLY WHEN TEMPERATURES ARE BELOW 60°F. (except for chickweed) OR WHEN MOISTURE IS LACKING IN THE SOIL.

## DIRECTIONS

**ESTABLISHED LAWNS** — It is most important to apply evenly. Contents will cover 6,000 sq. ft. For more effective results, lawn may be watered lightly before applying. Lawn seed may be sown 2 weeks after application of DACO WEED and FEED.

**MOWING** — Grass should not be mowed for several days before and after application.

The following weeds are controlled by DACO WEED and FEED. Chickweed (see below), white clover, henbit, thyme-leaved speedwell, yarrow, spotted spurge, dandelion, plantain, buckhorn, ragweed, burdock, lawn pennywort, mustard, nettles, pigweed, ground ivy, sow thistle and jimson weed.

**CHICKWEED** — Early spring control is preferred, however, late summer or early fall is also recommended to destroy weeds that may have developed during the summer or spring. Application can be made any time temperature is above freezing.

**IMPORTANT:** Use only on established lawns. Avoid drift of dust on vegetables, flowers, ornamental trees and shrubs, and other desirable plants. If some of the material comes in contact with desirable plants, it is recommended that plants be syringed thoroughly with water as soon as possible. Vapors from this product may injure susceptible plants in the immediate vicinity.

Do not store near other fertilizer, seed, insecticides, or fungicides. Do not use on carpetgrass, centipede grass, St. Augustine grass, or Dichondra nor on lawns or turf where desirable bent-grass or clovers are present.

**CAUTION:** Avoid prolonged or repeated inhalation of dust or contact with skin, eyes or clothing. Wash thoroughly after handling.

This product is toxic to fish; keep out of lakes, streams and ponds. Do not apply when weather conditions favor drift from areas treated as this product may injure susceptible plants in the immediate vicinity.

## SPREADER SETTING CHART

Cyclone .....	4
Milorganite .....	4
Jackson .....	6½
Modern .....	5
Davis-Daco .....	6
Scott .....	6

USDA REG. NO. 3442-674

**A QUALITY TURF PRODUCT**  
**For GOLF COURSES, PARKS, & HOME USE**

**WEED**

**14**

**KILLS WEED**

**WEED AND FEED  
KILLS WEEDS - FEEDS TURF**