



Trademark

ACCEPTED

APR 19 1973

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
ED UNDER NO. 3442-682

Toxaphene 6 EC

Delta

Contains 6 lbs. Toxaphene per gallon

ACTIVE INGREDIENTS	By Weight
Toxaphene*	58.5%
Xylene	37.5%
INERT INGREDIENTS	4.0%
TOTAL	100.0%

*Technical Chlorinated Camphene (Chlorine content 67% to 69%)

EPA Reg. No. 3442-682

WARNING

Keep Out of Reach of Children

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME

See other cautions on side panel

Net Contents

Gal.



Manufactured for

Agri-Chemicals

Division of United States Steel Corporation

P. O. Box 1685

Atlanta, Georgia

READ FOLLOWING CAREFULLY BEFORE APPLICATION

USE THIS PRODUCT ONLY FOR RECOMMENDED PURPOSES AND AT RECOMMENDED DOSAGE

DIRECTIONS FOR USE

SOYBEANS AND PEANUTS

Cutworms, Fall armyworms, Velvetbean caterpillars—Apply 2-2/3 pints per acre in sufficient water for good coverage. Apply as soon as small worms appear. Do not use on soybeans within 21 days of harvest.

PECANS

Pecan weevils—Use 2 gallons to 500 gallons of water (3.2 pts. per 100 gals.) Do not apply more than 400 gallons per acre. For number and frequency of applications, follow locally recommended spray schedule. Two applications are usually necessary, the first when the weevils begin to appear as determined by jarring and the second about 10 days to 2 weeks later. Do not apply after shucks begin to open. Do not graze livestock in treated groves.

COTTON

Thrips—For early season control, apply 1 to 2 pints per acre in sufficient water for good coverage, usually 2 to 3 gallons with 1 nozzle per row. Make the first application as soon as cotton is up to a stand followed by a second application 7 days later. For overwintered boll weevil and mid-season to late season control, use an alternate recommended insecticide.

BARLEY, OATS, RYE, WHEAT, CORN AND SORGHUM

Armyworms, Fall armyworms and Grasshoppers—Apply 2 to 2-2/3 pints per acre in sufficient water for good coverage. Do not apply to sorghum more than once after heads begin to form nor within 28 days of harvest. Do not ensile treated forage.

VEGETABLES

Do not apply to or allow drift to reach cucumbers, melons or squash as injury may result.

CUTWORMS on beans, broccoli, cabbage, cauliflower, celery, eggplant, kale, peppers and tomatoes—Apply 2-2/3 pints per acre in sufficient water for good coverage. Apply to soil surface before planting or immediately after setting plants. Do not apply when plants are past seedling stage.

COWPEAS: Cowpea curculio—Apply 1/2 gallon per acre in sufficient water for good coverage. Make first application at time of bloom or when first small peas appear. Make total of 2 to 4 applications at 4 to 5 days intervals as necessary. Do not apply within 10 days of

harvest. Apply only on peas which will be shelled. Do not apply to peas to be used as green snaps after pods begin to form. Do not feed treated units to dairy animals or animals being finished for slaughter.

EGGPLANTS, PEPPERS AND TOMATOES: Blister beetles, Hornworms—Apply 2-2/3 to 4 pints per acre on eggplants and 2-2/3 to 5-1/3 pints per acre on tomatoes in sufficient water for good coverage. Pepper weevil on peppers—Apply 4 pints per acre in sufficient water for good coverage. Do not apply on eggplants or peppers within 5 days of harvest nor on tomatoes within 7 days of harvest. Remove all residue at harvest by washing or other effective means.

LIVESTOCK

Do not use on dairy animals. Do not exceed recommended rate. Concentrations of twice the recommended rate may be fatal to young animals. Do not apply to animals within 28 days of slaughter. Do not use this product in dips. Do not use on calves under 3 months. Avoid treating animals in cold, stormy weather. Avoid treating overheated or sick animals.

BEEF CATTLE

Horn flies, Lice and Ticks—Use 5-1/3 pints per 100 gallons of water (1/2 cup to 5 gallons) and spray at the rate of 2 quarts per animal. For horn flies and ticks, apply at 3 weeks intervals or as needed.

WARNING

May be fatal if swallowed. Do not breath spray mist. Do not get in eyes, on skin or on clothing. Wash thoroughly after using. Do not store near feed or foodstuffs.

WILDLIFE: This product is poisonous to fish and wildlife. Keep out of lakes, ponds and streams. Do not apply in any manner not specified on the label. Birds feeding on treated areas may be killed.

BEES: This product is toxic to bees and should not be applied when bees are actively visiting the area.

Do not feed treated crop or crop refuse to poultry, dairy animals or animals being finished for slaughter. Do not allow grazing of treated crops by dairy animals or animals being finished for slaughter. Do not sell or ship treated forage.

DISPOSAL WARNING

Dispose of waste pesticide and spillage by burying in non crop area away from water supplies. Containers should be perforated or crushed and buried with waste.

NOTICE

USS Agri-Chemicals can accept no responsibility for injury to persons or animals or for loss or damage to crops, soil or other property resulting from failure to follow directions for use or to observe cautions printed on this label.