

Reg # 3442 - 688

PM - 19

194

4/22/71

**RESTRICTED USE PESTICIDE**

FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS OR  
PERSONS UNDER THEIR DIRECT SUPERVISION AND ONLY FOR THOSE  
USES COVERED BY THE CERTIFIED APPLICATOR'S CERTIFICATION

**+ FIRST /**

Call a doctor (physician)  
that the victim has been  
scribe his condition.

If breathing has stopped  
and maintain until doctor

If swallowed and victim  
quickly. Induce vomiting  
giving soapy or strong st  
clear. Never give anythi  
Have victim lie down an

In case of contact, imme-  
water for at least 15 mi  
ing and shoes. See doct

**NOT**

Antidote—administer at  
FOUR mg. intravenously  
is overcome. Repeat at  
atropinization appear. 2  
be administered in cor  
MORPHINE OR TRANQ  
cholinesterase inhibitor  
ous systems and produci  
first sign of pulmonary e  
mental oxygen and treat  
of the poison may occu  
after initial improve  
PATIENT IS INDICATED

For additional informati  
methyl parathion poiso  
Memoranda on Econom  
Department of Health,  
Service, Communicable  
booklet also available t  
Washington, D. C.

**PO:**

Consult your State Ag  
Station regarding postir

NOTICE: USS Agri-Chen  
jury to persons or anim  
or other property resul  
to observe warni

# Methyl Parathion 4EC

Contains 4 lbs. Methyl Parathion per gallon

**ACTIVE INGREDIENTS**

By Weight

O,O-dimethyl O-p-nitrophenyl thiophosphate\* 45.10%


**INERT INGREDIENTS**

54.90%

TOTAL 100.00%

\*Methyl Parathion

EPA Reg. No. 3442-688



EPA Est. 3442-MS-1

**STOP—READ LABEL BEFORE USING!**

## POISON

See Side Panels  
for Antidote &  
Precautions

Net Contents



## DANGER

Keep Out of  
Reach of  
Children.

Gal.

Not Reviewed. Registrant claims to be in accordance with  
Guidance for Phase 2  
Change in Status.

**BEST AVAILABLE COPY**