



CONTAINER

PRINT SPECIFICATIONS

USS Vertagreen Crabgrass Control	NUMBER: PAGE: 2 of 4 DATE: 5/7/84
----------------------------------	---

Copy for Back

The following copy must appear in the upper 1/3 of back of bag

USS	ACCEPTED 3442-712 5/7/84	10 lbs. net wt.
Vertagreen		
Crabgrass Control		
ACTIVE INGREDIENT		By Weight
N-Butyl-N-ethyl-a,a,a-trifluoro-2,6-dinitro-p-toluidine.....		2.41
INERT INGREDIENTS.....		97.59
TOTAL.....		100.00

This product contains 2.41 lbs. benetia per 100 lbs.

(Active Ingredient area should be printed on a yellow background.)

DIRECTIONS FOR USE

GENERAL CLASSIFICATION

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This product controls smooth and hairy crabgrass, green and yellow foxtail, barnyard grass and goosegrass (silver crabgrass or crowfoot) by killing plants as seeds germinate. It does not control established weeds.

Use only on established lawns listed below which are tolerant to the herbicide in this product when applied as directed. SEE PRECAUTIONS.

- | | |
|-----------------------------|--------------------|
| Bluestem (perennial) | Bermudagrass |
| Ryegrass (perennial) | Centipedegrass |
| Fescue (except fine-leaved) | St. Augustinegrass |
| Bartlett grass | Zoysiagrass |

WITH TO APPLY: Apply to established lawns in the spring before germination of annual weeds listed above.

AMOUNT TO APPLY: Use 1 lb. per acre equivalent to 2.0 lbs. of active ingredient per acre.



CONTAINER PRINT SPECIFICATIONS

USS Vertaqaen Crabgrass Control	NUMBER: 442 112 PAGE: 3 of 4 DATE: 5/7/84
---------------------------------	---

Copy for Lower 2/3 of Back (Continued)

HOW TO APPLY: Use a fertilizer spreader, avoiding types which would apply product in narrow concentrated bands. Apply evenly over the turf without skips or misses. A more uniform application can be made by spreading half the required amount of product over the area to be treated and then applying the remaining half at right angles or crosswise.

RESEEDING: Chemicals which effectively control annual weed seeds may also affect new desirable turfgrass seedlings. Reseeding should be delayed for at least 6 weeks after application. When reseeding, it is necessary that proper moisture be maintained until grass seedlings are established. To assure satisfactory results after using this product, the amount of seed should be increased.

PRECAUTIONS: Do not apply in the spring to turfgrass planted the previous fall. Do not apply to bentgrass putting greens or Dichondra turf. Do not apply to newly sprigged areas of Bermudagrass, St. Augustinegrass, or Centipedegrass until these turfs are well established. If verticutting is done, do not apply for 2 to 3 weeks.

CAUTION

KEEP OUT OF REACH OF CHILDREN.

Harmful if swallowed or absorbed through the skin. Avoid contact with skin, eyes or clothing. In case of contact, flush with water. Do not contaminate food or feedstuffs. Do not contaminate any body of water by direct application, cleaning of equipment or disposal of wastes. Do not let livestock graze treated areas. Do not feed forage from treated areas to livestock.

STORAGE AND DISPOSAL

Storage: Keep container closed and away from food, feedstuff, or domestic water supplies.

Disposal: Do not reuse empty container. Wash container and put in trash collection.

Agri-Chemicals
Division of United States Steel
P.O. Box 1000 - Atlanta, Georgia
© 1984 USS Agri-Chemicals, a registered trademark.



CONTAINER

PRINT SPECIFICATIONS

30110	USS Vertagreen Crabgrass Control	NUMBER: 3420 2
		PAGE: 4 of 4
		DATE: 5/7/84

Copy for Lower 2/3 of Back (Continued)

NOTICE:

USS Agri-Chemicals can accept no responsibility for injury to persons or animals or for loss or damage to turf or other property resulting from failure to follow directions for use or to observe cautions printed on this package.

EPA Reg. No.
EPA Est. No.

Vertagreen

Crabgrass Control

(Caution paragraph should be printed on a yellow background)