


Vertagreen Fertilizer for Professional Use 22-3-9

Copy for Face of Bag:

U.S.

Vertagreen

Fertilizer^R For Professional Use
with Balan^R



Active Ingredient:
N-butyl-N-ethyl-a,a,a-trifluoro-
2,6-dinitro-p-toluidine
Inert Ingredients
Total

By Weight

.857%
99.163%
100.000%

Balan^R - the trademark for benefin, Elanco Products Company,
a division of Eli Lilly and Company.

KEEP OUT OF REACH OF CHILDREN

CAUTION:

ADDITIONAL CAUTIONS: SEE REVERSE SIDE

See Reverse Side for Weeds Controlled and Tolerant Grasses

50 lbs Net Wt./22.68 Kg.

Fertilizer for Professional Use
with Balan

Copy for Gussets:

Vertagreen Fertilizer for Professional Use with Balan

Note: Regarding above copy, minimum type size is
18 pt. for CAUTION
12 pt. for KEEP OUT OF REACH OF CHILDREN

(2)

3442-113

Vertagreen Fertilizer for Professional Use 22-3-9

Copy for Lower 2/3 of Back continued

2. For pre-emergence control of crabgrass, foxtail, barnyard grass, and goose grass, apply in early spring prior to annual weed grass germination.

AMOUNT TO APPLY: Use 4.1 lbs. per 1000 sq.ft. or 1.75 lbs. per acre (equivalent to 1½ lbs. of active ingredient per acre).

HOW TO APPLY: Use a fertilizer spreader, avoiding types which would apply product in narrow concentrated bands. Apply evenly over the turf without skips or misses. A more uniform application can be made by spreading half of the required amount of product over the area to be treated and then applying the remaining half at right angles or crosswise.

RESEEDING: Chemicals which effectively control annual weed seeds may also affect new desirable turf grass seedlings. Reseeding should be delayed for at least 12 to 16 weeks after application. When reseeding, it is necessary that proper irrigation be maintained. To insure satisfactory reseeding results after using this product, the amount of seed used should be increased.

PRECAUTIONS: Do not apply in the spring to turf grass planted the previous fall. Do not apply to bentgrass putting greens or dichondra turf. Do not apply to newly sprigged areas of Bermuda grass, St. Augustine grass or Centipede grass until these turfs are well established. If verticulating is done, do not apply for 2 to 3 weeks. When applied as recommended, this product will thin fine-leaved fescues.