



ACCEPTED

MAY - 8 1984

United States Insecticide,
Fungicide and Rodenticide Act,
as amended for the pesticide
registered under
EPA Reg. No. 3442-716

Vertagreen Crabgrass Preventer Plus

Net Weight	100.00
Net Content	100.00
Active Ingredients:	By Weight
1,1-Dimethyl-3-methyl-2-propyl-4-pyridinium chloride	1.0%
1,1-Dimethyl-3-methyl-2-propyl-4-pyridinium chloride	98.98%
Total	100.00

CAUTION
KEEP OUT OF THE REACH OF CHILDREN

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS.
 Irritant if swallowed or absorbed through the skin. Avoid contact with skin, eyes or clothing. In case of contact, flush with water. Do not contaminate food or feed.

ENVIRONMENTAL HAZARDS
 This product is toxic to fish. Do not apply directly to water. Do not contaminate water by direct application, cleaning of equipment, or disposal of wastes. Apply this product only as directed on this label.

(Pictures)

Apply Vertagreen Crabgrass Preventer Plus only once - in spring - stop crabgrass before it has a chance to start. One application prevents crabgrass all year long. Betan is a highly effective crabgrass controller which inhibits seed germination when applied in spring before temperature reaches 65-75°F.

Vertagreen Crabgrass Preventer Plus cleans your lawn, and will give a generous feeding of essential nutrients necessary to get your lawn off to a healthy start when applied in early spring.

In addition to a full feeding, an early application of Crabgrass Preventer Plus will work to keep weed seeds from germinating, thus providing your lawn with full scale protection for the entire growing season.

Net Contents _____ lbs.

EPA Reg. No. 3442-
EPA Est. 3442-

U.S. Agricultural Chemicals
 Division of U.S. Steel
 P. O. Box 1055
 Atlanta, Georgia 30301

DOCUMENT AVAILABLE

lb per 1000 sq ft

3442-716

Vertamech Herbicide - Preventer Plus Turf Food

GUARANTEED ANALYSIS

Total Nitrogen (N)	17.5%	Plant Food, derived from Ammoniated
Total Available Nitrogen	17.5%	Plant Food, derived from Ammoniated
Total Phosphorus (P ₂ O ₅)	7.0%	Plant Food, derived from Superphosphate
Total Potassium (K ₂ O)	17.5%	Plant Food, derived from Sulfate of Potash and Sulfate
Total Sulfur (S)	5.0%	Plant Food, derived from Sulfate of Potash and Sulfate
Total Calcium (Ca)	17.5%	Plant Food, derived from Calcium Carbonate
Total Magnesium (Mg)	5.0%	Plant Food, derived from Calcium Carbonate

Vertamech Herbicide - Preventer Plus Turf Food is an effective pre-emergence herbicide. It is easy to apply with standard seed and fertilizer spreaders. Use on lawns, lawns, parks and golf courses.

DIFFICULTY LABEL

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

A selective pre-emergence herbicide for the control of crabgrass, annual bluegrass (poa annua), poa annua, and other annual seed germinants in established stands of the following: Bluegrass (perennial), Bermuda grass, Ky. blue (perennial), centipede grass, fescue, St. Augustine grass, buff bermuda, zoysia grass.

USE TO PREVENT

For crabgrass control, apply in the spring before crabgrass germinates. Re-seeding should be delayed for 30 to 40 days after application. To insure satisfactory re-seeding results after a top-dress or overseed operation, the amount of seed should be increased. For poa annua (Annual bluegrass) control: Apply late summer to early fall (mid-July to mid-September) prior to annual bluegrass (poa annua) germination. Chemicals that effectively control poa annua seedlings often delayed germination of fine leaved species. Re-seeding should be delayed 12 to 15 weeks after application.

HOW TO APPLY

Apply this product evenly over the turf area without skips or misses. A more uniform application can be made by spreading half of the required amount of this product over the area and then applying the remaining half at right angles or crosswise. Both drop type and rotary type spreaders may be used to apply.

Listed below are suggested spreader settings. These settings are only approximately. Differences may occur due to condition of spreader, speed at which applied, and roughness of area to be treated. This broadcast rate is for approximately 5,000 square feet (example: 50 ft. x 100 ft. area) of spray nozzle. Calibrate the rate of the area to be treated and do not over or under apply. Close spreader when halting, stopping or turning. DO NOT RE-ENTER

SUGGESTED BROADCAST SETTINGS

Model	Setting
Centra	5-1/4" - 6-1/4"
Centra Broadcast	4-1/4" - 5-1/2"
Ex-100	3-3/4" - 4"
GreenView	5-1/4" - 6-1/4"
Ortho 3900	5-1/2" - 6"
Scotts Drop	5-3/8" - 5-1/4"
Scotts Broadcast	5-0"
Scotts 3600	5-0"
Scotts Broadcast	3-1/4" - 3-1/2"

SAFETY PRECAUTIONS

Storage: Keep container closed and away from children, foodstuff, or domestic water supplies. Empty container, clean spreader, and put in trash collection.

NOTICE

U.S. Government will not accept responsibility for injury to persons or animals or for loss or damage to property, or other property resulting from failure to follow directions for use on the above container printed on this product.

BEST DOCUMENT AVAILABLE