


Vertagreen 55% Malathion Concentrate - 1 Gallon

**50101
 1 of 8
 05/13/1988**

**Vertagreen
 55% MALATHION
 Concentrate
 Premium Grade**

**For Household Insects-Fleas on Dogs
 & Cats-Ornamental Pest Control-Fruit
 Insect Control-Vegetable Gardens - Citrus-
 Livestock Pests-Stored Grain-Peanuts**



ACTIVE INGREDIENT:	By Weight
Malathion O,O-dimethyl phosphorodithioate of diethyl mercaptosuccinate.....	55.00%
INERT INGREDIENTS:*	45.00%
TOTAL	100.00%

*Contains Xylene Range Aromatic Solvent.

Contains CYTHION® insecticide "THE PREMIUM GRADE

MALATHION™. Registered Trademarks American Cyanamid Company

EPA Reg. No. 3442-736

D M K N
 EPA Est. No. 3442-FL-2, 6720-FL-1, 10370-TX-1, 572-NJ-1
 Superscript used is the first letter of the lot number at bottom of the package.

**KEEP OUT OF REACH OF CHILDREN
 CAUTION**

Note. Regarding above copy, minimum type size is:
 18 Pt. for CAUTION
 12 Pt. for KEEP OUT OF REACH OF CHILDREN

See Back Panel for Additional Precautionary Statements

STATEMENT OF PRACTICAL TREATMENTS

IF SWALLOWED - Call a physician or Poison Control Center immediately. Induce vomiting by giving victim 1 or 2 glasses of water and touching back of throat with finger. Never give anything by mouth to an unconscious or convulsing person. This product contains an aromatic petroleum solvent. Treat symptomatically.

IF INHALED - Remove victim to fresh air at once. Apply respiration if indicated.

IF ON SKIN - Wash thoroughly with soap and warm water.

IF IN EYES - Flush with plenty of water. Get medical attention.

Not Reviewed. Registrant claims to be in accordance with
 Reregistration of Pesticide Product Guidance for Phase 2
 Response, Page 2.12, Active to Inert Change in Status.

298

LAROCHE INDUSTRIES INC.—

CONTAINER PRINT SPECIFICATION

Vertagreen 55% Malathion Concentrate - 1 Gallon

50101
2 of 8
05/13/1988

NOTE TO PHYSICIAN- Malathion, upon repeated use, may cause cholinesterase inhibition. Atropine is antidotal.

Net Contents 1 Gallon

Manufactured and Registered by
LaRoche Industries Inc.
1100 Johnson Ferry Road N.E. - Atlanta, GA 30342
Vertagreen is a Registered Trademark
4Y30

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

HOW TO MIX

INSECTS

AMOUNT CONC. PER QUART WATER

FLEAS -

Contact Spray	1 teaspoonful per quart
Residual Spray	3 teaspoonfuls per quart
On Dogs & Cats	1-1/2 teaspoonful per quart
Animal Quarters	3 teaspoonfuls per quart

LICE -

Residual Spray	3 teaspoonfuls per quart
----------------	--------------------------

BROWN DOG TICK -

Residual Spray	3 teaspoonfuls per quart
----------------	--------------------------

ANTS, CLOVER MITES, CRICKETS,

EARMIGS, PANTRY PESTS (Saw

Toothed Grain Beetle, Rice

Weevil, Dermestids, Indian

Meal Moth larvae), ROACHES,

SCORPIONS, SILVERFISH, SPIDERS

(3 Teaspoonfuls = 1 tablespoonful)

After mixing apply as a coarse wet mist. Thoroughly clean, sweep and vacuum areas possible before spraying. Remove dead insects afterwards.

IN THE HOME - For Fleas, Lice and Brown Dog Ticks. Thoroughly treat areas where insects may enter or congregate such as door sills, window frames, baseboards, closets, storage cabinets, openings around water pipes, in the home or other buildings. Treat pet sleeping quarters and adjacent areas of floor and carpeting. Replace animal quarters with fresh bedding after treatment.

Vertagreen 55% Malathion Concentrate - 1 Gallon

50101
3 of 8
05/13/1988

FOR BEDBUGS - Mix 3 teaspoonfuls in 1 quart of deodorized base oil (Kerosene). Apply lightly to mattress surfaces in sufficient quantities to mist the fabric and generously to beds and woodwork with special attention to wet all possible hiding places. Do not contaminate food utensils, milk, milk equipment nor water. Repeat applications as needed for insect control.

IN AND AROUND HOMES, DAIRIES FOOD PROCESSING PLANTS - For the insects as listed above in chart mix as indicated and apply to window frames, baseboards, under sinks, drainboards, stoves, cracks and crevices and other areas where insects may breed and hide. Repeat as needed for control.

FLEAS ON DOGS AND CATS - Mix as indicated above and thoroughly wet animals by a hand sprayer or sponge bathing. Animals that run outside may need to be retreated in 2 to 3 weeks. Examine for Fleas before re-treating. Do not treat animals under one month of age. Wear rubber gloves when treating animals.

FLEAS AND TICKS IN ANIMAL QUARTERS - Mix as indicated above and spray kennels, pens, yards, and under houses and porches to control Ticks and Fleas. Apply one quart ready to use spray to cover approximately 250 sq. ft. Treat runways, soil and grass thoroughly. Replace treated bedding with fresh bedding. Remove manure and refuse.

LIVESTOCK PEST CONTROL

BEEF CATTLE, GOATS, and NON-LACTATING DAIRY ANIMALS - For control of Lice, Ticks, and Horn Flies mix 1 gallon of this product in 100 gallons of water and using a power sprayer apply thoroughly to animals with enough pressure to part the hair. Small dilution rate; (6 1/2 ounces in 5 gallons of water). NOTE - Do not treat during lactation or within 2 weeks of freshening. Do not apply to calves under one month of age. Repeat treatment if needed at two week intervals.

SWINE - For Lice mix 1 gallon of this product in 100 gallons of water (6 1/2 ounces per 5 gallons of water). Apply with a power sprayer as a complete spray to animals, pens, and litter. One treatment may be sufficient. Repeat only as needed. Do not contaminate feed, food containers nor watering troughs. Treated animals should be kept out of sun and "weather" for a few hours after treatment.

STORED GRAIN

INSECTS CONTROLLED - Cereal Leaf Beetle, Confused Flour Beetle, Flat Grain, Beetle, Granary Weevil, Indian Meal Moth, Lesser Grain Beetle, Red Flour Beetle, Rice Weevil, Rusty Grain Beetle, Sawtoothed Grain Beetle.

PROTECTION OF STORED GRAIN - Wheat, Oats, Rice, Corn, Rye, Barley, and Grain Sorghum; Mix 1 pint of this product in 2 to 5 gallons of water to spray treat each 1,000 bushels of grain as it is being loaded or turned into final storage.

Vertagreen 55% Malathion Concentrate - 1 Gallon

50101
4 of 8
05/13/1988

PROTECTION OF FIELD OR GARDEN SEEDS - Mix 1/2 pint of this product in 1 to 2½ gallons of water to spray treat 500 bushels of seed as it is being turned into final storage.

SURFACE TREATMENT FOR INDIAN MEAL MOTH - Mix 1/2 pint of this product in 1 to 2 gallons of water to spray each 1,000 sq. ft. of grain surface after the bin is filled. Repeat within 2 weeks if needed.

EMPTY BINS - Thoroughly clean inside and out by sweeping up old grain and removing refuse and debris. Mix 1 gallon of this product in 25 gallons of water. Spray walls, ceilings, floors, augers, & machinery by forcing spray into cracks and crevices where insects may hide. This applies also to truck beds, boxcars, and ships' holds. Spray to the point of run-off, inside and out.

PROTECTION OF STORED PEANUTS - For list of insects see paragraph above under **INSECTS CONTROLLED**.

Residual Warehouse Spray - Before Storing Peanuts - Clean warehouse thoroughly of trash and old remains of peanuts 1 to 2 weeks before new peanut crop is stored. Thoroughly spray with sufficient pressure, interior of empty warehouse (including cracks and protected places), outside walls to height of 6 to 8 feet and the ground to a distance of about 6 feet from warehouse by mixing 1 pint of this product in sufficient amount of water to make 2 1/2 gallons of spray or mix 1 gallon in 19 gallons of water. Apply finished spray at the rate of 2 gallons per 1,000 sq. ft. or to point of run-off.

Bulk Spray Treatment - Peanuts Going into Storage; Apply 1½ pints of this product in 5 gallons of water per 15 tons of farmers' stock peanuts, as they go into storage. Apply coarse spray uniformly. Preferably use suitable mechanical spray applicator which regulates rate of application to flow of peanuts. Otherwise, adjust pressure and size of nozzle opening to rate of flow of peanuts. Shield the nozzles from wind and air currents.

SUPPLEMENTAL SURFACE SPRAY - After the warehouse has been filled level surface of peanuts before spraying. Refer to Directions for Use on 25% Wettable Powder Malathion product for dilution and spraying directions. This treatment should be used in conjunction with the regular spray of this Emulsifiable Concentrate as outlined on this label.

FRUIT INSECT CONTROL

APPLES - To control Aphids, Bud Moth, Forbes scale, Codling Moth, Plum Curculio, Red-Banded Leaf Roller, and Spider Mites, mix 2 tsp. in 1 gallon of water (1 1/2 pint per 100 gallons). Make two or more applications for mites and one or more for aphids. Do not apply later than 3 days before harvest. This product may injure Cortland and McIntosh varieties in summer sprays.

528

LAROCHE INDUSTRIES INC.—

CONTAINER PRINT SPECIFICATION

Vertagreen 55% Malathion Concentrate - 1 Gallon

**50101
3 of 8
05/13/1988**

GRAPES - For Leaf hoppers, Spider Mites, Mealybugs, use 1 tsp. in 1 gallon of water (1 pint in 100 gallons water). Do not apply later than 3 days before harvesting. May cause injury to foliage of certain varieties. Check with local Agricultural Authorities.

CHERRIES - For Black Cherry Aphids and Fruit Tree Leaf Rollers; mix 1 tsp. in 1 gallon of water and thoroughly spray trees. Do not apply later than 3 days before harvest. May injure foliage in some varieties.

PEARS - To control Codling Moth, Fruit Tree Leaf Roller, Pear Psylla, Plum Curculio, and Red Banded Leaf Roller use 2 tsp. in 1 gallon of water. Do not apply later than 1 day before harvest. May injure Bosc pears under certain conditions in the Northwest.

PEACHES - For Spider Mites, Oriental Fruit Moth, and Plum Curculio, mix 2 tsp. in 1 gallon of water. Spray when damage is first seen or is anticipated. Do not apply later than 7 days before harvest.

PLUMS & PRUNES - For Mealy Plum Aphids use 1 tsp. in 1 gallon of water. Treat when insects first appear. Do not apply later than 3 days before harvest.

STRAWBERRIES - For the control of Spider Mites, and Aphids use 1 tsp. per gallon of water (1 pint per 100 gallons water). Treat when insects first appear. Do not apply later than 3 days before harvest.

NOTE - Do not apply when fruit is in bloom and bees are present. Plan on regular spray applications to fruit at 1 to 2 week intervals through the growing season to get best insect control.

PECANS - To control Spider Mites, Aphids, Pecan Nut Casebearer and Pecan Phylloxera, dilute 2 pints in 100 gallons of water (2 tsp. per 1 gal.) and apply with a pressure sprayer. Treat when insects are first seen or their damage is anticipated. Follow a spray schedule.

CITRUS INSECT CONTROL

CITRUS - For use on Grapefruit, Kumquats, Lemons, Limes, Oranges, Tangerines, and Tangelos. For control of Black Scale, California Scale, Citricola Scale, Soft Scale, and Yellow Scale use 1 to 1-1/2 pints per 100 gallons of water. For control of Florida Red & Purple Scale and the Mediterranean Fruit Fly use 2 pints per 100 gallons of water. For Thrips and Aphids use 1-1/2 pints per 100 gallons. (This spray will also suppress Spider Mites that may be present).

Apply sprays to give thorough coverage of fruit and the upper and lower sides of leaves and twigs when insects are present or their damage is eminent. Do not apply when trees are in bloom or within 7 days of harvest.

Vertagreen 55% Malathion Concentrate - 1 Gallon

50101
6 of 8
05/13/1988**ORNAMENTAL PEST CONTROL**

Use on a wide range of plants and insects as listed, however do not apply to Boston, Pteris or Maidenhair Ferns, Viburnum, Lantana, Crassula, Canaerti junipers, petunias, or small leaf Spirea since injury may result. The proper use of a spreader sticker will enhance insect control, especially with scale. Treat when insects are first seen and damage is anticipated.

Use 1 pint per 100 gallons water (1 teaspoonful per gallon). For control of Lace Bugs and Oyster Shell Scale. Apply when Scale crawlers have settled on foliage.

Use 1-1/2 pint per 100 gallons water (2 teaspoonfuls per gallon). For Aphids, Spider Mites, Mealybugs, Whiteflies, European Pine Shoot Moth, Four-lined Leaf Bug, Japanese Beetles, Potato Leafhopper, Rose Leafhopper, Tarnished Plant Bug, Thrips, and Euonymus Scale.

Use 2 pints per 100 gallons water (2 teaspoonfuls per gallon). For Bagworms, Birch Leaf Miner, Boxwood Leaf Miner, Oak Kermes, Tent Caterpillar, Scale insects as: Azalea, Magnolia, Pine leaf scale, Fletchers, Florida Red Scale, and Juniper Scale.

Use 2-1/2 pints per 100 gallons water (2 teaspoonfuls per gallon). For Black Scale Crawlers, Monterey Pine Scale, and Soft Scale.

Use 4 pints per 100 gallons water (4 teaspoonfuls per gallon). For Pine Needle Scale, and Wax Scale. Apply in the spring when crawlers are active. Repeat 1 to 2 full-coverage sprays at 10 day intervals.

VEGETABLE GARDEN INSECTS


Thoroughly spray plants particularly the under sides of leaves and the developing fruit. Mix the per acre rate in 100 gallons and apply to one acre. Use approximately 1 gallon of the ready to use spray per 500 sq. ft. of garden plants for the small dilution rate. Treat plants when insects are first seen and damage is noticed. Do not spray later than days shown til harvest.

BEANS - For Aphids, Spider Mites, Cucumber Beetles, Leaf Hoppers;
Use 2 tsp. per gallon - 2 pints per acre - 1 day to harvest.

CELERY - For Aphids, Spider Mites; Use 2 tsp. per gallon - 1 1/2
pint per acre - 7 days to harvest.

CUCUMBER, SQUASH, MELONS - For Aphids, Spider Mites, Cucumber Beetle,
Leaf Hopper; Use 2 tsp. per gallon - 1 1/2 to 2 pints per acre - 1 day to
harvest. Do not apply unless plants are dry.

CABBAGE, BROCCOLI, KALE, MUSTARD, TURNIPS, HEAD LETTUCE - For Aphids,
Imported Cabbage Worm, Cabbage Looper, Leaf Hopper; Use
2 tsp. per gallon - 2 pints per acre - 7 days to harvest.



Vertagreen 55X Malathion Concentrate - 1 Gallon

50101
7 of 8
05/13/1988

EGGPLANT - For Aphids, Spider Mites; Use 1 tsp. per gallon - 1 pint per acre - 3 days to harvest.

ONIONS - For Thrips; Use 2 tsp. per gallon - 1 1/2 pint per acre - 3 days to harvest.

PEPPERS - For Aphids; Use 1 tsp. per gallon - 1 pint per acre - 3 days to harvest.

POTATOES - For Aphids, Leaf Hoppers, Mealybugs; Use 1 tsp. per gallon - 1 pint per acre - No waiting time til harvest.

RADISHES - For Aphids; Use 2 tsp. per gallon - 2 pints per acre - 7 days to harvest.

TOMATOES - For Aphids, Spider Mites; Use 2 tsp. per gallon - 1 1/2 pint per acre - 1 day to harvest.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Avoid breathing of spray mist or vapors. Avoid contact with skin. Wash thoroughly after handling and before eating or smoking. Do not contaminate feed or foodstuffs. Do not apply to animals or in dwellings.

RE-ENTRY AND WORKER SAFETY STATEMENT: Use only with adequate ventilation. After using this product indoors, ventilate thoroughly before occupying enclosed spaces. Do not allow contact with treated surface until sprays have dried. Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons. Do not enter treated areas without protective clothing until sprays have dried.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish, aquatic invertebrates, and aquatic life stages of amphibians. Do not apply directly to water or wetlands (i.e., swamps, bogs, marshes, and potholes). Drift and runoff may be hazardous to aquatic organisms in areas near the application site. Do not contaminate water when disposing of equipment washwaters.

This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

Vertagreen 55% Malathion Concentrate - 1 Gallon

50101
8 of 8
05/13/1988

PHYSICAL OR CHEMICAL HAZARDS

Do not use, pour, spill or store near heat or open flame.

STORAGE & DISPOSAL

(Note: Heading must be minimum 12 Pt. type size)


Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Store product in a cool, dry, locked place out of reach of children.

PESTICIDE

DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER

DISPOSAL: Glass - Triple rinse (or equivalent). Then dispose of in a sanitary landfill or by other approved State and local procedures. Plastic - Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned stay out of smoke. Household - Do not re-use empty bottle. Wrap bottle and put in trash.**NOTE:** The solvent used in this material may stain certain fabrics, plastics, rubber and asphalt materials such as tiles, carpets, and floor coverings. Treat them sparingly and clean them after the insecticide has dried. Do not treat areas frequented by children and do not allow them in treated areas until the spray has dried.**NOTICE:** LaRoche Industries Inc. can accept no responsibility for injury to persons or animals or loss or damage to turf, soil or other property resulting from failure to follow directions for use or to observe cautions printed on this package.UPC Code
25704-50101
Proof of Purchase

Below for Internal Use Only

APPROVALS - MARKETING

DATE

REGULATORY SRVS. DATE

BY