


Subject	Vertagreen Turf Fungicide - 8 oz.	Account: USS50031 Page 1 of 4 Date: Rev. 6/12/85
---------	-----------------------------------	--

Copy for Front



Vertagreen Turf Fungicide
Anti-Dione* + Thiram
EIGHT OUNCES WILL TREAT UP
TO 4,000 SQ. FT. OF TURF AREA

For Use on Golf Greens
and Fine Lawn Grasses

A Turf Fungicide Containing the Antibiotic
Acti-dione*

Controls Disease on Bentgrass, Bermudagrass, Kentucky
Bluegrass, Merion Bluegrass, St. Augustinegrass, and Fescue.

ACTIVE INGREDIENTS:	By Weight
Cycloheximide*.....	0.75%
3-[2-(3,5-dimethyl-2-oxocyclohexyl)-2-hydroxyethyl] glutarimide.	
Thiram -- tetramethyl-thiuram disulfide.....	75.00%
INERT INGREDIENTS:.....	24.25%
TOTAL:.....	100.00%

Acti-dione The Upjohn Company, Kalamazoo, Mich.

KEEP OUT OF REACH OF CHILDREN

WARNING


See other warnings on side panel

Note: Regarding above copy, minimum type size is:
18 Pt. for WARNING
12 Pt. for KEEP OUT OF REACH OF CHILDREN

EPA Reg. No. 3442-756

EPA Est. No.

Net Contents -- 8 Ozs.



Subject	Vertagreen Turf Fungicide - 8 oz.	Account: USS50031
		Page 2 of 4
		Date: Rev. 6/12/85

Copy for Front (Continued)

Manufactured For

USS Agri-Chemicals
Division of United States Steel
P. O. Box 1685 - Atlanta, GA 30301
Vertagreen is a Registered Trademark.

Copy for Side Panel

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
WARNING

May be fatal if swallowed. This preparation produces skin irritation. Do not get on skin, in eyes or on clothing. Wash thoroughly after using. Do not breathe dust or spray mist. Do not contaminate feed or feedstuffs. Consumption of alcoholic beverages increases the toxic effects of thiram. Use rubber gloves when mixing and spraying Vertagreen Turf Fungicide.

ENVIRONMENTAL HAZARDS

This pesticide product is toxic to fish. Do not contaminate water by cleaning of equipment or disposal of wastes. Drift or runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not apply directly to water or wetlands. Do not use on pasture grasses.

STATEMENT OF PRACTICAL TREATMENT

In case of contact, remove contaminated clothing and wash skin immediately with soap and water. For eyes, flush with water for at least 15 minutes and get medical attention.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

The contents of this package have been accurately weighed. Head space is provided to facilitate withdrawal and avoid loss due to spilling.

Do not rotate treated crop with crops other than those with registered thiram uses.

Do not reenter treated areas within 24 hours following application unless protective clothing is worn.

Subject	Vertagreen Turf Fungicide - 8 oz.	Account:	USS50031
		Page	3 of 4
		Date:	Rev. 6/12/85

Copy for Side Panel (Continued)

HOW TO APPLY

Vertagreen Turf Fungicide may be applied as a liquid spray with a hand or power sprayer, or a hose-attached sprayer. The solution should be sprayed the same day it is prepared. Do not "Water-in". When used in a preventive schedule, it should be applied before the earliest symptoms of the disease are expected to appear. Applications should be repeated at 7 to 10-day intervals to maintain satisfactory disease control. Curative control may be obtained by increasing the amount of Vertagreen Turf Fungicide and shortening the interval between applications. Curative applications should be made at the first sign of disease. Avoid mowing the grass for 24 hours following each application.

COMPATIBILITY: Vertagreen Turf Fungicide may be applied with most insecticides commonly used on grasses. Do not use or mix with lime or other alkaline materials.

AMOUNT TO USE

For Large Brown Patch, Dollar Spot, Pink Patch, Melting-Out, Fading-Out, Leaf Spot, Rust, Powdery Mildew, and Gray Leaf Spot. If possible, time applications to allow 12 hours minimum before rain or watering. The addition of Spreader-Sticker will improve control.

PREVENTIVE - Use 2 ounces (10 level tablespoonfuls) in 10 gallons of water and apply to 1,000 sq. ft. of turf area. During hot, humid conditions favorable for disease development, use 3 ounces in 10 gallons of water and apply to 1,000 sq. ft. of turf area.

CURATIVE - Use 3 to 4 ounces (15 to 20 level tablespoonfuls) in 10 gallons of water and apply to 1,000 sq. ft. of turf area. After control is obtained, the preventive schedule (see above) may be followed.

FOR SNOW MOLD

Spray Application - Use 6 to 8 ounces (30 to 40 level tablespoonfuls) in 10 gallons of water and apply to 1,000 sq. ft. of turf area immediately preceding the expected permanent snow cover. If possible, repeat winter applications following each thaw, using 3 ounces (15 level tablespoonfuls) per 1,000 sq. ft. of turf area.

Dry Application - Thoroughly mix 6 to 8 ounces (30 to 40 level tablespoonfuls) with 10 quarts of screened compost or sand, and spread evenly over 1,000 sq. ft. of turf area immediately preceding the expected permanent winter snow cover. If possible, repeat winter application following each thaw, using 3 ounces (15 level tablespoonfuls) per 1,000 sq. ft. of turf area. To obtain uniform coverage use a mechanical spreader and distribute half the required amount of product over the area and then the remaining half at right angles or crosswise.

Subject	Vertagreen Turf Fungicide - 8 oz.	Account: USS50031 Page 4 of 4 Date: Rev. 6/12/85
---------	-----------------------------------	---

Copy for Side Panel (Continued)

STORAGE AND DISPOSAL

(Note: Heading must be minimum 12 Pt. type size.)

STORAGE: Store in a dry place.

✓ DISPOSAL: Do not reuse empty container. Wrap container and put in trash.

NOTICE: USS Agri-Chemicals can accept no responsibility for injury to persons or animals or loss or damage to turf, soil or other property resulting from failure to follow directions for use or to observe cautions printed on this package.

Below for Internal Use Only

APPROVALS - MARKETING	DATE	REGULATORY SRVS.	DATE
BY <u>J. M. [Signature]</u>	<u>6/13/85</u>	<u>C. D. [Signature]</u>	<u>6-13-85</u>