




ow to use this  
sterilized soil.

To be effective  
moved in early  
in freshly culti-  
per 100 square  
firmly as possi-  
location. Do not  
to not cultivate  
A second late  
on gives year  
in the listed or-  
wers. See cau-  
NOT be treated  
ail, purslane,  
weed common  
as well as many  
badly treat weeds  
are not con-

der shade trees  
ubs and hedge  
such as ivy and  
on dichondra  
o treated areas

ly in rose beds  
antings  
r transplanting  
and initial weed-  
ly after seeding  
similar flowers.

**ES OF ORNA-  
FLOWERS CAN**

**GREEN TREES**  
rber vitae, Ash  
xwood, Cam-  
e. Ornamental  
za, Dogwood  
Fir, Forsythia  
Hydrangea  
niper, Lantana  
Maple, Mock-  
ak, Pachista-  
rpus, Poplar,  
ron, Rose, Rus-  
Sycamore, Tree  
ed privet, Vitur-  
ow, Yew.

Ivy, Boston Ivy,  
Parhysandra  
Ageratum, Alys-

Garden Mistle-  
cupressus, Mar-  
Mother of thyme,  
ing pinks, Nas-  
Petunia, Pink  
coneflower, Re-  
sage, Snapdra-  
flower, Sundro-  
Verbena, Virgin  
Wormwood, Zin-

**Strawberries:** (1)  
transplanting a  
square feet. (2)  
in the fall and ear-  
per 60 square ft.  
bloom.

**Vegetables:** App-  
product per 500  
square feet. E-  
brusser sprouts,  
iards garlic, ka-  
whole and cut-  
nip, yam — App-  
mediate after  
Apply one app-  
weeks after  
cucumber, he-  
watermelon —  
seeding. Egg p-  
promptly after  
transplants ap-  
planting.

**THE FOLLOW-  
JURED — DO N**  
rauga, button-  
germander, ge-  
thium, pansy,  
telanthera,  
anarys and h-

**PRECAUTIONS  
HAZARDS  
AND DANGERS**

Do not take into  
eyes. Wash after  
contaminated. If  
store near see.  
Do not graze or  
parts to livestock.

**ENVIRONMENTAL**  
Keep out of lake  
not contaminated  
equipment or

**Read ALL the Label FIRST!** To be effective existing weeds must be removed in early spring or treatment made on freshly cultivated ground. Apply 1 pound per 100 square feet S<sup>®</sup> to the ground as uniformly as possible. Water lightly after application. Do not treat dry, hard ground. Do not cultivate ground after application. A second late Summer or Fall application gives year around weed free pastures in the listed ornamentals, shrubs and flowers. See cautioning plants that should NOT be treated. Controls crabgrass, foxtail, purslane, lambsquarters, red root pigweed, common chickweed, spotted spurge, as well as many other annual grasses and broadleaf weeds. Ragweed and smartweed are not controlled.

May be used very effectively in rose beds and in iris and *Hydriolus* plantings

THE FOLLOWING SPECIES OF ORNAMENTALS, SHRUBS AND FLOWERS CAN BE TREATED

**GROUND COVERS:** Baltic Ivy, Boston Ivy, English Ivy, Moneywort, Parthenocissus, Wilson Ivy.

HERBACEOUS PLANTS: *Ageratum* Alys

**Strawberries:** (New Plantings) Treat after transplanting at the rate of 1/2 lb. per 60 square feet. Established plantings: Treat in the fall and early spring at the rate of 1/2 lb. per 60 square feet. Do not apply after first bloom.

**THE FOLLOWING PLANTS MAY BE INJURED — DO NOT USE NEAR** *bulb weeds*, *rajuga*, *button plums*, *carnation*, *chrysanthemum*, *germander*, *geum*, *ice plants*, *mesquero*, *theurim*, *pansy*, *phlox*, *sweet william*, *terranthera*, *verys*, *use ar*, and *the* *primary* and *herb* *is* *quant* *it*.

**CAUTION**

Do not take internally. Avoid contact with eyes. Wash after handling. Do not eat or contaminate feed or forage. Do not store near seeds that you might eat. Do not place treated grain or forage in parts to livestock.

pollution out of lakes, streams, rivers, ponds, and  
contaminate water by dumping, pouring, or  
placement or disposal of wastes.