221	Agri-unemicais	
	Division of United State	

1. 1. H 344) - 8/4 1. 25

C CONTAINER PRINT SPECIFICATIONS

lor 5

;		Account:	99033PL
	Vertagreen ProTurf Sprayable Herbicide - Pocket Label	Page	1 of 5
		Date:	Rev. 7/16/85
_1			

Copy for Top of Pocket Label

USS Vertagreen Sprayable Herbicide for Professional Turf with Teamtm II

Net Wt. 24 Lbs.

Covers Up to 5.33 Acres

DEC 2 0 1985

Under the Federal Insecticide. Fungicide, and Rodenticide Act, as amended, for the pesticide

registered unde EPA Rog. No

Copy for Left Column

A selective preemergence herbicide for the control of crabgrass, goosegrass and other annual weed grasses in lawns, golf courses and other turf grass areas in established:

Bentgrass (nonputting green) Bluegrass (perennial) Fescues (fine and tall) Ryegrass (perennial) Bahiagrass

Bermudagrass Centipedegrass St. Augustinegrass Zoysiagrass

ACTIVE INGREDIENTS: By	Weight
N-buty1-N-ethy1-a,a,a-trifluoro-2,6-dinitro-p-toluidine	14.00%
trifluralin: a,a,a-trifluoro-2,6,-dinitro-N,N-dipropy1-p-toluidine	14.00%
INERT INGREDIENTS:	72.00%
TOTAL:	100.00%

Contains 4.4 ounces active ingredient per pound of product.

*Teamtm - and the diagonal color bar are the registered trademarks of Elanco Products Company. ACCEPTEL

EPA Reg. No.

EPA Est. No.

KEEP OUT OF REACH OF CHILDREN

CAUTION

Note: Regarding above copy, minimum type size is: 18 Pt. for CAUTION 12 Pt. for KEEP OUT OF REACH OF CHILDREN

> See righthand column for additional precautionary statements.

Division of United State

Account:



Vertagreen ProTurf Sprayable Herbicide ~ Pocket Label Date:

Rev.7/16/85

99033PL

2 of 5

Copy for Left Column (Continued)

U_SS Agri-Chemicals Division of United States Steel P. O. Box 1685 - Atlanta, GA 30301 USS and Vertagreen are Registered Trademarks

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Read all Directions Carefully Before Applying

Vertagreen Sprayable Herbicide is a preemergence herbicide which is applied to established turfgrasses for the control of crabgrass, foxtails, goosegrass, annual bluegrass and other annual weed grasses. Vertagreen Sprayable Herbicide kills these weeds as they germinate. This product does not control emerged or established weeds.

Precautions: Vertagreen Sprayable Herbicide may thin established annual bluegrass (Poa annua) turfs. Vertagreen Sprayable Herbicide should not be applied in the spring to turfgrasses planted in previous fall unless the turf achieved full establishment and cover and was mowed at least 4 times during the fall. Vertagreen Sprayable Herbicide may thin winter grasses overseeded to dormant Bermudagrass areas if applied during the early spring.

Do not apply Vertagreen Sprayable Herbicide to golf course putting greens, Dichondra, newly sprigged areas of Bermudagrass, St. Augustinegrass or centipedegrass.

APPLICATION DIRECTIC:

Rates, Frequency and Timing of Application.

Apply Vertagreen Sprayable Herbicide one to two weeks prior to the germination of annual weed grasses.

Optimum weed control performance will be obtained if treated areas are irrigated soon after application.

Any cultural practices that disturb the soil such as aerification or verticutting, should be completed prior to application.



CONTAINER PRINT SPECIFICATIONS

•

1.1.

H		Account:	99033PL
Subje	Vertagreen ProTurf Sprayable Herbicide - Pocket Label	Page	3 of 5
Ň		Date:	-Rev.7716785

Copy for Right Column

Rates:

Cool Season Grasses 5.25 - 10.5 lbs/acre (1.9-3.8 ounces/1000 sq. ft.)

Warm Season Grasses 7-14 lbs/acre (2.6-5.2 ounces/1000 sq. ft.)

Within the listed rate ranges, use the lowest rate in areas with a short growing season or if only 6 to 8 weeks of control is desired. In areas with a long growing season or where season long control is desired, use rates at the upper end to the maximum rate in the range.

Do not reseed treated areas with desirable perennial grasses for a minimum of 42 days following treatment. Areas treated at the maximum rates should not be reseeded for a minimum of 90 days.

APPLICATION EQUIPMENT AND DIRECTIONS

Vertagreen Sprayable Herbicide should be mixed with water and applied as a spray. Apply in a minimum of 100 gallons of water per acre using any properly calibrated spray unit.

Vigorous, continuous agitation is required to keep Vertagreen Sprayable Herbicide in suspension. Sparger pipe agitation generally provides the best agitation.

When mixing Vertagreen Sprayable Herbicide with water, start by filling the spray tank 3/4 full of clean water. Start agitation and increase pump speed to operating pressure. Add the correct quantity of Vertagreen Sprayable Herbicide, continue agitation and finish filling the tank with water.

Tank-Mixing Instructions

Vertagreen Sprayable Herbicide is compatible with most commercially available turf grass pesticides and sprayable plant nutrients. Compatibility between specific products should be determined by mixing a small quantity of each additive in an appropriate quantity of water. Any changes in color, an abnormal precipitate, or rapid settling may indicate incompatibility.

When tank mixing several products always add to the spray tank in the following order:

Dry Flowables--Wettable/Dispersible Powders--Aqueous Suspensions--Flowables--Liquids--Solutions--Emulsifiable or Liquid Concentrates.

CONTAINER PRINT SPECIFICATIONS

ччгчсь С я С,

U U Vontagnoon DesTunf Sperieble Hautisid – Date is i	Account:	99033PL	
Vertagreen ProTurf Sprayable Herbicide - Pocket Label	Page	4 of 5	ļ
<u> </u>	Date:	-Rev.7/16/85-	-

Copy for Right Column (Continued)

Agri-Chemicals Division of United State

JUSS)

Always add materials to the spray tank while the agitation is operating and ensure that all products are completely dispersed in water.

Do not allow spray mixtures to stand overnight. If agitation must be stopped for any period of time, be certain that all material in the bottom of the tank is resuspended prior to resuming spray operations.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

Harmful if swallowed. Avoid contact with skin, eyes, or clothing.

Statements of Practical Treatment

In case of contact, immediately flush eyes or skin with plenty of water. If swallowed, drink large quantity of water and induce vomiting by placing finger in back of throat. Never give anything by mouth to an unconscious person.

If accidentally swallowed, or if irritation persists, get medical attention.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not contaminate any body of water by direct application, cleaning of equipment or disposal of waste.

STORAGE AND DISPOSAL (Heading must be minimum 12 pt. type size)

- <u>Storage</u>: Do not store in direct sunlight. Store in a cool dry place in original container only.
- Pesticide Disposal: Do not contaminate water, food, or feed by storage vr. a disposal. Wastes resulting from use may be disposed of on site to or at any approved waste disposal facility.

<u>Container Disposal</u>: Dispose of overpack in a sanitary landfill or incinerate, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Agri-Chemicels Division of United State wel

Ter C

CONTAINER PRINT SPECIFICATIONS

ect	·	Account;	99033PL
	ver owgreen frontin opragable nerbicide - rocket Laber	Page	5 of 5
		Date:	-Rev.7/16/85

Copy for Right Column (Continued)

NOTICE:

:

USS Agri-Chemicals can accept no responsibility for injury to persons or animals or for loss or damage to turf or other property resulting from failure to follow directions for use or to observe cautions printed on this package.

			Below For Intern		
APPROVALS	-	MARKETING	DATE	REGULATORY SRVS.	DATE
BY				CD DUMBUND	7-16-85

