PAI 23 3442-818

LAROCHE INDUSTRIES INC.

CONTAINER PRINT SPECIFICATIONS

ject	Vertagreen Weed & Feed 26-4-6	Account:	- V870200
Subj		Page Date:	17/1//60

Copy for Face:

A PREMIUM GRADE HERBICIDE AND TURE FOOD COMBINATION -- CONTROLS A WIDE SPECTRUM OF BROADLEAF WEEDS WHITE PROVIDING FAST AND LONG FEEDING ACTION. --

VERTAGREEN^E

(Pictorial)

26-4-6

Pictorials

3442.876

For use on Blue Grass, Fescue, Rye, Bermuda, Bahia, Ventipede, Zoysia and St. Augustine Lawns

Copy for Right Column:

KEEP OUT OF REACH OF CHILDREN

WARNING

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Get Medical Attention

IF ON SKIN:

Wash Skin with Soap and Water

IF IN EYES:

Flush Eyes with Water for 15 minutes

and Get Medical Attention

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

Note: Regarding above copy, minimum type size is:

18 Pt. for WARNING

12 Pt. for KEEP OUT OF REACH OF CHILDREN

Copy for Left Column

			۰
ACTIVE INGREDIENTS:	By Weight	•	•
Dimethylamine Salt of 2,4-dichlorophenoxyacetic acid **Dimethylamine Salt of 2-(2-methyl-4-chlorophenoxy)	0:313%	•	•
propionic Acid* ***Dimethylamine Salt of 2-(2,4-dichlorophenoxy)	0.314%	•••••	
propionic Acid	0.3102	••••	•
TOTAL:	99.063%	••	•
	*** :	••••	•

lift

CONTAINER PRINT SPECIFICATIONS

1000	Vertagreen Weed & Feed 26-4-6	Account: Page	7670200 2 of 6 12/17/86
1		Date:	

EQUIVALENT TO:

*0.26% of 2,4-Dichlorophenoxyacetic Acid

**0.26% of 2-(2-Methyl-4-Chlorophenoxy) propionic Acid

***0.26° of 2-(2,4-Dichlorophenoxy) propionic Acid

BY ISOMER SPECIFIC ASSAY

EPA Reg. No. 3442-878

A B C D
EPA EST. NO. 3442-IL-1, 3442-TX-2, 3442-NC-2, 3442-FL-2

Superscript used is the first letter of lot number at the top or the bag.

Net Wt. 20 Lbs.

Copy for Gussets - Vertagreen Weed & Feed 26-4-6

Copy for Upper 1/3 of Back:

NET WT. 20 LBS.

vertagreen^R

WEED & FEED 26-4-6

Copy for Upper 1/3 of Back (Continued)

GUARANTEED ANALYSIS

Total Nitrogen (N)	
Chlorine (C1), not more than 6.00-	
Sources of Plant Nutrients: Urea, Sulfur Coated Urea, Amm Phosphate, Muriate of Potash	nontum
Iron (Fe) 1.00% Manganese (Mn) 0.05% Zinc (Zn) 0.05%	
	•

^{*}Contains 4.0% Coated Slow Release Nitrogen derived from Sulfur Coated Urea.

CONTAINER PRINT SPECIFICATIONS

hject	Vertagreen Weed & Feed 26-4-6	Account: Page	V670200 3 of 6 12/17/86
S		Date:	1

Copy for Top of Back

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING

Avoid contact with skin, eyes or clothing. Avoid breathing dust. Wash thoroughly after handling. Do not get in eyes or on clothing. Caused substantial but temporary eye injury. Wear goggles or safety glasses.


ENVIRONMENTAL HAZARDS

Do not apply directly to water. Do not apply when weather conditions favor drift from target area. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Vertagreen Weed and Feed 26-4-6 is a stable mixture of three phenoxy herbicides. It is for use on Lawns, Turf (not pastures) and Golf Fairways, Aprons, Tees and Roughs.



346

CONTAINER PRINT SPECIFICATIONS

1	Vertagreen Weed & Feed 26-4-6	Account: Page	VG/0200 4 of 6 12/17/86
		Date:	<u></u>

Copy for Left Column: (Continued):

Broadleaf weeds controlled:

This product is designed for control of broadlear weeds in lawns such as:

Black Medic, Buckhorn, Burdock, Chickory, Chickweed, Clover, Beck, Dandelion, English Daisy, Ground Ivy, Heal-All, Knotweed, Lambsquarter, Mallow, Morning Glory, Pigweed, Plantain, Poison Ivy, Poison Oak, Purslane, Radweed, Red Clover, Red Sorrel, Stitchwort, Speedwell, Spurge, White Clover, Wild Carrot, Wild Garlic, Wild Onion, Wild Lettuce, Yarrow and other Broadleaf weeds.

Read the entire label before using this product.

This product may be used on Blue Grass, Fescue, Rye, Centipede, Bermuda, Bahia, Zoysia Lawns and St. Augustine grass. DO NOT USE on Dichondra and Carpet Grass. DO NOT USE this product where desirable clovers are present or allow the product to "wind drift" on flowers, vegetables, ornamentals, shrubs, trees or other desirable plants.

NORTHERN GRASSES:

WHEN TO APPLY: Best results are obtained when weeds are growing rapidly in spring or early fall. IMPORTANT: Apply when grass is moist (after rain, sprinkling or dew). For most effective broadleaf weed killing action, avoid unnecessary disturbances, including watering, on treated areas for 48 hours after application. Watering will wash off weed killing material. Do not reseed for at least 3 to 4 weeks after application. Do not apply to newly seeded or sodded lawns until after 3 mowings.

SOUTHERN GRASSES:

WHEN TO APPLY: Best results are obtained when weeds are growing rapidly in spring. IMPORTANT: Apply when grass is moist (after rain, sprinkling or dew). DO NOT water lawn after material is applied. Watering will wash off weed killing material. For most effective broadleaf weed killing action, avoid unnecessary disturbance of treated areas for 48 hours after application. Avoid contact with valuable shrubs, plants, or flowers. Possible retreation ment may be necessary if lawn is heavily overrun with weeds or if adverse low moisture conditions prevail and weeds are in a state of poor growth. Do not overapply. Under conditions of adequate moisture, high termperature and high humidity, over-application may cause minor transitory discoloration on St. Augustine.

NOTE: It is recommended that during hot, any conditions, application rates should be reduced 50% to avoid possible dehydration or browning of grass.

Do not reseed for at least 3 to 4 weeks after application. Do not apply to newly seeded or sodded lawns until after 3 mowings.

_/

CONTAINER PRINT SPECIFICATIONS

ect	l Vertagreen Weed & Feed 26-4-6	Account:	VG70200 5 of 6
ubje		Page	12/17/86
Su		Date:	

Copy for Left Panel Continued)

HOW TO APPLY: This 20 pound bag should be used on 5000 square feet. Apply evenly with a lawn spreader either as a one pass application or a two pass application, criss-crossing the area at half the rate each time. Lawns that are heavily overrun by weeds and that are in poor condition may require more than one application of this product. A suggested second application should be made in approximately 30 days.

The following settings are approximate for applying contents of this package at suggested rate of 4.0 pounds per 1,000 square feet. Settings are based on an average walking speed. Reduce setting for slower speed; raise setting for higher speed. Hand spreading is not recommended.

SUGGESTED SPREADER SETTINGS

MAKE	SETTINGS	
Kmart	6 - 7	
K Mart Proadcast	4 - 5	
Cyclore Broadcast	5-1/2 - 5-3/4	
Central	6	
Central Broadcast	4-1/2	
Precision	6	
Precision Broadcast	4 - 5	
Scotts	6	
Scotts Broadcast	D - £	
Ortho Broadcast	5 -6	
Ortho Drop	8 - 8-1/2	
Wards	6-3/4	
Wards Broadcast	5	
Sears	5-1/2 - 6	
Sears Broadcast	4-4-1/2	

	STORAGE AND DISPOSAL	
STORAGE:	(Note: Heading must be minimum 12 pt. type size) Store in a dry place. Do not contaminate water, food, or feed by storage or disposal. Do not store near other fertilizers or seeds.	• • • • • • • • • • • • • • • • • • • •
DISPOSAL:	Do not reuse empty bag. Wrap bag and discard in trash. ••••••••••••••••••••••••••••••••••••	•
NOTICE:	••••	•••••

LaRoche Industries Inc. can accept no responsibility for injury to persons or animals, or for loss or damage to turf, soil or other property resulting from failure to follow directions for use or to observe cautions printed on this package.

CONTAINER PRINT SPECIFICATIONS

0	Vertagreen Weed & Feed 26-4-6	Account	• VG70200 • 6 of 6
	.⊸i	Page	12/17/86
J	ans ans	Date:	

Copy for Back (Continued)

Manufactured and registered by:

taRoche Industries Inc. 1100 Johnson Ferry Road NE- Atlanta, GA 3034° Vertagreen is a Registered Trademark 4H86

UPC Code 25704-70200

STORE IN A DRY PLACE

Below for Internal Use Only				
APPROVALS	- MARKETING	DATE	REGULATORY SRVS.	DATE
ВҰ	allen J Foot	2/17/84	CD Surder	12.17.84

16