

Vertagreen for Professional Turf with TEAM[®]

99049
1 of 4
03/12/1987

Net Wt. 50 Lbs.

Copy for Left Column

| ACTIVE INGREDIENTS: | BY WEIGHT |
|--|---------------|
| N-butyl-N-ethyl-alpha, alpha, alpha trifluoro-2,6-dinitro-p-toluidine..... | 0.76% |
| Trifluralin: Alpha, alpha, alpha, trifluoro-2, 6-dinitro-N,N-dipropyl-O-toluidine..... | 0.38% |
| INERT INGREDIENTS:..... | <u>98.86%</u> |
| TOTAL:..... | 100.00% |

TEAM[®] is the Registered Trademark of Elanco Products Company, Benefin plus Trifluralin.

EPA Reg. No. 3442-XXX

| | | | | | |
|--------------|-----------|-----------|-----------|-----------|-----------|
| | A | B | C | D | H |
| EPA Est. No. | 3442-IL-1 | 3442-TX-2 | 3442-NC-2 | 3442-FL-2 | 3442-TX-1 |

Superscript used is the first letter of the lot number at the top of the bag.

GUARANTEED ANALYSIS

| | |
|---|--------|
| Total Nitrogen (N)..... | 30.00% |
| Available Phosphoric Acid (P ₂ O ₅)..... | 0.00% |
| Soluble Potash (K ₂ O)..... | 0.00% |

ACCEPTED

JUN 22 1987

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 3442-885

Source of Plant Nutrients: Urea

KEEP OUT OF REACH OF CHILDREN!
WARNING
STATEMENTS OF PRACTICAL TREATMENT

- IF IN EYES: Hold eyelids open and flush with water with a steady, gentle stream of water for 15 minutes. Call a physician.
- IF SWALLOWED: Call a physician or Poison Control Center immediately. Drink promptly a large quantity of milk, egg white, gelatin solution, or, if these are not available, large quantities of water. Avoid alcohol.
- IF ON SKIN: Wash with plenty of soap and water. Get medical attention if skin irritation persists.
- IF INHALED: If breathing difficulties occur, move individual to fresh air and get medical attention.

NOTE: Regarding above copy, minimum type size is:
18 Pt. for WARNING
12 Pt. for KEEP OUT OF REACH OF CHILDREN

BEST AVAILABLE COPY

2004

Vertagreen for Professional Turf with TEAM

99049
2 of 4
03/12/1987

Copy for Left Column (Continued)

PRECAUTIONARY STATEMENTS
WARNING
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Causes eye irritation. Do not get in eyes. Harmful if swallowed, inhaled, or absorbed through the skin. Avoid breathing dust and contact with skin or clothing. the granular formulation may cause skin sensitization reactions in certain individuals. Use eye protection and protective clothing such as coveralls, a long-sleeved shirt, and impermeable gloves when handling this product. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

ENVIRONMENTAL HAZARDS

Do not contaminate any body of water by direct application, cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL
(Note: Heading must be minimum 12 pt. type size.)

- STORAGE: Store in original container only. In case of leak or spill, contain material and dispose of waste.
- PESTICIDE Do not contaminate water, food or feed by storage or disposal. Wastes
- DISPOSAL: resulting from use of this product may be disposed of on site or at an approved waste disposal facility.
- CONTAINER Completely empty bag into application equipment. Then dispose of empty
- DISPOSAL: bag in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

WHERE TO APPLY

Apply to turf prior to germination of weed seeds since established weeds will not be controlled. Apply only to healthy established turf grass lawns.

IMPORTANT: Do not apply in the spring to Bermuda Grass which has been overseeded with winter grasses (annual rye) since thinning will occur. DO NOT use on Dichondra nor to newly sprigged areas of Bermudagrass, St. Augustine Grass or Centipede Grass until these turfs are well established, nor to Bentgrass or Putting Greens. DO NOT apply in spring to Turfgrass planted the previous fall. This product may thin established Annual Bluegrass (Poa Annua) & fine-leaved Fescue lawns when used at higher rates than 176 lbs. per acre (2.0 lb. A. I. per acre). Do not overdose.

WHEN AND HOW TO TIME THE APPLICATION

BEST AVAILABLE COPY

Do not apply to sidewalks, driveways, patios, or other concrete surfaces as staining may occur. Successful pre-emergence control of the annual grass weeds on this label requires proper timing of application. This product must be applied one to two weeks prior the germination of annual grass weeds. All summer annual grasses do not germinate at the same time. If application timing does not coincide with the normal germination period

99049

3 of 4

03/12/1987

Vertagreen for Professional Turf with TEAM®

Copy for Left Column (Continued)

of any of the annual grassy weeds listed on this label, weed control results may be erratic or poor.

WATER THE LAWN AFTER APPLICATION: It is not necessary but optimum weed control performance will be obtained if treated areas are irrigated with $\frac{1}{4}$ to $\frac{1}{2}$ inch water soon after application. Any cultural practices that disturb the soil, such as verticutting, should be done prior to application of this product.

Copy for Right Column

"SPRING GERMINATING" SUMMER ANNUAL GRASSES: CRABGRASS, FOXTAIL, BARNYARDGRASS

- A. Northern cool season grasses- For pre-emergence control, apply 132 to 176 lbs. per acre (equal to 1.5 to 2.00 lbs. A. I. per acre) applied in the spring one to two weeks prior to the onset of conditions favorable for annual weedy grass seed germination. One 50 lb. bag will cover 12,000 to 16,000 Sq. Ft. Use for 12,000 Sq. Ft. where turf is completely overrun with weeds. Use for 16,000 Sq. Ft. when weeds are sparse and the turf is in a Yearly Lawn Maintenance Program.
- B. Southern warm season turfgrass- For pre-emergence control apply 176 to 263 lbs. per acre (equal to 2 to 3 lbs. Active Ingredient per acre) in the late winter or early spring one to two weeks prior to the onset of conditions favorable for annual weedy grass seed germination. One 50 lb. bag will cover 8,000 to 12,000 Sq. Ft. Use for 8,000 Sq. Ft. where turf is completely overrun by weeds. Use for 12,000 Sq. Ft. when weeds are sparse or the turf is in a Yearly Crabgrass Control Program.

"LATE SPRING, EARLY SUMMER GERMINATING" ANNUAL GRASSES: GOOSEGRASS (SILVER CRAB)

Although classified as a summer annual grass, goosegrass normally germinates 4 to 6 weeks later than the crabgrass, foxtails and other summer annual grass species. Applications of this product should therefore be timed accordingly.

- A. Northern cool season turfgrasses- For pre-emergence control apply 176 lbs. per acre (equal to 2 lbs. Active Ingredient per acre) of this product in late spring or early summer ~~one to two weeks~~ prior to the expected germination period of goosegrass. Use 50 lbs. to cover 12,000 Sq. Ft.
- B. Southern warm season turfgrasses- For pre-emergence control, apply 176 to 263 lbs. per acre (equal to 2 to 3 lbs. Active Ingredient per acre) of this product in late spring or early summer one to two weeks prior to the expected germination of goosegrass. Use one 50 lb. bag to cover 8,000 to 12,000 Sq. Ft. Use 8,000 Sq. Ft. where turf is completely overrun by weeds. Use for 12,000 Sq. Ft. when weeds are sparse and the turf is in a Yearly Lawn Maintenance Program.

APPLY TWICE FOR FULL SEASON CONTROL IN NORTHERN AND SOUTHERN TURFGRASSES

BEST AVAILABLE COPY

In areas of the United States where pre-emergence control of both early germinating annual grasses (Crabgrass, Foxtail, and Barnyardgrass) and the later germinating Goosegrass is desired. Two applications of this product at the recommended rate spaced 8 to 10 weeks apart may be made to provide full season control of these annual grass species.

4064

Vertagreen for Professional Turf with TEAM[®]

99049
4 of 4
03/12/1987

Copy for Right Column (Continued)

POA ANNUAL (ANNUAL BLUEGRASS) CONTROL

In areas of heavy infestation of Annual Bluegrass, its elimination will leave areas of thin turf. Employ proper cultural practices to establish a new turf.

- A. Northern cool season turfgrasses- For pre-emergence control apply 176 lbs. per acre (equal to 2 lbs. Active Ingredient per acre) of this product per acre in late summer or early fall and again in late winter or spring one to two weeks prior to expected germination of annual bluegrass. One 50 lb. bag will treat 12,000 Sq. Ft. of turfgrass.
- B. Southern warm season turfgrasses- For pre-emergence control apply 263 lbs. per acre (equal to 3 lbs. Active Ingredient per acre) of this product per acre in the later summer or early fall one to two weeks prior to expected germination of poa annua. In southern areas of the United States where germination of annual bluegrass continues for extended periods of time, a second application at 263 lbs. per acre can be made 10 to 12 weeks after the initial treatment. One 50 lb. bag will treat and feed 8,000 Sq. Ft. of turfgrass.

RESEEDING TREATED AREAS: 1. Delay reseeding - 6 weeks when applying at the rate of 132 or 176 lbs. of product per acre (equal to 1.5 to 2.0 lbs. Active Ingredient per acre or when using 50 lbs. to cover 16,000 to 12,000 Sq. Ft.).

2. Delay reseeding - 12 to 16 weeks when applying at the rate of 263 lbs. of product per acre (equal to 3.0 lbs. of Active Ingredient per acre or when using 50 lbs. to cover 8,000 Sq. Ft. of turfgrass).

NOTICE: LaRoche Industries Inc. can accept no responsibility for injury to persons or animals or for loss or damage to turf, soil or other property resulting from failure to follow directions for use or to observe cautions printed on this package.

Manufactured and Registered by
LaRoche Industries Inc.
1100 Johnson Ferry Road, N. E.
Atlanta, Georgia 30342

BEST AVAILABLE COPY

Below for Internal Use Only

| APPROVALS- | MARKETING | DATE | REGULATORY SRVS. | DATE |
|------------|-----------|------|---------------------|----------------|
| BY | | | <i>CD Henderson</i> | <i>3-16-87</i> |