

ACCEPTED  
7-10  
UNDER THE FEDERAL  
FUNGICIDE AND RO  
FOR ECONOMIC POLI  
ED UNDER NO. 34

REC

**Pest**

**Beef Cattle**  
Horn Flies, Lice,  
Sarcoptic Mange, Ticks

**Swine**  
Lice, Sarcoptic Mange

**Sheep**  
Lice, Keds (Sheep Tick),  
Psoroptic or Sarcoptic Shee  
Scabs, Ticks

Horn Flies—spray back of a  
Lice, Ticks, Sarcoptic Man  
spray animals thoroughly. R

Do not treat sick animals. I  
6 months of age. Do not sp  
within 60 days of slaughter  
Animals should have acces  
in warm weather. Do not s

**NOTICE:** The manufacturer makes  
implied warranty concerning this n  
property damage, or other loss re  
The buyer assumes all risk and lia

**OBSERVE THE FOLLOWING USE PRECAUTIONS**

In order to protect fish and wildlife, do not contaminate lakes, streams, or ponds with this material.

The Federal Food and Drug Administration has set official tolerances for certain pesticide residues. In order that residues will not exceed established tolerances, use only at recommended rates.

**DIRECTIONS FOR USE**

**SPRAYING**


Pour recommended amount of this material into nearly filled spray tank. Add balance of water to fill tank. Keep agitator running during filling, mixing, and spraying operations. Do not allow mixture to stand. If mixture does not mix readily, but tends to separate as an oily layer, do not use as injury to animals may result. Do not combine with other materials.

**LIVESTOCK OILERS OR BACKRUBBERS**

For control of horn flies and as an aid to reduce lice infestations on beef cattle and swine, dilute this material at the rate of 1 gallon to 10 of diesel fuel. Place the mixture in oil reservoir or on burlap or oiler device and keep wet at all times. Place livestock oiler equipment where animals have free access to rubbing. Primarily near watering facilities or wherever animals congregate. Do not permit animals access to treatment within 30 days of slaughter or animals under 6 months of age.

**RESIDUAL SPRAY**

For control of House Flies, Mosquitoes and Stable Flies. Use 1 gal. of this material to 9 gals. of water. Apply spray at the rate of 1/2 to 1 gal. per 1000 sq. ft. of surface. Thoroughly treat exteriors of barns, stock sheds and other farm buildings as well as manure, vegetation and refuse. Do not use on dairy barns. Repeat as needed.



.....GAL. NET

**LIVESTOCK SPRAY, DIP & OILER CONC.**

For Use on Beef Cattle, Sheep and Swine. Controls Horn Flies, Lice, Sarcoptic and Psoroptic Mange and Ticks.  
Contains .18 Lbs. Lindane and 4 Lbs. Toxaphene Per Gallon .

**ACTIVE INGREDIENTS:**


Gamma isomer of benzene hexachloride (from Lindane).....	1.9%
Toxaphene (Technical chlorinated camphene)*.....	42.0%
Xylene range aromatic solvent .....	49.7%
<b>INERT INGREDIENTS:</b> .....	6.4%
*Chlorine Content 67% to 69%	
	100%

**WARNING**

**MAY BE FATAL IF SWALLOWED. MAY BE ABSORBED THROUGH SKIN. Do not breathe spray mist. Avoid contact with eyes, skin and clothing. Wash thoroughly after using. Do not contaminate feed or foodstuffs. In case of contact, wash immediately with soap and water. Do not use on household pets or humans.**

**Keep Out of Reach of Children  
DO NOT USE OR STORE NEAR HEAT OR OPEN FLAME  
Protect From Temperatures Below 0°F.**

Manufactured by  
**SCHALL CHEMICAL, INC. Monte Vista, Colorado**



GAL. NET

# LIVESTOCK SPRAY, DIP & OILER CONC.

For Use on Beef Cattle, Sheep and Swine. Controls Horn Flies, Lice, Sarcoptic and Psoroptic Mange and Ticks.

Contains .18 Lbs. Lindane and 4 Lbs. Toxaphene Per Gallon .

### ACTIVE INGREDIENTS:

Gamma isomer of benzene hexachloride (from Lindane)	1.9%
Toxaphene (Technical chlorinated camphene)*	42.0%
Xylene range aromatic solvent	49.7%
<b>INERT INGREDIENTS:</b>	6.4%

\*Chlorine Content 67% to 69%  
100%

## WARNING

MAY BE FATAL IF SWALLOWED. MAY BE ABSORBED THROUGH SKIN. Do not breathe spray mist. Avoid contact with eyes, skin and clothing. Wash thoroughly after using. Do not contaminate feed or foodstuffs. In case of contact, wash immediately with soap and water. Do not use on household pets or humans.

Keep Out of Reach of Children  
DO NOT USE OR STORE NEAR HEAT OR OPEN FLAME  
Protect From Temperatures Below 0°F.

Manufactured by  
**SCHALL CHEMICAL, INC.** Monte Vista, Colorado

**ACCEPTED**  
7-10-67  
UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENT CIDE ACT  
FOR ECONOMIC POISON REGISTER-  
ED UNDER NO. 3468-5

### RECOMMENDATIONS

Pest	50 Gal. of Water	5 Gal. of Water
<b>Beef Cattle</b>		
Horn Flies, Lice,	2 qt. for Sprays	1/2 cup
Sarcoptic Mange, Ticks	2 qt. for Dips	1/2 cup
<b>Swine</b>		
Lice, Sarcoptic Mange	2 qt. for Sprays	1/2 cup
	2 qt. for Dips	1/2 cup
<b>Sheep</b>		
Lice, Keds (Sheep Tick), Psoroptic or Sarcoptic Sheep Scabs, Ticks	2 qt. for Sprays	1/2 cup
	1 qt. for Dips	1/2 cup

### APPLICATION

Horn Flies -- spray back of animals every 3 weeks.  
Lice, Ticks, Sarcoptic Mange and Sarcoptic Sheep Scab. Immerse or spray animals thoroughly. Repeat after 2-3 weeks if needed.

### CAUTION

Do not treat sick animals. Do not use on dairy animals or animals under 6 months of age. Do not spray animals within 30 days of slaughter or dip within 60 days of slaughter. Do not treat emaciated or lactating animals. Animals should have access to the shade following dipping or spraying in warm weather. Do not spray when weather is excessively cold or hot.

**NOTICE:** The manufacturer makes no warranty of merchantability or any other express or implied warranty concerning this material. It shall not be held responsible for personal injury, property damage, or other loss resulting from the handling, storage or use of this material. The buyer assumes all risk and liability resulting from such handling, storage or use.

ACCEPTED WITH COMMENTS