

SUPERIOR 70 OIL

NET CONTENTS
Gallons

DIRECTIONS FOR MIXING

Fill the spray tank about one-third full of water, start the agitator and pour in the required amounts of Superior 70, Oil. If other spray materials are to be used with the oil, first mix the other materials thoroughly with the water in the spray tank, and then add the oil last as the filling of the tank is being completed.

DIRECTIONS FOR USE

GALLONS INDICATED ARE AMOUNTS TO USE PER 100 GALLONS OF WATER.

APPLES, PEARS: European

Fruit Lecanium Scale, San Jose Scale—Apply 2 gallons in the delayed dormant stage. Fruit Tree Leaf Roller, Scurfy Scale — Apply 3 gallons in the delayed dormant stage. European Red Mite—Apply 2 gallons in the green tip to delayed dormant stage. Pear Leaf Blister Mite—Apply 3 gallons in the dormant stage. Pear Psylla—Apply 2 gallons in the late dormant stage.

PEACHES: European Red Mite, San Jose Scale, European Fruit Lecanium—Use 2 gallons in the delayed dormant stage. Terrapin Scale—Use 3 gallons in the delayed dormant stage.

ORNAMENTALS and SHADE TREES (Conifers (see Note), Holly, Pyracantha, Sycamores, Hawthorne, Flowering Peach, Flowering Cherry, Apple, Magnolia, Maple, Dogwood, Oak, Privet, Birch): Soft Brown Scale, Oyster Shell Scale, Brown Apricot Scale, Black Scale, San Jose Scale, Red Spider Mites and Eggs—1 gal. to 100 gals. water. Dormant Spray on Deciduous Varieties—2 gals. per 100 gals. water. Dormant Spray on Other Varieties—1 gal. per 100 gals. water. NOTE: May cause discoloration of blue spruce. Do not use on conifers or ferns not tolerant to oil sprays.

INSECTICIDE



ACTIVE INGREDIENT:

Petroleum Oil (Superior Type)97.0%

INERT INGREDIENTS:3.0%

100.0%

Petroleum Oil Specifications:

Un sulfonated Residue ... (Min) 92%

Viscosity70 Seconds

CAUTION

KEEP OUT OF REACH OF CHILDREN.

Harmful if swallowed

DO NOT STORE AT TEMPERATURES BELOW 32° F.

DO NOT USE IN UNDILUTED FORM.

AVOID INHALATION OF VAPORS AND SPRAY MIST

NOTICE: Seller's guarantee shall be limited to the terms of the label, and subject there to the buyer assumes any risk to persons or property arising out of use or handling and accepts the product on these conditions.

Manufactured by

SCHALL CHEMICAL, INC. Monte Vista, Colorado

E. P. A. Reg. No. 3468-9

DIRECTIONS FOR USE

When a gun has been shut off or the outfit stopped for any length of time, spray on the ground for a few moments until the spray appears creamy white. This is desirable also where there are dead ends in the pipe lines of stationary outfits. All the material in stationary outfits should be sprayed out each day.

Spray on warm days in late Winter or early Spring, and before the buds separate into the clusters. Do not spray in freezing weather, or apply strong dosages after the buds have burst to any extent. Do not apply when buds have separated in the clusters.

Do not apply within 2 weeks of a sulfur spray or within 7 days of Captan.

Do not use if this spray oil does not emulsify. Correct emulsification is indicated if, when this product is added to the water in the spray tank, a uniform milk white spray mixture results provided the spray mixture is being agitated. If the spray mixture, after adding the oil to the spray tank with the agitation running, is not milk-white, do not use the resulting spray or the remainder of this product.

Dealers Should Sell in Original Packages Only.

DO NOT ALLOW THIS MATERIAL TO DRIFT ONTO NEIGHBORING CROPS OR NON-CROP AREAS OR USE IN A MANNER OR AT A TIME OTHER THAN IN ACCORDANCE WITH LABEL DIRECTIONS BECAUSE ANIMAL, PLANT OR CROP INJURY, ILLEGAL RESIDUES OR OTHER UNDESIRABLE RESULTS MAY OCCUR.

3468-9

2-4-72

1 of 2

EPAN

12472

ACCEPTED
FEB 07 1972
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO. 3468-9

2072